

A New Synthesis of Formycin *via* Nitropyrazole Derivatives

By J. GRANT BUCHANAN,* ALAN R. EDGAR, RODERICK J. HUTCHISON, ALAN STOBIE, and RICHARD H. WIGHTMAN
(Department of Chemistry, Heriot-Watt University, Riccarton, Currie, Edinburgh EH14 4AS)


Summary 3-(2,3,5-Tri-*O*-benzyl- β -D-ribofuranosyl)pyrazole (**4**) has been converted into formycin (**15**) by a sequence involving, as the key step, *cine* substitution, by cyanide ion, of the 1-nitro group in the 1,4-dinitropyrazole (**10**).

THE C-nucleoside antibiotic formycin^{1,2} (**15**) is a naturally occurring isomer of adenosine, containing a pyrazole in place of an imidazole ring. It has antiviral and anti-tumour properties and is able to replace the adenosine unit in a number of enzymic reactions at the nucleotide level. Although formycin has already been synthesised by two distinct routes^{3,4} we were interested in applying our own general method,⁵ which involves acetylenic intermediates, to the problem.

The ribofuranosylpyrazole^{6,7} (**4**) has been prepared by an improved method. Reaction of the ribofuranose (**1**) and 3,3-diethoxyprop-1-ynylmagnesium bromide formed a mixture of *D-altro* and *D-allo* diols (**2**) which underwent ring-closure with toluene-*p*-sulphonyl chloride in pyridine⁵ to give the β -D-*ribo*-isomer (**3**) [52% from (**1**)] together with the α -isomer (**8**). Acidic hydrolysis of (**3**) followed by reaction with hydrazine gave the pyrazole (**4**) (71%) identical with that from earlier preparations.

Although nitration of a pyrazole ring normally takes place at C-4,⁸ forcing conditions are required when a nitric-sulphuric acid mixture is used, owing to the formation of the pyrazolium cation. Acetyl nitrate, which acts on the neutral pyrazole ring, causes *N*-nitration. We therefore prepared the *N*-2,4-dinitrophenyl derivative (**5**) by means of 1-fluoro-2,4-dinitrobenzene and triethylamine in benzene solution.⁹ The benzyl ether groups were easily removed by boron trichloride and the resulting triol was converted into the triacetate (**6**) in 80% overall yield from (**4**). Nitration of (**6**) with cupric nitrate.3H₂O in acetic anhydride¹⁰ then gave the 4-nitro-derivative (**7**) (93%); the location of the nitro group was shown by the disappearance of the characteristic signal due to H-4 in the ¹H n.m.r. spectrum. Removal of the protecting groups⁹ with methanolic sodium methoxide yielded the triol (**8**) (76%) which was converted into the triacetate (**9**) (94%). *N*-Nitration¹¹ of (**9**) to give a single isomer (**10**) in 87% yield was achieved with the copper nitrate reagent; in the ¹H n.m.r. spectrum the signal for the pyrazole ring proton was deshielded further by the *N*-nitro group, thus establishing structure (**10**).

Habraken and Poels¹² have recently shown that 1,4-dinitropyrazoles react with secondary amines at C-5 in a *cine* substitution with expulsion of the *N*-nitro group as nitrite. When (**10**) was added to an excess of potassium cyanide in aqueous ethanol a very rapid reaction occurred at room temperature to give the key nitrile (**12**) (82%, as a crystalline benzene solvate). It may be noted, in addition, that thermal rearrangement¹¹ of (**10**) (in boiling anisole for 12 h) afforded the dinitropyrazole (**11**) (60%), a potential intermediate for elaboration of a fused pyrazine or other ring.



Catalytic hydrogenation of (12) gave the 4-aminopyrazole (13) (84%). Subsequent reaction of (13) with formamidine acetate in boiling 2-ethoxyethanol yielded formycin triacetate (14) (79%) which was converted by methanolysis into formycin (15) (90%), indistinguishable from an authentic sample.

The method described should be capable of extension to

the synthesis of a number of analogues of formycin containing modifications in the sugar and heterocyclic portions.

We thank the S.R.C. for studentships (to R. J. H. and A. S.)

(Received, 10th December 1979; Com. 1284.)

- ¹ R. J. Suhadolnik, 'Nucleoside Antibiotics,' Wiley-Interscience, New York, 1970.
- ² G. D. Daves and C. C. Cheng, *Progr. Medicin. Chem.*, 1976, **13**, 303.
- ³ E. M. Acton, K. J. Ryan, D. W. Henry, and L. Goodman, *Chem. Comm.*, 1971, 986; J. Farkaš and F. Šorm, *Coll. Czech. Chem. Comm.*, 1972, **37**, 2798; R. A. Long, A. F. Lewis, R. K. Robins, and L. B. Townsend, *J. Chem. Soc. (C)*, 1971, 2443.
- ⁴ L. Kalvoda, *Coll. Czech. Chem. Comm.*, 1978, **43**, 1431.
- ⁵ J. G. Buchanan, A. R. Edgar, and M. J. Power, *J.C.S. Perkin I*, 1974, 1943, and subsequent papers.
- ⁶ J. G. Buchanan, A. R. Edgar, M. J. Power, and G. C. Williams, *Carbohydrate Res.*, 1977, **55**, 225.
- ⁷ J. G. Buchanan, A. D. Dunn, A. R. Edgar, R. J. Hutchison, M. J. Power, and G. C. Williams, *J.C.S. Perkin I*, 1977, 1786.
- ⁸ K. Schofield, M. R. Grimmett, and B. R. T. Keene, 'The Azoles,' Cambridge University Press, 1976, p. 60ff.
- ⁹ J. G. Buchanan, M. E. Chacon-Fuertes, and R. H. Wightman, *J.C.S. Perkin I*, 1979, 244.
- ¹⁰ J. B. Menke, *Rec. Trav. chim.*, 1925, **44**, 141, 269.
- ¹¹ J. W. A. M. Janssen and C. L. Habraken, *J. Org. Chem.*, 1971, **36**, 3081, and subsequent papers.
- ¹² C. L. Habraken and E. K. Poels, *J. Org. Chem.*, 1977, **42**, 2893.