

Preparation of 1,5-Dienes by Photolysis of η^3 -Allylpalladium Complexes

By JACQUES MUZART and JEAN-PIERRE PETE

(Laboratoire de Photochimie, Equipe de Recherche Associée au C.N.R.S. No. 688, U.E.R. Sciences 51062 Reims Cedex, France)

Summary 1,5-Dienes are formed in good yields when dilute solutions of η^3 -allylpalladium complexes in aceto- nitrile are irradiated at 366 nm.

THE use of η^3 -allylpalladium complexes in organic synthesis has been extensively investigated over the past few years¹ However, there has been no report on the photochemical behaviour of these compounds From the photoreactivity of many organometallic complexes, it was expected that radical coupling reactions might be observed with η^3 -allylpalladium complexes We report here that 1,5-dienes are the main products formed when acetonitrile solutions of η^3 -allylpalladium complexes are irradiated

In a typical experiment a carefully deoxygenated solution of di- μ -chloro-bis[1-3- η -cycloheptylpalladium] (1) (150 mg) in MeCN (60 ml) was irradiated at 366 nm for 2 h Precipitation of palladium was observed and owing to the resulting filter effect, filtration through Florisil and re-irradiation were necessary to obtain a 52% conversion The 1,5-diene (7) (87% yield), and unchanged starting material were isolated from the reaction mixture Similarly the dienes (8)—(12) were produced from complexes (2)—(6) (Table).


Other functions, such as acetate or even a conjugated enone, do not interfere with the dimerization of the allylic group, which seems to be very regioselective These results contrast with those reported for the photolysis of nickel complexes Whereas the coupling of η^3 -allyl nickel

TABLE^a

Substrate	Conversion/%	Product	Isolated yield/%
(1)	52	(7)	87 ^b
(2)	100	(8)	55 ^c
(3)	78	(9)	32 ^d
(4)	80	(10)	80 ^{b,e}
(5)	62	(11)	78 ^{bf}
(6)	39	(12)	96 ^b

^a The starting complexes were prepared according to H C Volger, *Recl Trav Chim Pays-Bas*, 1969, **88**, 225 and B M Trost P E Strege, L Weber, T J Fullerton, and T J Dietsche, *J Am Chem Soc*, 1978, **100**, 3407 Irradiation was carried out using 2.5—6 mm l⁻¹ of substrate in MeCN under argon with light of wavelength λ 366 nm ^b Satisfactory i r, n m r, and mass or microanalytical data were obtained ^c Identified by comparison with authentic material (see P de Mayo, J B Stothers, and W Templeton, *Can J Chem*, 1961, **39**, 488) ^d Identified by comparison with literature data (see M Kobielski and H Pines, *J Am Chem Soc* 1957, **79** 5820) ^e Obtained as a mixture of 1,5-dienes with the head to head dimer predominating ^f The major product (62% yield) appears to result from head to head coupling

and hafnium⁴ suggests that homolytic cleavage of the carbon-palladium bond might be involved in the reaction of η^3 -allylpalladium complexes Further work, now in



complexes to give 1,5-dienes under thermal conditions in the presence of CO or phosphines is well known,² dimers are formed as only very minor products in the photolysis of these complexes³ The recent report on radicals and hexa-1,5-diene formation during the photolysis of tetra- (η^3 -allyl)-complexes of molybdenum, tungsten, zirconium,

progress, is required before the mechanism can be defined exactly.

We thank la Société Roure Bertrand Dupont for a gift of citronellol

(Received 17th December 1979, Com 1306.)

¹ B M Trost *Tetrahedron* 1977, **33**, 2615

² G Wilke, B Bogdanovic, P Hardt, P Heimbach, W Keim, M Kroner, W Oberkirch, K Tanaka, E Steinrucke, D Walter, and H Zimmermann, *Angew Chem Int Ed Engl*, 1966, **5**, 151, E J Corey, M F Semmelhack, and L S Hegedus, *J Am Chem Soc*, 1968, **90**, 2416

³ V A Kormer, L F Shelokhneva, and N A Kartsivadze, *Zh Obshch Khim*, 1974, **44**, 710

⁴ R Benn and G Wilke, *J Organomet Chem*, 1979, **174**, C 38