

Dimethyl- α -styrylsulphonium Bromide as a Reaction Intermediate


By YUAN L. CHOW,* BERT H. BAKKER, and KIYOSHI IWAI

(Department of Chemistry, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6)

Summary The addition product of styrene, bromine, and dimethyl sulphide was treated with bases to give the dimethyl- α -styrylsulphonium ion which reacted rapidly with various nucleophiles and also underwent slower base-catalysed rearrangement.

VINYLSULPHONIUM salts are versatile electrophilic alkylating agents that can serve in multistep reactions, and are generally prepared by alkylation of the corresponding alkyl vinyl sulphides.^{1,2} We report a simple method for the generation of a dimethylvinylsulphonium ion and its reactions.

Styrene reacted with a mixture of bromine and dimethyl sulphide in *ca.* 1 : 5 molar ratio in methylene chloride to give the crystalline bromosulphonium bromide (**1**) in 85% yield; m.p. 145—158 °C. The structure of (**1**) was supported by i.r., ¹H n.m.r. (ABX system at δ 4.24 and 5.12, $J = 8.5, 6.5,$ and 14.5 Hz, and singlets at δ 2.69 and 2.98), and ¹³C n.m.r. (t, δ 29.6; d, 62.05 p.p.m.) spectroscopy and analysis. The compound was obviously formed by the attack of the bromonium ion on the double bond followed by dimethyl sulphide as suggested by extensive skeletal rearrangements in a similar addition to norbornene.



Using ^1H n m r spectroscopy to monitor the reaction, treatment of (1) with potassium carbonate, triethylamine, or potassium hydroxide in deuterated solvents, such as D_2O , $(\text{CD}_3)_2\text{SO}$, or CD_3OD , rapidly gave the dimethyl- α -styrylsulphonium ion (2) which was gradually transformed into other compounds when left at room temperature. Much slower deuterium exchange of the methyl group was also observed. For example, in D_2O a clean spectrum of three singlets at 3.04, 6.40, and 7.55 (6 : 2 : 5 ratio, sodium 3-trimethylsilylpropanesulphonate as the internal standard)

was observed for (2). Brief heating of an aqueous solution of (1) containing 5–10% potassium carbonate gave a 1:1 mixture of styrene oxide and styrene glycol without a trace of other products. A similar reaction in the two phase carbon tetrachloride–aqueous system gave a nearly quantitative yield of styrene oxide. Under similar conditions the vinylsulphonium ion (2) reacted with CH_2XY (*e.g.* $\text{X} = \text{Y} = \text{CN}$, COMe , or CO_2Et , $\text{X} = \text{CN}$, $\text{Y} = \text{CO}_2\text{Et}$, $\text{X} = \text{COMe}$, $\text{Y} = \text{CO}_2\text{Et}$) smoothly at 50°C in several minutes, to give the corresponding cyclopropane derivatives (4).¹ Sodium cyanide also reacted rapidly at room temperature to give *cis*- and *trans*-cinnamitrile (3)² in 2:3 ratio. The reaction of (1) in aqueous methylamine gave aziridine (6).⁴ Most of these compounds are known compounds, in the present syntheses they are obtained cleanly in *ca.* 80% yields without tedious separation processes.

In methylene chloride, triethylamine reacts with (1) instantaneously at room temperature to give α -methylmercaptostyrene (5)⁵ without a trace of other products. In methanol–potassium carbonate, (2) reacted slowly to give (8) as the major product. The structure of (8) was supported by its n m r, i r, and mass spectra and analysis, in particular by the proton signals at δ 2.95 and 3.45 (A_2B_2 system) and singlets at δ 1.95, 3.22, and 3.62. While the formations of compounds other than (5) and (8) involve efficient straightforward attacks of nucleophiles, that of (8) follows the Sommelet–Hauser rearrangement⁶ *via* the ylide (7), the deprotonation step of which rearrangement is shown to be relatively slow. As a result, the crude product is contaminated by various by-products arising from nucleophilic substitutions, *e.g.*, 1,2-dimethoxyphenylethane, 1-methylthio-2-methoxy-1-phenylethane, and other unidentified compounds.

(Received, 3rd March 1980, Com 230)

¹ J. Gosselck, L. Béress, and H. Schenk, *Angew Chem, Int Ed Engl*, 1966, 5, 596

² J. Gosselck, L. Béress, H. Schenk, and G. Schmidt, *Angew Chem, Int Ed Engl*, 1965, 4, 1080, H. Braun and A. Amann, *ibid.*, 1975, 14, 755, 756, H. Braun, A. Amann, and M. Richter, *ibid.*, 1977, 16, 471, M. C. Caserio, R. E. Bratt, and R. J. Holland, *J. Am. Chem. Soc.* 1966, 88, 5747

³ B. Deschamps, G. Lefebvre, and J. Seyden-Penne, *Tetrahedron*, 1972, 28, 4209

⁴ A. P. Barsetti and D. R. Crist, *J. Heterocycl. Chem.*, 1975, 12, 1287

⁵ L. Léger and M. Saquet, *Bull. Soc. Chim. Fr.*, 1975, 657

⁶ K. W. Ratts and A. N. Yao, *J. Org. Chem.*, 1968, 33, 70, Y. Hayashi and R. Oda, *Tetrahedron Lett.*, 1968, 5381, S. E. Potter and J. O. Sutherland, *Chem. Commun.*, 1970, 520