

Unusual Effect of a Mixed Solvent on the Asymmetric Reduction of Chiral α -Keto-amides with Sodium Borohydride

Kenso Soai,* Kazuyuki Komiya, Yasuhiko Shigematsu, Hitoshi Hasegawa, and Atsuhiko Ookawa

Department of Applied Chemistry, Faculty of Science, Science University of Tokyo, Kagurazaka, Shinjuku-ku, Tokyo 162, Japan


The mixed solvent tetrahydrofuran + methanol (99 : 1) was much more effective than the individual solvents in the asymmetric reduction of chiral α -keto-amides with sodium borohydride; this solvent effect in asymmetric induction is unprecedented.

Despite the importance of the role of solvents in asymmetric synthesis, mixed solvents have been used very rarely.¹ We report the first example of a solvent effect on asymmetric induction in which a mixed solvent caused much higher asymmetric induction than either of the individual solvents.

In general, diastereoselectivities are low in the asymmetric reduction of chiral α -keto-esters or -amides with complex metal hydrides,² especially with sodium borohydride.

α -Keto-amides (**1a**, **b**),³ derived from (*S*)-proline methyl ester and α -keto-acids using dicyclohexylcarbodi-imide, were reduced with readily available NaBH₄ at 0 °C to afford (**2a**, **b**). Acidic hydrolysis of (**2a**, **b**) afforded the optically active α -hydroxy-acids (**3a**, **b**). When (**1a**) was reduced in tetrahydrofuran (THF) or methanol alone, the resulting enantiomeric excess (e.e.) of the mandelic acid (**3a**) produced was 36% (THF) and 4% (MeOH).[†]

A striking solvent effect was observed, however, when a THF + methanol (99:1 by volume) co-solvent was used.



The optical purity of the resulting (*S*)-(+)-mandelic acid $\{[\alpha]_D^{25} +101^\circ$ (c 1.9, H₂O) $\}$ increased to 64% e.e.[†] Using the same mixed solvent, (*S*)-(+)-lactic acid $\{[\alpha]_D^{25} -7.4^\circ$ (c 2.9, 1.5 M NaOH) $\}$ was obtained in 55% e.e.[‡] from (**1b**). Water, instead of MeOH, was also found to be effective.

In either mixed solvents, a catalytic amount of the protic solvent (MeOH or H₂O) in the aprotic THF was found to be essential for good asymmetric induction. The results are summarized in Table 1.

[†] Enantiomeric excesses were determined by ¹H-n.m.r. analysis of (**2a**) and/or by reported specific optical rotation of (**3a**). (*S*)-(+)-Mandelic acid, $[\alpha]_D +158^\circ$ (H₂O), S. Mitsui and A. Kanai, *Nippon Kagaku Zasshi*, 1965, **86**, 627.

[‡] Based on the value of (*S*)-(+)-lactic acid $[\alpha]_D^{20} -13.5^\circ$ (c 2.5, 1.5 M NaOH), 'Aldrich Catalog Handbook of Fine Chemicals,' Aldrich Chemical Co. Inc., Wisconsin, 1980.

Table 1. Effect of solvent on asymmetric reduction.

Ratio (v/v) THF:MeOH	(3a) (% e.e.)
100: 0	36
99: 1	64
0:100	4
THF:H ₂ O	
99: 1	50
85: 15	0

This work was partially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture.

Received, 9th August 1982; Com. 948

References

- 1 Y. Inouye, S. Inamasu, M. Horiike, M. Ohno, and H. M. Walborsky, *Tetrahedron*, 1968, **24**, 2907; T. Mukaiyama, K. Soai, T. Sato, H. Shimizu, and K. Suzuki, *J. Am. Chem. Soc.*, 1979, **101**, 1455.
- 2 J. D. Morrison and H. S. Mosher, 'Asymmetric Organic Reactions,' Prentice-Hall, Englewood-Cliffs, N.J., 1972, ch. 2. For the asymmetric hydrogenation or hydrosilylation of chiral α -keto-amides, see K. Harada, T. Munegumi, and S. Nomoto, *Tetrahedron Lett.*, 1981, **111**; I. Ojima, T. Tanaka, and T. Kogure, *Chem. Lett.*, 1981, 823.
- 3 Cf., B. W. Bycroft and G. R. Lee, *J. Chem. Soc., Chem. Commun.*, 1975, 988.