

A Maximum in a Grunwald–Winstein Plot for a Limiting S_N1 Solvolysis

Dennis N. Kevill and Mohd S. Bahari

Department of Chemistry, Northern Illinois University, DeKalb, Illinois 60115, U.S.A.

The Grunwald–Winstein Y values for acetone–methanol mixtures vary appreciably and in a uniform manner but the specific rates of solvolysis of 2-adamantyl perchlorate vary by less than threefold over a range of from 5 to 100% methanol content and a shallow maximum is observed.

Recently, it was reported¹ that the S_N2 solvolysis^{2,3} of methyl perchlorate in aqueous acetone or aqueous dioxan exhibits a maximum in the Grunwald–Winstein plot. Methyl derivatives

are good models for S_N2 solvolyses and, indeed, a Q -parameter scale^{4,5} has been developed for situating the solvolysis mechanism for alkyl derivatives within a spectrum intermediate

Table 1. Specific rates of solvolysis of *t*-butyl chloride^a and Grunwald-Winstein *Y* values for acetone-methanol mixtures at 25.0 °C.

MeOH (v/v)	10 ⁶ <i>k</i> ₁ (s ⁻¹) ^b	<i>Y</i>
100	7.53 ^c	-1.099
80	5.77 ± 0.15	-1.206
60	2.84 ± 0.06	-1.513
40	1.01 ± 0.02	-1.962
20	0.187 ± 0.012	-2.695

^a[Bu^tCl] = *ca.* 0.01 M. ^b With estimated standard deviations and all runs performed in duplicate. ^c From A. H. Fainberg and S. Winstein, *J. Am. Chem. Soc.*, 1956, **78**, 2770.

Table 2. Specific rates of solvolysis of 2-adamantyl perchlorate^a in hydroxylic solvents at 0.0 °C.

Solvent ^b	10 ⁶ <i>k</i> ₁ (s ⁻¹) ^c	<i>Y</i> ^d
CF ₃ CH ₂ OH ^e	610.3 ± 4.0	+1.045
MeOH	4.77 ± 0.14	-1.090
AcOH	(1.48) ^f	-1.675
EtOH	0.451 ± 0.011	-2.033
Pr ⁱ OH	0.123 ± 0.003	-2.73
Bu ^t OH	(0.071) ^g	-3.26

^a [2-AdOClO₃] = *ca.* 0.005 M. ^b On volume-volume basis, 98% of indicated solvent plus 2% pentane. ^c With estimated standard deviations and all runs performed in duplicate. ^d From P. R. Wells, *Chem. Rev.*, 1962, **62**, 171, except for value for CF₃CH₂OH which is from V. J. Shiner, Jr., W. Dowd, R. D. Fisher, S. R. Hartshorn, M. A. Kessick, L. Milakofsky, and M. W. Rapp, *J. Am. Chem. Soc.*, 1969, **91**, 4838. ^e 2% dioxan replacing 2% pentane. ^f Extrapolated value obtained from Arrhenius plot of four determinations at 17–42 °C. ^g Extrapolated value obtained from Arrhenius plot of three determinations at 26–50 °C.

between methyl derivatives and 2-adamantyl (2-Ad) derivatives, considered as limiting S_N1 substrates.⁶

Especially since 2-adamantyl methanesulphonate in aqueous acetone mixtures gives a shallow curve with gradually increasing *m* value,⁷ it is of interest to see whether perchlorate esters exhibit a maximum in Grunwald-Winstein plots for solvolyses at the S_N1 mechanistic extreme. The 2-adamantyl perchlorate⁸ was prepared by heterogenous reaction of 2-adamantyl bromide with silver perchlorate in pentane. Unfortunately, 2-adamantyl perchlorate (after transfer to dioxan) is of very limited solubility and very short half-life in solvents containing in excess of 50% water. However, it is soluble in all compositions of acetone-methanol mixtures, a system for which specific rates of solvolysis of methyl perchlorate have been reported.⁹


Methanol is considerably more polar than acetone as is indicated by higher *E*_T, *Z*, and log *k*_{ion} values¹⁰ and Grunwald-Winstein *Y* values for acetone-methanol mixtures (Table 1) show the expected trend of decreasing value with increasing acetone content. As with the acetone-water system, a Grunwald-Winstein plot of the methyl perchlorate data⁹ has a maximum, showing that the presence of water is not essential for this phenomenon to be observed.

The specific solvolysis rates (acid formation) for 2-adamantyl perchlorate in five hydroxylic solvents (Table 2) show the normal behaviour for a limiting S_N1 solvolysis with an *m* value of 0.989 ± 0.020 (standard error), inclusion of the *t*-butyl alcohol solvolysis (requiring a rather lengthy Arrhenius plot extrapolation) leads to a value of 0.941 ± 0.041. It is noteworthy that the acetolysis specific rate falls on the plot described by the alcoholyses; for 1-adamantyl toluene-*p*-sulphonate, the acetolysis rate¹¹ was faster than predicted based

Table 3. Specific rates of solvolysis^a of 2-adamantyl perchlorate^b in acetone-methanol mixtures.^c

Methanol % (v/v)	10 ⁶ <i>k</i> ₁ (25.0 °C)	10 ⁶ <i>k</i> ₁ (0.0 °C)
100	227 ± 4	4.77 ± 0.14
90		5.12 ± 0.12
80	299 ± 5	5.96 ± 0.19
60	345 ± 4	
40	317 ± 2	
20	236 ± 2	
10	171 ± 3	
5	138 ± 2	

^a In units of s⁻¹ and with estimated standard deviations; all runs performed in duplicate. ^b [2-AdOClO₃] = *ca.* 0.005 M. ^c On volume-volume basis, solvent is 98% of mixture indicated plus 2% pentane.

**Figure 1.** Solvolyses in acetone-methanol mixtures at 25.0 °C of methyl perchlorate (data from ref. 9), 2-adamantyl perchlorate, and *t*-butyl chloride.

upon the alcoholyses.¹² Presumably, the specific electrophilic assistance for the perchlorate leaving group is less than that for toluene-*p*-sulphonate and comparable to that for the chloride of the standard substrate.

Surprisingly, the specific rates of solvolysis of 2-adamantyl perchlorate in acetone-methanol mixtures (Table 3) show characteristics very similar to those for methyl perchlorate (Figure 1). In the range of 5 to 100% methanol, there is a less than three-fold variation in specific rate with a shallow maximum in plots against either solvent composition or *Y* values. The specific rates with 100 and 20% methanol content are essentially identical, despite a difference in *Y* value of 1.6.

We are currently investigating 2-adamantyl perchlorate in other mixed systems in an attempt to rationalize this highly unusual behaviour for a solvolysis involving a substrate of a

type universally accepted as a good model for limiting S_N1 character.

Received, 10th February 1982; Com. 141

References

- 1 D. N. Kevill and A. Wang, *J. Chem. Soc., Chem. Commun.*, 1981, 83.
 - 2 D. N. Kevill and H. R. Adolf, *Tetrahedron Lett.*, 1976, 4811.
 - 3 M. H. Abraham, A. Nasehzadeh, and J. J. Moura Ramos, *Tetrahedron Lett.*, 1981, **22**, 1929.
 - 4 F. L. Schadt, T. W. Bentley, and P.v.R. Schleyer, *J. Am. Chem. Soc.*, 1976, **98**, 7667.
 - 5 T. W. Bentley, C. T. Bowen, D. H. Morten, and P.v.R. Schleyer, *J. Am. Chem. Soc.*, 1981, **103**, 5466.
 - 6 P.v.R. Schleyer, J. L. Fry, L. K. M. Lam, and C. J. Lancelot, *J. Am. Chem. Soc.*, 1970, **92**, 2542.
 - 7 T. W. Bentley, C. T. Bowen, H. C. Brown, and F. J. Chloupek, *J. Org. Chem.*, 1981, **46**, 38.
 - 8 D. N. Kevill and B. W. Shen, *J. Am. Chem. Soc.*, 1981, **103**, 4515.
 - 9 J. Koskikallio, *Suom. Kemistil. B*, 1967, **40**, 131.
 - 10 For a discussion of these solvent polarity scales and the relevant values, see C. Reichardt, *Angew. Chem., Int. Ed. Engl.*, 1965, **4**, 29.
 - 11 P.v.R. Schleyer and R. D. Nicholas, *J. Am. Chem. Soc.*, 1961, **83**, 2700; M. L. Sinnott, H. J. Storesund, and M. C. Whiting, *Chem. Commun.*, 1969, 1000; B. R. Ree and J. C. Martin, *J. Am. Chem. Soc.*, 1970, **92**, 1660.
 - 12 D. N. Kevill, K. C. Kolwyck, and F. L. Weitzl, *J. Am. Chem. Soc.*, 1970, **92**, 7300.
-