

An Efficient Method for the Preparation of Carbapenem Derivatives with C-2 Carbon Side Chains from PS-5 and OA-6129 Compounds

Takeo Yoshioka,* Ken-ichi Yamamoto, Yasutaka Shimauchi, Yasuo Fukagawa, and Tomoyuki Ishikura


Sanraku-Ocean Co., Ltd., Central Research Laboratories, 9-1 Johnan 4-chome, Fujisawa 251, Japan

Treatment of *S*-oxides of PS-5 and OA-6129 esters with nitromethane gives 2-(nitromethyl)carbapenems which are converted into nitrile oxides leading to isoxazolyl, isoxazolinyll, or isoxadiazolinyll derivatives.

We have already established a new versatile method for displacement of the C-2 sulphur side chains of naturally-occurring carbapenems by a variety of thiols *via* the corresponding *S*-oxides.¹

Recently we have found that the sulphanyl group at the C-2 position can be replaced by active methylene compounds such as nitromethane or methyl cyanoacetate. When the *S*-oxide of PS-5 *p*-nitrobenzyl ester in dimethyl formamide (DMF) was treated ($-35\text{ }^{\circ}\text{C}$ for 1.5 h) with methyl cyanoacetate (10

equiv.) as a nucleophile in the presence of tetramethyl guanidine (TMG, 9 equiv.) as a base, a compound possessing the cyanoacetyl side chain at C-2 was obtained as a mixture of the geometrical isomers in 56% yield. The structure of the compound (1) was determined by mass spectrometry [m/z 413 (M)] and by observation of the C-3 proton [δ 5.75 and 5.48 (isomeric ratio 5 : 1) d, $J_{1\beta,3}$ 2.0 Hz] in the n.m.r. spectrum. The use of strong bases such as NaH, NaOMe, or KO^tBu in DMF resulted in the rupture of the β -lactam ring.



2-(Nitromethyl)carbapenems, (2) and (3),[†] were prepared in 58 and 54% isolated yields, respectively, by treatment (-25 °C, 30 min) of the sulphoxides of PS-5 and isopropylidene-OA-6129B2 *p*-nitrobenzyl esters² with nitromethane as a solvent in the presence of TMG. These nitromethyl compounds were so unstable towards acids that the purification could be achieved by extraction with organic solvents and column chromatography on Bio-Beads S-X3³ instead of silica gel.

[†] *Spectroscopic data:* (2) λ_{\max} (CH₂Cl₂), nm (ϵ) 268 (12 100); ν_{\max} (CHCl₃), cm⁻¹ 1780 (β -lactam), 1725 (ester), 1555, 1520, 1350 (nitro); ¹H n.m.r. (CDCl₃, Me₄Si) δ 3.02 (2H, d, *J* 9 Hz, 1-H₂), 3.22 (1H, m, 6-H), 4.02 (1H, dt, *J* 3, 9 Hz, 5-H), 5.31 (1H, d, *J* 15.5 Hz, CHH-NO₂), 5.78 (1H, d, *J* 15.5 Hz, CHH-NO₂). (3) λ_{\max} (CH₂Cl₂), nm (ϵ) 268 (13 100); ν_{\max} (CHCl₃), cm⁻¹ 1780 (β -lactam), 1720 (ester), 1555, 1520, 1350 (nitro); ¹H n.m.r. (CDCl₃) δ 2.98 (2H, d, *J* 9 Hz, 1-H₂), 3.30 (1H, m, 6-H), 3.85–4.35 (2H, m, 5-H, 8-H), 5.29 (1H, d, *J* 15 Hz, CHH-NO₂), 5.66 (1H, d, *J* 15 Hz, CHH-NO₂). (4) (purified by silica gel column chromatography) λ_{\max} (CH₂Cl₂), nm (ϵ) 324 (10 600), 267 (12 000); ν_{\max} (CHCl₃), cm⁻¹ 2295 (nitrile), 1792 (β -lactam), 1730 (ester); ¹H n.m.r. (CDCl₃) δ 2.98 (1H, dd, *J* 10, 19 Hz, 1-HH), 3.17 (1H, dd, *J* 10, 19 Hz, 1-HH), 3.28 (1H, m, 6-H), 4.08 (1H, dt, *J* 3, 10 Hz, 5-H). (5) λ_{\max} (THF), nm (ϵ) 325 (12 300), 269 (12 000), 262.5 (14 200), 257.5 (13 600), 251 (12 200); ν_{\max} (CHCl₃), cm⁻¹ 1785 (β -lactam), 1740 (ester); ¹H n.m.r. (CDCl₃) δ 3.0–3.8 (5H, m, 1-H₂, 6-H, 4'-H₂), 3.75 (3H, s, OMe), 3.93 (1H, dt, *J* 3, 9 Hz, 5-H), 5.10 (1H, m, 5'-H); *m/z* 443 (*M*). (6) λ_{\max} (THF), nm (ϵ) 268 (13 400), 260 (12 900), 253 (12 000), 248 (11 300); ν_{\max} (CHCl₃), cm⁻¹ 1790 (β -lactam), 1740 (ester); ¹H n.m.r. (CDCl₃) δ 3.27 (3H, m, 1-H₂, 6-H), 3.73 (3H, s, OMe), 3.97 (3H, s, OMe), 4.1 (1H, dt, *J* 3, 9 Hz, 5-H); *m/z* 499 (*M*). (7) λ_{\max} (THF), nm (ϵ) 270 (11 000), 264 (10 700), 261 (10 000), 255 (8800); ν_{\max} (CHCl₃), cm⁻¹ 1790 (β -lactam), 1735 (ester); ¹H n.m.r. (CDCl₃) δ 2.50 (3H, s, NMe), 3.16 (3H, m, 1-H₂, 6-H), 4.06 (1H, m, 5-H), 5.83 and 5.90 (1H, s respectively, 5'-H); *m/z* 476 (*M*).

Compound (2) was allowed to react (0 °C, 30 min) with methyl chloroformate (2 equiv.) in the presence of triethylamine (2 equiv.) in DMF, leading to a nitrile oxide (4)[†] which, without purification, was trapped with 1,3-dipolarophiles to afford isoxazolyl, isoxazolinyll, and isoxadiazolinyll derivatives. Namely, the cycloaddition reaction (room temp., 4 h) of (4) to methyl acrylate (3 equiv.) provided an isoxazoline compound (5) in 51% isolated yield from PS-5 ester sulphoxide. The use of dimethyl acetylenedicarboxylate (1.5 equiv.) or *N*-benzylidenemethylamine (4 equiv.) under the same reaction conditions gave the corresponding isoxazolyl or isoxadiazolinyll compound (6)[†] or (7)[†] in 36 or 34% isolated yield, respectively, from the sulphoxide.

These carbapenem derivatives having the carbon side chains at C-2 showed substantial antimicrobial activities against *Comamonas terrigena* B-996 and *Staphylococcus aureus* 209P on bioassay agar plates containing horse serum.

We thank Prof. Y. Yamada, Tokyo College of Pharmacy, for his helpful advice and discussions on this work.

Received, 13th August 1984; Com. 1172

References

- 1 K. Yamamoto, T. Yoshioka, Y. Kato, K. Isshiki, M. Nishino, F. Nakamura, Y. Shimauchi, and T. Ishikura, *J. Antibiot.*, 1983, **31**, 407.
- 2 T. Yoshioka, I. Kojima, K. Isshiki, A. Watanabe, Y. Shimauchi, M. Okabe, Y. Fukagawa, and T. Ishikura, *J. Antibiot.*, 1983, **36**, 1473.
- 3 Chromatography, Electrophoresis, Immunochemistry, and HPLC. Catalogue I, 1983, Bio-Rad Laboratories.