


Synthetic Use of Lithium Nitride, an Unusual Reducing Agent. Formation of Ti₄ and Ti₆ Complexes

Melvyn Kilner,* Gerard Parkin, and Andrew G. Talbot


Chemistry Department, The University of Durham, South Road, Durham DH1 3LE, U.K.

Lithium nitride acts as a reducing agent in co-ordinating solvents liberating dinitrogen, reducing TiCl₄ to LiTiCl₄(THF)₄ (THF = tetrahydrofuran), (η⁵-C₅H₅)₂TiCl₂ to [(η⁵-C₅H₅)₂TiCl]₂ and (C₅H₅)₆Ti₆(C₅H₄)₂N, and (η⁵-C₅H₅)TiCl₃ to (η⁵-C₅H₅)TiCl₂(THF), [(η⁵-C₅H₅)TiCl₂]_n, and (C₅H₅)₆(C₅H₄)₂Ti₆N₃; in the presence of CO, Li₃N reacts with (η⁵-C₅H₅)₂TiCl₂ (2 : 3 molar ratio) to form (η⁵-C₅H₅)₂Ti(CO)₂.

Except at temperatures close to its melting point (840–845 °C), lithium nitride is regarded as generally unreactive.¹ Water is unusual in that it reacts vigorously at ambient temperature to yield ammonia.^{2a} Lithium nitride has a hexagonal ionic structure³ and its poor solubility contributes much to its limited known chemistry.⁴ We have found that its solubility in tetrahydrofuran (THF), though low, is sufficient to enable it to react with halides of metals in their high oxidation states, and we report here a series of reduction reactions involving titanium(IV). Lithium nitride is not normally regarded as a reducing agent. We find it to be a useful synthetic reagent for converting Ti^{IV} into Ti^{III} and lower states in good yield. Lithium nitride reductions, which proceed according to equation (1), are often found to be preferable to other laboratory methods, in requiring non-forcing conditions, and in producing products easily separated from unwanted materials.




Titanium(III) chloride is usually prepared by hydrogen reduction of the tetrachloride^{2b} at elevated temperatures, though it can also be made using LiAlH₄,⁵ aluminium alkyls,^{2b} etc. An active form suitable for McMurry coupling reactions⁵ is formed when titanium(IV) chloride (99 mmol) in THF (40 cm³) is added to Li₃N (33 mmol) at –196 °C, and the mixture stirred at ambient temperature. The adduct TiCl₄(THF)₂,⁶ formed immediately, is replaced over 12 h by a pale green precipitate of LiTiCl₄(THF)₄ [equation (2)], which is recrystallised by Soxhlet extraction to produce pale blue-green microcrystals. The lithium cation may be replaced by tetra-alkylammonium and bis(triphenylphosphino)iminium cations in THF solution, and the corresponding bromides prepared by starting with TiBr₄. Authentic samples of the complexes were prepared also by the reaction of commercial titanium(III) halides with lithium halides in THF. Other solvates were prepared using other solvents such as dioxane or monoglyme. Further reactions of Li₃N with the titanium(III) compounds occur to produce grey Ti^{II} compounds, but these have not yet been fully characterised. Preliminary results⁷ on coupling of aryl ketones suggest that LiTiCl₄(THF)₄ can be used as a convenient and effective substitute for TiCl₃ in the conventional McMurry reactions.^{5,8}




[(η⁵-C₅H₅)₂TiCl]₂, (η⁵-C₅H₅)TiCl₂(THF), and [(η⁵-C₅H₅)TiCl₂]_n may all be prepared conveniently using Li₃N as the reducing agent. (η⁵-C₅H₅)₂TiCl₂ (4 mmol) and Li₃N (1.35 mmol) in THF (25 cm³) react at ambient temperature over 30 min to produce a green solution from which [(η⁵-C₅H₅)₂TiCl]₂ is isolated in 88% yield [equation (3)]. This preparative method has distinct advantages over the standard

method which uses zinc dust as the reductant, in that it avoids the formation of [(η⁵-C₅H₅)₂TiCl]₂ZnCl₂, which is often a major product. The method has advantages also over the reaction between (η⁵-C₅H₅)₂TiCl₂ and (η⁵-C₅H₅)₂TiBH₄ since it avoids the prior preparation of the latter complex.



With (η⁵-C₅H₅)TiCl₃ (2.4 mmol) in THF (35 cm³), Li₃N (0.81 mmol) reduction produces a green solution from which blue-green crystals of (η⁵-C₅H₅)TiCl₂(THF)⁹ are produced. Alternatively if toluene is added to the reaction mixture, and THF removed under reduced pressure, [(η⁵-C₅H₅)TiCl₂]_n is obtained by fractional crystallisation in 67% yield [equation (4)].



Further reduction of (η⁵-C₅H₅)TiCl₃ using Li₃N in a higher molar proportion (3Ti:2Li₃N) produces green (η⁵-C₅H₅)₄Ti₄Cl₄ (*m/z* 591). This diamagnetic complex has a simple ¹H n.m.r. spectrum consisting of a sharp singlet at δ 6.58 at ambient temperature, suggesting only one type of cyclopentadienyl group. Available evidence suggests a tetrahedral cluster with face bridging chloro-ligands as a probable structure {cf. [(cyclo-octatetraene)TiCl]₄¹⁰}.

The reactions so far described, which involve (η⁵-C₅H₅)TiCl₃ and Li₃N in 3:1 and 3:2 molar ratios, and (η⁵-C₅H₅)₂TiCl₂ and Li₃N in a 3:1 molar ratio, have demonstrated clearly that Li₃N acts as a reducing agent without incorporation of nitrogen into the products. Use of the appropriate molar ratio of Li₃N to remove all the chloride from the complexes would therefore be expected to produce the [C₅H₅Ti], or the titanocene [(C₅H₅)₂Ti] moieties and hence lead to known chemistry.^{2b} However reaction between (η⁵-C₅H₅)₂TiCl₂ (4.0 mmol) and Li₃N (2.8 mmol) in THF produced a very air-sensitive blue solid containing nitrogen. Mass spectral studies show the highest titanium-containing ion to be centred at *m/z* 820 corresponding to [(C₅H₅)₆(C₅H₄)₂Ti₆N]⁺. Fragmentation occurred by loss of nitrogen followed by sequential loss of cyclopentadienyl groups. A different blue complex is formed in the reaction between (η⁵-C₅H₅)TiCl₃ (2.3 mmol) and Li₃N (2.3 mmol) in THF (25 cm³). The ion of highest mass number observed in the mass spectrum occurred at *m/z* 848, corresponding to [(C₅H₅)₆(C₅H₄)₂Ti₆N₃]⁺, the fragmentation pattern being similar to that of the previous blue complex.

Attempts to obtain derivatives of the blue complexes were unsuccessful, the complexes failing to react with 2,2'-bipyridine, triphenylphosphine, 1,2-bisdiphenylphosphinoethane (dppe), and carbon monoxide under a variety of

conditions. Also the reactions between $(\eta^5\text{-C}_5\text{H}_5)_2\text{TiCl}_2$ or $(\eta^5\text{-C}_5\text{H}_5)\text{TiCl}_3$ and Li_3N in the presence of PPh_3 or dppe resulted in no change in the course of the reaction. However when the reaction between $(\eta^5\text{-C}_5\text{H}_5)_2\text{TiCl}_2$ and Li_3N was undertaken in the presence of carbon monoxide, $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO})_2$ ^{11,12} was obtained.

The synthetic utility of Li_3N as a reducing agent has been well demonstrated and exploratory work indicates that the reagent achieves a wide range of reductions of metals in high oxidation states.

We thank the University of Durham for research student-ships (to G. P. and A. G. T.).

Received, 17th February 1984;† Com. 211

References

- 1 'Gmelins Handbuch der Anorganischen Chemie. Lithium,' Verlag Chemie, Berlin, 1927, vol. 20, p. 86; 'Inorganic Syntheses,' ed. J. C. Bailar, McGraw-Hill, New York, 1953, vol. 4, p. 1.
- 2 (a) W. A. Hart and O. F. Beurnel, in 'Comprehensive Inorganic Chemistry,' eds. J. C. Bailar, H. J. Emeléus, R. Nyholm, and A. F. Trotman-Dickenson, Pergamon, Oxford, 1973, vol. 1, p. 357; (b) R. J. H. Clark, *ibid.*, vol. 3, p. 368.
- 3 E. Zintl and G. Brauer, *Z. Elektrochem.*, 1935, **41**, 102.
- 4 J. M. Morris, R. B. Dunmire, P. E. Koenig, and G. R. Newkome, *J. Org. Chem.*, 1972, **37**, 1244, and references therein; A. J. Gordon and R. L. E. Ehrenkauffer, *ibid.*, 1971, **36**, 44; C. J. Porritt, *Chem. Br.*, 1979, **15**, 282.
- 5 J. E. McMurry, *Acc. Chem. Res.*, 1983, **16**, 405; J. E. McMurry and M. P. Fleming, *J. Am. Chem. Soc.*, 1974, **96**, 4708.
- 6 P. M. Hamilton, R. McBeth, W. Bekebrede, and H. H. Sisler, *J. Am. Chem. Soc.*, 1953, **75**, 2881; B. Hessett and P. G. Perkins, *J. Chem. Soc. A*, 1970, 3229.
- 7 W. J. Feast and I. S. Millichamp, unpublished results.
- 8 W. J. Feast and I. S. Millichamp, *Polymer Commun.*, 1983, **24**, 102.
- 9 G. Fachinetti, G. Fochi, and C. Floriani, *J. Chem. Soc., Chem. Commun.*, 1976, 230.
- 10 H. R. van der Wal, F. Overzet, H. O. van Oven, J. I. de Boer, H. J. de Leifde Meijer, and F. Jellinek, *J. Organomet. Chem.*, 1975, **92**, 329.
- 11 B. Demerseman, G. Bouquet, and M. Bigorgne, *J. Organomet. Chem.*, 1975, **101**, C24.
- 12 L. P. Battaglia, M. Nardelli, C. Pelizzi, G. Predieri, and G. P. Chiusoli, *J. Organomet. Chem.*, 1983, **259**, 301.

† Received in revised form 9th October 1984.