## Corrigenda

## Diastereoselective Addition to an $\alpha$ -Alkoxyaldehyde under Dipolar (Cram–Felkin) and Chelation (Cram–Cyclic) Controlled Conditions; A Stereocontrolled Synthesis of (+)-Blastmycinone

Jun-ichi Uenishi, Hideo Tomozane, and Masatoshi Yamato

J. Chem. Soc., Chem. Commun., 1985, 717.

On p. 718 the structures of the chelation model (Scheme 1), compounds (4a,b) (Scheme 2), and compound (10) (Scheme 4) should be as below:

## The Nature of Hydrogen Bonding in Hydrotrioxides of Methyl $\alpha$ -Methylbenzyl Ether and $\alpha$ -Methylbenzyl Alcohol. Are Hydrotrioxides Self-associated in Solutions?

Božo Plesničar, Franci Kovač, Milan Hodošček, and Jože Koller

J. Chem. Soc., Chem. Commun., 1985, 515.

Dihedral angles,  $\Phi_1$  and  $\Phi_2$ , calculated for methyl hydrotrioxide (STO 4-31G), are defined as shown in the corrected Figure 1. H<sup>1</sup> is in the plane C–O<sup>1</sup>–O<sup>2</sup> (A); O<sup>1</sup>, O<sup>2</sup>, and O<sup>3</sup> are in the plane B; O<sup>2</sup>, O<sup>3</sup>, and H<sup>4</sup> are in the plane C. Bond lengths are in Å. [Figure 1 (HO<sub>3</sub>H) in ref. 1(c) should be corrected accordingly.]




Figure 1