

Facile Generation and Trapping of α -Oxo-*o*-quinodimethane; a Synthesis of 3-Aryl-3,4-dihydroisocoumarins


Satinder V. Kessar,* Paramjit Singh, and D. Venugopal

Department of Chemistry, Panjab University, Chandigarh-160014, India

Treatment of *o*-(trimethylsilylmethyl)benzoyl chloride (**1**) with fluoride ions in the presence of aromatic aldehydes and alkyl fumarates gives 3-aryl-3,4-dihydroisocoumarins (**3**) and substituted α -tetralones (**4**), respectively.

α -Oxo-*o*-quinodimethanes (**2**) are intermediates of considerable interest which on suitable [2 + 4] trapping give functionalised bicyclic systems. These intermediates (**2**) have been generated by thermolysis of benzocyclobutenones¹ and homophthalic anhydrides.² We have found that desilylation of *o*-(trimethylsilylmethyl)benzoyl chloride (**1**) provides a convenient route to them.³

Treatment of the acid chloride⁴ (**1**) with CsF in refluxing acetonitrile (or Bu₄NF in tetrahydrofuran or CH₂Cl₂) containing aromatic aldehydes furnished 3-aryl-3,4-dihydroisocoumarins.⁵ With benzaldehyde, (**3a**) was obtained [38%, m.p. 88–90 °C (lit.⁵ 87–89 °C)]; *p*-anisaldehyde, (**3b**) [51%, m.p. 108–109 °C (lit.⁵ 109 °C)]; veratraldehyde (**3c**) [62%, m.p. 101–102 °C, δ_{H} (CDCl₃) 3.19 (m, 2H, CH₂), 3.99



(s, 6H, 2 \times OMe), 5.55 (dd, 1H, CH), 7.35 (m, 6H, ArH), 8.25 (d, 1H, ArH)].

Under similar conditions, alkyl fumarates formed adducts (4) in moderate yield: with dimethyl fumarate, (4a) [48%, m.p. 87–88 °C (lit.¹, 86–88 °C)]; diethyl fumarate, (4b)†

† Obtained as a keto–enol mixture which on crystallisation from hexane gave the pure keto form.

[60%, m.p. 54–55 °C, δ_{H} (CCl₄) 1.15 (t, 3H, CO₂Et), 1.35 (t, 3H, CO₂Et), 3.24 (m, 2H, ArCH₂), 3.68–4.53 (m, 6H, 2 \times CO₂CH₂, COCHCO, COCH), 7.32 (m, 3H, ArH), 7.96 (d, 1H, ArH)]. When dimethyl acetylenedicarboxylate was employed trapping proceeded inefficiently (ca. 10% yield by n.m.r. spectroscopy). In desilylations carried out in the presence of methyl acrylate and acrylonitrile, or without any addend, the isocoumarin (6) was the predominant product [40%, m.p. 63–65 °C, δ_{H} (CCl₄) 2.48 (s, 3H, Me), 6.49 (s, 1H, =CH), 7.31 (m, 7H, ArH), 8.25 (d, 1H, ArH)]. Formation of (6) can be rationalised in terms of self trapping of (2) followed by a 1,5-hydrogen shift in the dimer (5).

The reactions were carried out in refluxing acetonitrile (4.5 h) using the acid chloride, CsF, and the dienophile in the molar ratio 1:1.2:3. The yields are for pure compounds isolated through chromatography and/or crystallisation.‡

Received, 17th May 1985; Com. 680

References

- 1 P. Schiess, M. Eberle, M. H. Francotte, and J. Wirz, *Tetrahedron Lett.*, 1984, **25**, 2201.
- 2 Y. Tamura, A. Wada, M. Sasho, and Y. Kita, *Tetrahedron Lett.*, 1981, **22**, 4283.
- 3 Generation of an imino analogue of (2) has been described recently; Y. Ito, E. Nakajo, and T. Saegusa, *Tetrahedron Lett.*, 1984, **25**, 5139. We are grateful to a referee for bringing to our notice a related publication on reaction of F⁻ with methyl-*o*-(trimethylsilylmethyl)benzoate; A. Aono, Y. Terao, and K. Achiwa, *Chem. Lett.*, 1985, **3**, 339. We have also observed that products derived from the intermediate (2) are formed on desilylation of corresponding esters and benzene sulphonic anhydrides.
- 4 W. R. Bergmark, M. Medaor, J. Issacs, and M. Thim, *Tetrahedron*, 1983, **39**, 1109.
- 5 For importance and methods for synthesis of this class of compounds see; M. Watanabe, M. Sahara, M. Kubo, S. Furukawa, R. J. Billedeau, and V. Snieckus, *J. Org. Chem.*, 1984, **49**, 742.

‡ All new compounds gave satisfactory analytical and spectral data.