

## The Very High Activity of $(\text{Ph}_2\text{P})_2\text{C}=\text{CH}_2$ to Michael Additions when complexed to Platinum(II) or Platinum(IV)

Xavier L. R. Fontaine, Fatma S. M. Hassan, Simon J. Higgins, Grant B. Jacobsen, Bernard L. Shaw,\* and Mark Thornton-Pett

School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.


Complexes of the types  $[\text{PtMe}_2(\text{vdpp})]$ ,  $[\text{PtMe}_3\text{I}(\text{vdpp})]$ , or  $[\text{PtX}_2(\text{vdpp})]$  ( $\text{X} = \text{Cl}$  or  $\text{I}$ ,  $\text{vdpp} = (\text{Ph}_2\text{P})_2\text{C}=\text{CH}_2$ ) undergo Michael additions with various amines, hydrazines, or carbon nucleophiles (ethyl acetoacetate or acetone) to the vinylidene double bond;  $[\text{PtMe}_3\text{I}(\text{Ph}_2\text{P})_2\text{CHCH}_2\text{NHCH}_2\text{Ph}]$  isomerizes to  $[\text{PtMe}_3\text{I}\{\text{Ph}_2\text{PCH}(\text{PPh}_2)\text{CHNHCH}_2\text{Ph}\}]$ .

We have reported previously that the complex  $[(\text{OC})_4\text{W}\{(\text{Ph}_2\text{P})_2\text{C}=\text{CH}_2\}]$  undergoes Michael type addition with a variety of amines, hydrazines, and carbon nucleophiles to give functionalized phosphine complexes whereas the free ligand,  $(\text{Ph}_2\text{P})_2\text{C}=\text{CH}_2$  ( $\text{vdpp}$ ), is inactive towards these additions.<sup>1,2</sup> We now report that  $\text{vdpp}$ , when complexed to  $\text{Pt}^{\text{II}}$  or  $\text{Pt}^{\text{IV}}$  is very highly activated towards Michael type additions and a variety of novel complexes† can be prepared.

Treatment of  $[\text{Me}_2\text{Pt}(\text{cyclo-octa-1,5-diene})]$  with  $\text{vdpp}$  gave  $[\text{Me}_2\text{Pt}(\text{vdpp})]$  (**1**), which in toluene solution reacted smoothly with  $\text{H}_2\text{NNH}_2 \cdot \text{H}_2\text{O}$  or  $\text{H}_2\text{NNHMe}$  to give (**2**),  $\text{R} = \text{H}$  or  $\text{Me}$ , in high (80–90%) yield. Treatment of (**1**) with an excess of allylamine at 20 °C gave, after a few minutes, a 60% yield of the adduct (**3**),  $\text{R}^1 = \text{H}$ ,  $\text{R}^2 = \text{CH}_2\text{CH}=\text{CH}_2$ . Other adducts of type (**3**) were prepared similarly using  $\text{NHMeCH}_2\text{CH}=\text{CH}_2$ ,  $\text{H}_2\text{NCH}_2\text{C}\equiv\text{CH}$ ,  $\text{NHMeCH}_2\text{C}\equiv\text{CH}$ ,  $\text{H}_2\text{NCH}_2\text{Ph}$ ,  $\text{H}_2\text{NCH}_2\text{C}_6\text{H}_4\text{OMe-4}$ , and the chiral amines (*R*)- and (*S*)- $\alpha$ -methylbenzylamine. The products of type (**3**) were

isolated in 55–90% yields and fully characterized. We expected that  $\text{Pt}^{\text{IV}}$  would activate  $\text{vdpp}$  towards nucleophilic attack more than  $\text{Pt}^{\text{II}}$ . Treatment of  $[\text{Me}_2\text{Pt}(\text{vdpp})]$  with an excess of  $\text{MeI}$  at *ca.* 20 °C gave the oxidative addition product  $[\text{Me}_3\text{IPt}(\text{vdpp})]$  (**4**). This  $\text{Pt}^{\text{IV}}$  complex (**4**) reacted with an excess of benzylamine at *ca.* 20 °C in  $\text{CDCl}_3$  to give a single, readily isolable, product  $[\text{Me}_3\text{IPt}\{(\text{Ph}_2\text{P})_2\text{CHCH}_2\text{NHCH}_2\text{Ph}\}]$  which we formulate, because of the subsequent reactions described below, with the stereochemistry (**5**),  $\text{R} = \text{CH}_2\text{Ph}$ , with the CH hydrogen and the iodine on opposite sides of the  $\text{P}_2\text{Pt}$  plane. Treatment of (**4**) with an excess of benzylamine in  $\text{CDCl}_3$  immediately led to complete conversion into (**5**) but then (**5**) gradually changed into two other similar products (A and B) each showing an AX  $^{31}\text{P}$ - $\{^1\text{H}\}$  n.m.r. pattern with satellites due to  $^{195}\text{Pt}$ . Gradually (over a period of 72 h at *ca.* 20 °C) (A) was converted completely into (B) and (B) was isolated, characterized, and its crystal structure (details to be published elsewhere) determined as (**6**). We suggest (A) has the structure (**7**) and the initial adduct has the structure (**5**). The isomerization (**5**)  $\rightarrow$  (**6**) is presumably promoted by the sterically hindered  $\text{Pt}^{\text{IV}}$  forming the less sterically demanding 5-membered ring with the 'harder' (N) donor atom.

† The complexes described in this Communication have been characterized by elemental analyses (C, H, N, halogen) and by  $^1\text{H}$ ,  $^1\text{H}$ - $\{^{31}\text{P}\}$ , and  $^{31}\text{P}$ - $\{^1\text{H}\}$  n.m.r. and i.r. spectroscopy; selected data were available to the referees.



We thought initially that  $\text{PtCl}_2$  would activate  $\text{vdpp}$  but that a nucleophile might, preferentially, displace the Cl rather than attack the vinylidene double bond. However, we have found this not to be the case. Thus, such is the remarkably activating effect of  $\text{PtCl}_2$  that a 0.02 M solution of  $[\text{Cl}_2\text{Pt}(\text{vdpp})]$  (8) in  $\text{CD}_2\text{Cl}_2$  at  $-80^\circ\text{C}$  reacts immediately and quantitatively with the equivalent amount of  $\text{H}_2\text{NCH}_2\text{C}_6\text{H}_4\text{OMe}$ -4 to give the adduct (9),  $\text{R} = \text{CH}_2\text{C}_6\text{H}_4\text{OMe}$ -4 [ $^{31}\text{P}$ - $^1\text{H}$ ] n.m.r. evidence]. The adduct (9) has been fully characterized.  $\text{H}_2\text{NNMe}_2$  immediately adds to (8) at  $-40^\circ\text{C}$  to give (9),  $\text{R} = \text{NMe}_2$ . Other amines have been added to (8) and their adducts of type (9) fully characterized, viz.  $\text{H}_2\text{NCH}_2\text{CH}=\text{CH}_2$ , the chiral amine (*R*)- $\alpha$ -methylbenzylamine, and  $\text{H}_2\text{NCH}_2\text{CO}_2\text{Et}$ . Also, and remarkably, ethyl acetoacetate and (8), both 0.012 M, in  $\text{CH}_2\text{Cl}_2$ , in the presence of anhydrous  $\text{Na}_2\text{CO}_3$ , gave the compound (10) in good (86%) yield. We find that although the dichloride (8) does not react with acetone, the corresponding di-iodide  $[\text{I}_2\text{Pt}(\text{vdpp})]$  reacts with acetone to give the C-alkylated product (11). This reaction is promoted by sodium iodide. Recently Schmidbauer<sup>3</sup> and co-workers have shown that methanol is added reversibly to  $\text{vdpp}$  in the 8-membered

ring  $[(\text{vdpp})\text{AuCl}]_2$  but  $\text{vdpp}$  complexes of Ag or Cu did not add methanol. We suggest that  $\text{vdpp}$ , when chelated to a single metal with a 4-membered ring, is more likely to undergo Michael type additions than in larger rings or in open-chain structures because the strain in the 4-membered chelate ring will be relieved somewhat as the central carbon goes from  $\text{sp}^2 \rightarrow \text{sp}^3$  hybridisation on Michael addition.

We thank the S.E.R.C. for support and Johnson Matthey PLC for the generous loan of platinum salts.

Received, 12th August 1985; Com. 1203

## References

- 1 G. R. Cooper, D. M. McEwan, and B. L. Shaw, *Inorg. Chim. Acta*, 1983, **76**, L165.
- 2 G. R. Cooper, F. Hassan, B. L. Shaw, and M. Thornton-Pett, *J. Chem. Soc., Chem. Commun.*, 1985, 614.
- 3 H. Schmidbauer, R. Herr, G. Müller, and J. Riede, *Organometallics*, 1985, **4**, 1208.