

## A Novel Route to Metallacyclopropene ( $\eta^2$ -Vinyl) Complexes from Alkynes: Synthesis and X-Ray Crystal Structure of $trans$ - $[\text{ReCl}\{\overline{\text{C}}(\text{CH}_2\text{Ph})\text{CH}_2\}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2][\text{BF}_4]$

Armando J. L. Pombeiro,<sup>\*a</sup> David L. Hughes,<sup>b</sup> Raymond L. Richards,<sup>b</sup> Jérôme Silvestre,<sup>c</sup> and Roald Hoffmann<sup>d</sup>

<sup>a</sup> Centro de Química Estrutural, Complexo I, Instituto Superior Técnico, Av. Rovisco Pais, 1096 Lisbon Codex, Portugal

<sup>b</sup> AFRC Unit of Nitrogen Fixation, University of Sussex, Brighton BN1 9RQ, U.K.

<sup>c</sup> Centre National de la Recherche Scientifique, Institut de Recherches sur la Catalyse, 2 Avenue Albert Einstein, 69626 Villeurbanne Cedex, France

<sup>d</sup> Department of Chemistry and Materials Science Centre, Cornell University, Ithaca, N.Y. 14853, U.S.A.


Treatment of the allene complex  $trans$ - $[\text{ReCl}(\overline{\text{H}_2\text{C}-\text{C}=\text{CHPh})(\text{dppe})_2]$  (dppe =  $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ ) with acids, HA (A =  $\text{BF}_4$  or  $\text{PF}_6$ ), in  $\text{CH}_2\text{Cl}_2$  or tetrahydrofuran gives the metallacyclopropene ( $\eta^2$ -vinyl) complex  $trans$ - $[\text{ReCl}\{\overline{\text{C}}(\text{CH}_2\text{Ph})\text{CH}_2\}(\text{dppe})_2][\text{A}]$  whose structure (A =  $\text{BF}_4$ ) has been authenticated by X-ray crystallography; the site of protonation is in agreement with extended Hückel calculations which show, in the HOMO, a high electronic density on the allene carbon atom which is not directly bonded to the metal.

There is increasing interest in the synthesis and reactions of compounds with metal-carbon multiple bonds, particularly when derived from alk-1-yne complexes whose conversion into vinylidene complexes is well documented.<sup>1</sup> Metallacyclopropene ( $\eta^2$ -vinyl) compounds,  $\overline{\text{M}}=\text{C}(\text{R})-\text{C} \lt$ , although few are known to date, are also products of metal-promoted reactions of alkynes, through intra- or inter-molecular nucleophilic attack at a ligating alkyne.<sup>2,3</sup> These compounds are possible reaction intermediates in insertion, oligomerisation, and co-cyclisation reactions of alkynes.<sup>2</sup> The possibility of their rearrangement into carbyne-type species has also been demonstrated.<sup>3</sup>

During our studies of the activation of nitrogenase substrates at electron-rich dinitrogen-binding metal centres, we have observed a novel route for the conversion of an alkyne into a metallacyclopropene ( $\eta^2$ -vinyl) species by electrophilic attack on an alkyne-derived allene ligand.

We have already reported<sup>4</sup> the metal-promoted 1,3-hydrogen migration of  $\text{PhC}\equiv\text{CMe}$  in its reaction with  $trans$ - $[\text{ReCl}(\text{N}_2)(\text{dppe})_2]$  (dppe =  $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$ ) to give the

allene complex  $trans$ - $[\text{ReCl}(\overline{\text{H}_2\text{C}-\text{C}=\text{CHPh})(\text{dppe})_2]$  (1), Scheme 1. A related rearrangement has been reported<sup>5</sup> for some areneacetylene complexes of chromium.



We now report that the allene ligand of (1), activated by the electron-rich {ReCl(dppe)<sub>2</sub>} centre, undergoes electrophilic attack by acids (HA) (A = BF<sub>4</sub> or PF<sub>6</sub>) in CH<sub>2</sub>Cl<sub>2</sub> or tetrahydrofuran (thf), at 20°C, to afford the first metallacyclopropene ( $\eta^2$ -vinyl) complex of rhenium, *trans*-[ReCl{=C(CH<sub>2</sub>Ph)CH<sub>2</sub>}(dppe)<sub>2</sub>][A] (2, A = BF<sub>4</sub> or PF<sub>6</sub>) as pink, diamagnetic crystals in high yield. In the <sup>13</sup>C{<sup>1</sup>H} n.m.r. spectrum (CD<sub>2</sub>Cl<sub>2</sub>, 298 K), the carbene carbon [Re=C(12) of Figure 1] resonance is observed at  $\delta$  258.2.


The molecular structure of (2, A = BF<sub>4</sub>) has been authenticated by X-ray crystallography and the complex cation is shown in Figure 1.<sup>†</sup>

The conformation of the rhenium complex cation (2) is very similar to that of the rhenium-allene complex (1),<sup>4</sup> but the dimensions of the ReC<sub>2</sub> ring system show a contrasting pattern of bonding. The Re—C(12) distance in (2) [1.947(6) Å] is much shorter than Re—C(11) [2.193(6) Å] and the corresponding distance in (1) [2.087(6) Å], and is indicative of an Re=C(12) double bond. The C—C bond lengths in the ligand and the location and satisfactory refinement of the hydrogen atoms on both the benzylic, C(13), and terminal, C(11), carbon atoms confirm the co-ordination bonding arrangements shown in Scheme 1.


The electronic factors driving the protonation of the allene complex were investigated by means of extended Hückel

<sup>†</sup> *Crystal data:* (2)·CH<sub>2</sub>Cl<sub>2</sub>, C<sub>62</sub>H<sub>59</sub>BCl<sub>3</sub>F<sub>4</sub>P<sub>4</sub>Re, *M* = 1307.4, monoclinic, space group P2<sub>1</sub>/n (equivalent to no. 14), *a* = 13.553(8), *b* = 19.014(4), *c* = 22.308(5) Å,  $\beta$  = 99.82(3), *U* = 5664.5 Å<sup>3</sup>, *Z* = 4, *D<sub>c</sub>* = 1.533 g cm<sup>-3</sup>;  $\lambda(\text{Mo-K}\alpha)$  = 0.71069 Å,  $\mu(\text{Mo-K}\alpha)$  = 24.8 cm<sup>-1</sup>. Diffraction intensities of a clear yellow square prismatic crystal, size 0.08 × 0.08 × 0.18 mm, were measured on an Enraf-Nonius CAD4 diffractometer with monochromated radiation. Lorentz-polarisation and absorption corrections were applied, and of 9961 independent reflections (to  $\theta = 25^\circ$ ), 7047 with *I* > 2 $\sigma$ (*I*) were considered observed. Structure determination was from Patterson, electron density, and difference Fourier maps. Preliminary refinement was in SHELX,<sup>6</sup> with most H-atoms riding (in idealized positions) on their bonded C-atoms. The four H-atoms on the {CH<sub>2</sub>-C-CH<sub>2</sub>} group were identified in difference maps and refined independently. The BF<sub>4</sub><sup>-</sup> ion and a molecule of solvent, CH<sub>2</sub>Cl<sub>2</sub>, are disordered, and the solvent molecule has not been fully resolved. In the final cycles of refinement by block-diagonal-matrix least-squares methods,<sup>7</sup> all the non-H atoms were refined anisotropically. Final *R* and *R<sub>w</sub>* values were 0.0761 and 0.0534 for the 9903 reflections with positive net intensities, each weighted *w* =  $\sigma_F^{-2}$ .

Atomic co-ordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1, 1986.




**Figure 1.** View of the Re-complex cation of (2), *trans*-[ReCl{=C(CH<sub>2</sub>Ph)CH<sub>2</sub>}(dppe)<sub>2</sub>]<sup>+</sup>, in the same projection as that shown for (1) in ref. 4. Selected dimensions (with e.s.d.s in parentheses): Re—C(11) 2.193(6), Re—C(12) 1.947(6), Re—P(2) 2.487(2), Re—P(3) 2.457(2), Re—P(4) 2.490(2), Re—P(5) 2.509(2), Re—Cl 2.440(2), C(11)—C(12) 1.412(9), C(12)—C(13) 1.500(8), C(13)—C(14) 1.511(9) Å; P(2)—Re—P(5) 164.5(1), P(3)—Re—P(4) 175.3(1), C(11)—C(12)—C(13) 130.0(6), C(12)—C(13)—C(14) 118.0(5)°.



**Figure 2.** A molecular orbital diagram for a model of the allene complex *trans*-[ReCl(H<sub>2</sub>C=C=CHPh)(dppe)<sub>2</sub>].

calculations.<sup>8</sup> Figure 2 is the molecular orbital diagram of the system before protonation. The interactions are typical of those for an alkene binding to a transition metal fragment.<sup>9</sup>

The point of interest is the composition of the HOMO. Although formally metal d-centered (*d<sub>xy</sub>*), this orbital turns out to be delocalised over the carbon framework of the allene unit with a substantial contribution (36%) from the *p<sub>y</sub>* orbital of C<sub>γ</sub>. The rest of the HOMO is made up of Re-*d<sub>xy</sub>*/*d<sub>yz</sub>* (54%) and C<sub>β</sub>-*p<sub>y</sub>* (10%). The high density on C<sub>γ</sub> can be explained using standard perturbation theoretical arguments: *d<sub>yz</sub>* (along



Scheme 2

with  $d_{xy}$ ) mixes with  $\pi_1$  to produce the HOMO, which is antibonding between Re and  $C_\beta$ . A crucial feature of this orbital is the subsequent mixing of  $\pi_2^*$  with it, which reduces the Re- $C_\beta$  antibonding character (dashed line in Figure 2). This requirement diminishes the contribution from  $C_\beta$  and reinforces that from  $C_\gamma$  as summarised in Scheme 2.

On the basis of simple frontier orbital concepts, a proton approaching the allene complex will protonate either the metal atom, which is co-ordinatively saturated, with rather diffuse electronic density, or the  $\gamma$  carbon atom with a highly directional and electron rich p orbital. The carbon atom is protonated. The protonation is predicted to be easier if  $d_{yz}$  is high in energy; that is, with a strong  $\pi$ -donor *trans* to the allene, as in the complex discussed here.

Calculations on (1) predict a shorter bond between Re and  $C_\beta$  than between Re and  $C_\alpha$ , in agreement with the structure. The overlap populations (computed with identical distances) are 0.373 and 0.286 electrons for Re- $C_\beta$  and Re- $C_\alpha$  respectively. From Figure 2 it can be seen that the partial double bond character of the Re- $C_\beta$  bond is a consequence of the second order mixing shown in Scheme 2; there is more Re- $C_\beta$  bonding character in the second M.O. (in-phase mixture of  $d_{yz}$  and  $\pi_1$ ) than there is antibonding character between these atoms in the HOMO.

The two main steps in the conversion of an alkyne into an  $\eta^2$ -vinyl ligand (alkyne isomerisation to a ligating allene<sup>4</sup> and its subsequent protonation) are induced by the relatively high electron-rich character and the  $\pi$ -electron releasing nature of the  $\{\text{ReCl}(\text{dppe})_2\}$  centre, and the tendency of rhenium to form multiple bonds to carbon.

Electrophilic attack at the  $\beta$ -carbon (relative to Re) of the allene ligand converts it from a two- into a three-electron donor, as also has been observed for ligating isocyanides<sup>10,11</sup> and alkyne-derived vinylidene ligands<sup>1</sup> at related metal centres, to give carbyne-type complexes.

Both alkynes and allenes are known substrates of nitro-genase<sup>12</sup> and although the mechanisms of their reductions (to alkenes, *etc.*) have not been ascertained,  $\eta^2$ -vinyl compounds of the type described in this work are conceivable intermediates in these reductions.

We thank I.N.I.C., The Calouste Gulbenkian Foundation (Portugal) and The British Council for financial support, Dr. A. G. Avent and Mr. C. J. Macdonald for recording the n.m.r. spectra, and Mr. A. Hills for assistance with the X-ray analysis.

Received, 26th March 1986; Com. 402

## References

- 1 For recent references, see, *e.g.*, K. R. Birdwhistell, T. L. Tonker, and J. L. Templeton, *J. Am. Chem. Soc.*, 1985, **107**, 4474; R. Weinand and H. Werner, *J. Chem. Soc., Chem. Commun.*, 1985, 1145; F. J. G. Alonso, A. Höhn, J. Wolf, H. Otto, and H. Werner, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 406; J. Silvestre and R. Hoffmann, *Helv. Chim. Acta*, 1985, **68**, 1461.
- 2 J. L. Davidson, M. Shiralio, L. M.-Muir, and K. W. Muir, *J. Chem. Soc., Dalton Trans.*, 1984, 2167, and references cited therein; L. Carlton, J. L. Davidson, J. C. Miller, and K. W. Muir, *J. Chem. Soc., Chem. Commun.*, 1984, 11; J. L. Davidson and L. Carlton, *ibid.*, 1984, 964; J. L. Davidson, W. F. Wilson, and K. W. Muir, *ibid.*, 1985, 460.
- 3 M. Green, N. C. Norman, and A. G. Orpen, *J. Am. Chem. Soc.*, 1981, **103**, 1267; S. R. Allen, M. Green, A. G. Orpen, and I. D. Williams, *J. Chem. Soc., Chem. Commun.*, 1982, 826.
- 4 D. L. Hughes, A. J. L. Pombeiro, C. J. Pickett, and R. L. Richards, *J. Chem. Soc., Chem. Commun.*, 1984, 992.
- 5 V. V. Krivykh, E. S. Taits, and M. I. Rybinskaya, 'XII International Conference Organometallic Chemistry,' Abstract No. 102, Vienna, 1985.
- 6 G. M. Sheldrick, SHELX-76. Program for Crystal Structure Determination, University of Cambridge, 1976.
- 7 J. D. Owen, BLOKLS, Block-diagonal least-squares refinement program, Rothamsted Experimental Station, 1982.
- 8 R. Hoffmann, *J. Chem. Phys.*, 1963, **39**, 397; R. Hoffmann and W. N. Lipscomb, *ibid.*, 1962, **36**, 3179. Calculations performed on a model system with all ligands taken as hydrogens. The parameters are from A. Dedieu, T. A. Albright, and R. Hoffmann, *J. Am. Chem. Soc.*, 1979, **101**, 3141.
- 9 See, *e.g.* T. A. Albright, J. K. Burdett, and M.-H. Whangbo, 'Orbital Interactions in Chemistry,' Wiley, New York, 1985.
- 10 A. J. L. Pombeiro, M. F. N. N. Carvalho, P. B. Hitchcock, and R. L. Richards, *J. Chem. Soc., Dalton Trans.*, 1981, 1629.
- 11 A. J. L. Pombeiro and R. L. Richards, *Transition Met. Chem. (Weinheim, Ger.)*, 1980, **5**, 55; J. Chatt, A. J. L. Pombeiro, and R. L. Richards, *J. Chem. Soc., Dalton Trans.*, 1980, 492; *J. Organomet. Chem.*, 1980, **184**, 357.
- 12 C. E. McKenna, T. Nakajima, J. B. Jones, C. Huang, M. C. McKenna, H. Eran, and A. Osumi, in 'Molybdenum Chemistry of Biological Significance,' eds. W. W. Newton and S. Otsuka, Plenum Press, London, 1980, p. 39.