

## Synthesis of Linear Coumarins *via para*-Claisen Rearrangement of Coumarate Ester Derivatives: Total Syntheses of Suberosin, Demethylsuberosin, and Ostruthin

Nicholas Cairns,<sup>a</sup> Laurence M. Harwood,<sup>\*a</sup> and David P. Astles<sup>b</sup>

<sup>a</sup> Dyson Perrins Laboratory, University of Oxford, South Parks Road, Oxford OX1 3QY, U.K.

<sup>b</sup> Shell, Sittingbourne Research Centre, Sittingbourne ME9 8AG, U.K.

Heating 2'-*O*-prenylated derivatives of 4'-*O*-methyl and 4'-*O*-benzyl methyl coumarates (**3**) furnishes the corresponding linear prenylated umbelliferones (**1**) directly *via* sequential *para*-Claisen rearrangement and relactonisation.

We have recently reported the regioselective Lewis acid catalysed *ortho*-Claisen rearrangement of 4'-allyloxycoumarate esters<sup>1</sup> and have successfully applied this approach to the synthesis of the naturally occurring linear coumarin demethylsuberosin (**1a**),<sup>2</sup> isolated from *Ruta graveolens*.<sup>3</sup> This procedure utilises 7-(1,1-dimethyl)allyloxycoumarin, derived initially from the readily available 3-chloro-3-methylbut-1-yne.<sup>4</sup> However, extension of this approach to the synthesis of geranyl and farnesyl prenylogues of (**1**) would necessitate the

use of the less readily obtained dehydrolinaloyl or dehydronerolidyl halides. Since the double inversion of a *para*-Claisen rearrangement would permit the use of ethers derived from more readily available prenyl, geranyl, and farnesyl halides, we have investigated the application of such an approach to the synthesis of structures (**1**).<sup>5</sup>

The 2'-*O*-allyl ethers (**4d-f**) were readily prepared from 7-methoxycoumarin (**2**, X = Me) *via* cleavage to the coumarate ester (**3**, X = Me) (NaOMe, MeOH, reflux, 92%)



Table 1.

Substrate	Product, % isolated yield		
	(1)	(5)	(6)
(4)			
d	80	12	0
g	78	10	0
j	4	7	84
k	6	8	76
l	28	9	53
m	27	8	50

from *Pastinaca ostruthium*<sup>8</sup>), and the farnesyl derivative (1c) in 84, 82, and 77% purified yields respectively.

Examination of the <sup>1</sup>H 300 MHz n.m.r. spectra of the geranyl and farnesyl *para* rearrangement products (1) and (5) indicated that the double inversion had regenerated the initial 2'-(*E*)-double bond geometry of the side chain, as evidenced by the absence of duplication of absorptions, particularly the doublet corresponding to the side chain benzylic CH<sub>2</sub> group.

The allyl ethers (4j), (4k) and crotyl ethers (4l), (4m) were also prepared in order to assess the effect of  $\gamma$ -substitution of the allyl ether upon the *para*-rearrangement process. The allyl ethers furnished largely the 8-substituted products (6j) and (6k) (Table 1), identified by a mutual *ortho* coupling (10 Hz) of the aromatic protons, whereas the crotyl ethers furnished a mixture of 8-(1'-methyl)allyl [(6l), (6m)] and 6-crotyl umbelliferones [(1l), (1m)], [(6):(1) ca. 2:1]. These results appear to reflect the degree of steric crowding at the benzylic position of the 8-substituted products. Interestingly, the yield of

3-substituted umbelliferones (5) produced appeared insensitive to the nature of the migrating group in all instances.

The *para*-Claisen rearrangement approach described above permits the ready and efficient preparation of 6-prenylated derivatives of umbelliferone, particularly the higher prenyl-ogues, and is complemented by our *ortho*-Claisen rearrangement approach to 6-allylumbelliferone.<sup>1</sup>

We thank the S.E.R.C. and Shell Research, Sittingbourne for financial support (to N. C.).

Received, 28th April 1986; Com. 563

## References

- 1 N. Cairns, L. M. Harwood, D. P. Astles, and A. Orr, *J. Chem. Soc., Chem. Commun.*, 1986, 182.
- 2 N. Cairns, L. M. Harwood, and D. P. Astles, *J. Chem. Soc., Chem. Commun.*, 1986, 750.
- 3 B. E. Ellis and S. A. Brown, *Can. J. Biochem.*, 1974, **52**, 734. For a complete survey of naturally occurring coumarins see R. D. H. Murray, J. Mendez, and S. A. Brown, 'The Natural Coumarins, Occurrence, Chemistry and Biochemistry,' Wiley-Interscience, New York, 1982.
- 4 R. D. H. Murray, M. M. Ballantyne, and K. P. Mathui, *Tetrahedron*, 1971, **27**, 1247; see also, J. Hlubucek, E. Ritchie, and W. C. Taylor, *Aust. J. Chem.*, 1971, **24**, 2347, 2355.
- 5 Others have used the *para*-Claisen rearrangements of prenyl ethers to introduce a prenyl group into an aromatic nucleus, e.g.: R. D. H. Murray, M. M. Ballantyne, T. C. Hogg, and P. H. McCabe, *Tetrahedron*, 1975, **31**, 2960.
- 6 G. A. Kuznetsova, *Ombelliferes: Contrib. Pluridiscip. Syst., Actes Symp. Int., 2nd.*, 1977, 515; *Chem. Abstr.*, 1978, **89**, 126088r.
- 7 For a related triple rearrangement leading to 3-allylated products see: M. M. Ballantyne, P. H. McCabe, and R. D. H. Murray, *Tetrahedron*, 1971, **27**, 871.
- 8 E. von Gorup-Besanez, *Liebigs Ann. Chem.*, 1876, **183**, 321.