

Evidence for Generation of the Unsaturated Sila-acetate Species $\text{Me}(\text{O}^-)\text{Si}=\text{O}$ by Dissociation of the Silanediolate Dianion $m\text{-ClC}_6\text{H}_4\text{CH}_2\text{SiMe}(\text{O}^-)_2$

Jadwiga Chmielecka,^a Julian Chojnowski,^a Colin Eaborn,^b and Włodzimierz A. Stańczyk^a

^a Polish Academy of Sciences, Centre of Molecular and Macromolecular Studies, Boczna 5, 90-362 Łódź, Poland


^b School of Chemistry and Molecular Sciences, University of Sussex, Brighton BN1 9QJ, U.K.

Kinetic studies indicate that in the cleavage of $m\text{-ClC}_6\text{H}_4\text{CH}_2\text{SiMe}(\text{OH})_2$ by NaOH in $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ there is a major contribution by unimolecular dissociation of the dianion $m\text{-ClC}_6\text{H}_4\text{CH}_2\text{SiMe}(\text{O}^-)_2$ to give the acetate ion analogue $\text{Me}(\text{O}^-)\text{Si}=\text{O}$.

We previously suggested that when the compounds RSiMe_2OMe ($\text{R} = m\text{-ClC}_6\text{H}_4\text{CH}_2$ or $\text{PhC}\equiv\text{C}$) are dissolved in 5% v/v $\text{H}_2\text{O}-\text{MeOH}$ containing NaOH, the cleavages of the $\text{R}-\text{Si}$ bonds involve concurrent reactions of the neutral species RSiMe_2OMe and RSiMe_2OH with base anion and the unimolecular dissociation of the anionic species RSiMe_2O^- to give a silanone intermediate $\text{Me}_2\text{Si}=\text{O}$.^{1,2} For those cleavages, the plots of the observed first-order rate constants (k_{obs}) against base concentration, $[\text{OH}^-]$, initially rise steeply with increase in $[\text{OH}^-]$ and then tail off to almost constant values of k_{obs} , and can be nicely accounted for in terms of a substantial contribution by the unimolecular decomposition. However, the plots of k_{obs} against $[\text{OH}^-]$ could have the same form even if only cleavages of the neutral species were significant, and so (even though there is independent evidence for ejection of an anion from the silanolate ion to give a silanone^{1,3}) we decided to seek unambiguous evidence for the unimolecular dissociation of a negatively charged species by examining the cleavage of the diol $m\text{-ClC}_6\text{H}_4\text{CH}_2\text{SiMe}(\text{OH})_2$ [subsequently denoted by $\text{A}(\text{OH})_2$] in $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ mixtures containing NaOH.

In basic solutions of the diol $\text{A}(\text{OH})_2$ equilibria (i) and (ii) will be rapidly established. That the second ionization, to give $\text{A}(\text{O}^-)_2$, would be likely to be significant was suggested by the fact that the second dissociation constant for orthosilicic acid, $\text{Si}(\text{OH})_4$, is only about 100 times smaller than the first, and that in more concentrated aqueous alkali there is virtually complete conversion into $\text{Si}(\text{OH})_2(\text{O}^-)_2$.⁴ A similar difference can be assumed between the first and second dissociation constants of $\text{A}(\text{OH})_2$, and so conversion into the dianions $\text{A}(\text{O}^-)_2$ would be essentially complete in the very strongly basic solutions used in the present studies.

Bimolecular cleavage of the doubly charged species $\text{A}(\text{O}^-)_2$ by attack of OH^- at silicon must be negligibly slow, but there is the possibility of unimolecular cleavage, analogous to that recognised for RSiMe_2O^- species, as depicted in equation (iii).



$$k_{\text{obs}} = k_{\text{A}(\text{OH})_2}[\text{A}(\text{OH})_2][\text{OH}^-] + k_{\text{A}(\text{OH})(\text{O}^-)}[\text{A}(\text{OH})(\text{O}^-)] + k_{\text{A}(\text{O}^-)_2}[\text{A}(\text{O}^-)_2] \quad (1)$$

$$k_{\text{obs}} = \left[\frac{B}{[\text{OH}^-]} + k_{\text{A}(\text{O}^-)_2} \right] \frac{K_1 K_2 [\text{OH}^-]^2}{1 + K_1 [\text{OH}^-] + K_1 K_2 [\text{OH}^-]^2} \quad (2)$$

$$\text{where } K_1 = \frac{[\text{A}(\text{OH})(\text{O}^-)][\text{OH}^-]}{[\text{A}(\text{OH})_2]}, \quad K_2 = \frac{[\text{A}(\text{O}^-)_2][\text{OH}^-]}{[\text{A}(\text{OH})(\text{O}^-)]}, \\ \text{and } B = (k_{\text{A}(\text{OH})_2}/K_1 K_2) + (k_{\text{A}(\text{OH})(\text{O}^-)}/K_2).$$

If such a process occurs, then in appropriate regions of base concentration three cleavage processes should contribute to

the observed rate constant k_{obs} , the value of which, provided that the concentration of base is much greater than that of $\text{A}(\text{OH})_2$, will be given by equation (1). By use of the definitions shown of the equilibrium constants K_1 and K_2 for the first and second ionizations of $\text{A}(\text{OH})_2$, equation (1) can be transformed into equation (2). It follows that when much of the substrate, say 90%, is present as the dianion, doubling of the base concentration will reduce the concentration of $\text{A}(\text{OH})_2$ by a factor approaching four and that of $\text{A}(\text{OH})(\text{O}^-)$ by a factor approaching two. At the same time the rate of cleavage of the remaining $\text{A}(\text{OH})_2$ should be doubled, but the contribution from this cleavage will still be halved, as will that from the unimolecular cleavage of $\text{A}(\text{OH})(\text{O}^-)$. Consequently, at the higher base concentrations the contributions from the cleavages of $\text{A}(\text{OH})_2$ and $\text{A}(\text{OH})(\text{O}^-)$ will fall off rapidly, and in the limit the observed rate constant will be that for the unimolecular cleavage of $\text{A}(\text{O}^-)_2$. Thus a plot of k_{obs} against $[\text{OH}^-]$ should show a steep initial steep rise in k_{obs} as the rate of bimolecular cleavage of the $\text{A}(\text{OH})_2$ species rises in line with $[\text{OH}^-]$, but then should begin to tail off (as in the cleavage of RSiMe_2OH) as the monoanion $\text{A}(\text{OH})(\text{O}^-)$ is formed. As $[\text{OH}^-]$ is increased further and significant amounts of the dianion $\text{A}(\text{O}^-)_2$ are formed, then if the dianion





Figure 1. Plot of k_{obs} against concentration of KOH for cleavage of $\text{A}(\text{OH})_2$ in 1 : 1 v/v $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ (a) at 50°C and (b) at 25°C. The lines shown are those generated from equation (2) by use of the values of K_1 , K_2 , B , and $k_{\text{A}(\text{O}^-)_2}$ specified in the text.




Figure 2. Plot of k_{obs} against concentration of KOH for cleavage of $A(\text{OH})_2$ in 3:1 v/v $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ at 25°C. The line shown is that generated from equation (2) by use of the values of K_1 , K_2 , B , and $k_{A(\text{O}^-)_2}$ specified in the text.

were unreactive, k_{obs} would begin to fall off sharply, ultimately falling effectively to zero, which is not observed. If, however, the dianion undergoes cleavage at a significant rate, then the value of k_{obs} should level off to a plateau, or pass through a maximum and subsequently level off at a constant value corresponding to the rate constant for the unimolecular cleavage of the dianion alone.

The plots of k_{obs} against $[\text{OH}^-]$ for cleavage of $A(\text{OH})_2$ in 1:1 and 3:1 v/v $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ are shown in Figures 1 and 2, and in our view their shapes must mean that the dianionic species undergoes spontaneous cleavage at a substantial rate. This conclusion remains valid irrespective of the limitations of the rough quantitative analysis which follows.

Rough values of K_1 (based largely on measurements of the dissociation constant for $\text{PhCH}_2\text{SiMe}_2\text{OH}$ in the two media) of 400 and $6000 \text{ dm}^3 \text{ mol}^{-1}$ can be estimated for $A(\text{OH})_2$ in 1:1 and 3:1 $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ respectively. The values of K_2 are then assumed to be smaller by a factor of 10^2 , the ratio between the first and second ionization constants of $\text{Si}(\text{OH})_4$.⁴ A computer analysis then gives values of $k_{A(\text{O}^-)_2}$ and B which produce the best fit to the plots of k_{obs} against $[\text{OH}^-]$, and in Figures 1 and 2 the lines shown are those based on equation (2) with values of $10^4 k_{A(\text{O}^-)_2}$ of 2.7 and 30 s^{-1} in 1:1 and 3:1 $\text{Me}_2\text{SO}-\text{H}_2\text{O}$ and corresponding values of $10^5 B$ of 4.5 and $14 \text{ mol dm}^{-3} \text{ s}^{-1}$.

We suggest that the results provide convincing evidence for the operation of the process depicted as reaction (iii). The postulated intermediate $\text{Me}(\text{O}^-)\text{Si}=\text{O}$ [a derivative of the hypothetical metasilicic acid $\text{H}(\text{OH})\text{Si}=\text{O}$] is an analogue of acetate ion, and should be stabilized in the same way as the latter (though to a smaller extent) by delocalization of the negative charge over both oxygen atoms.

We thank Dr. Marek Cypryk for the computer analyses, the Polish Academy of Sciences for support (for Problem CPBP 01.13), and S.E.R.C. for support (to C. E.).

Received, 13th May 1987; Com. 645

References

- 1 Z. H. Aiube, J. Chojnowski, C. Eaborn, and W. A. Stańczyk, *J. Chem. Soc., Chem. Commun.*, 1983, 493.
- 2 C. Eaborn and W. A. Stańczyk, *J. Chem. Soc., Perkin Trans. 2*, 1984, 2099.
- 3 Z. H. Aiube, D. Phil. Thesis, University of Sussex, 1984.
- 4 N. Ingri in 'Aqueous Silicic Acid, Silicates, and Silicate Complexes in Biochemistry of Silicon and Related Problems,' eds. S. Bendz and I. Lindquist, Plenum Press, New York, 1978, pp. 3–51.