

Intramolecular Wittig Reaction of α,α' -Dioxocycloalkylidene-tributylphosphoranes: Formation and Trapping of Cycloalkyn-2-ones with Five- and Six-membered Rings

Hidenobu Ohmori, Hatsu Maeda, Chihiro Ueda, and Masaichiro Masui*

Faculty of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565, Japan

Reaction of the title phosphoranes with homophthalic anhydride or 1,3-diphenylisobenzofuran in the presence of Me_3SiCl gave the corresponding Diels-Alder adducts, indicating the transient intermediacy of cycloalkyn-2-ones with five- and six-membered rings.


The existence of short-lived cycloalkynes such as cyclopentynes and cyclohexynes has been demonstrated by several methods, and their cycloaddition reactions with dienes have been used in synthesis.¹ However, the corresponding conjugated cycloalkynones of type (2) are not known;^{2†} cyclo-octyn-2-one is the homologue of smallest ring size hitherto reported.³ We report here preliminary results on the trapping of the cycloalkynones (2) generated by intramolecular Wittig reaction of the title phosphoranes (1). Electrochemical oxidation of Bu_3P in the presence of cyclic 1,3-diketones has provided a convenient one-step procedure for

the preparation of the phosphoranes (1),[‡] which would be difficult to obtain by conventional methods for non-cyclic α,α' -dioxoalkylidene-triphenylphosphoranes.⁴

When a mixture of (1) (1 mmol), homophthalic anhydride (diene equivalent)⁵ (3.3 mmol), and Me_3SiCl (3 mmol) in toluene was heated at 150 °C for 40 h in a sealed tube, the

† Trapping of benzynequinone [C. W. Rees and D. E. West, *Chem. Commun.*, 1969, 647; *J. Chem. Soc. (C)*, 1970, 583] and 4,5-didehydrotropone (T. Nakazawa and I. Murata, *Angew. Chem., Int. Ed. Engl.*, 1975, **14**, 711) has been reported.

‡ The phosphoranes (1) were prepared by essentially the same electrochemical procedure as that reported for $\text{Ph}_3\text{P}=\text{C}(\text{COR})_2$ (H. Ohmori, H. Maeda, M. Tamaoka, and M. Masui, *Chem. Pharm. Bull.*, 1988, **36**, 613). Constant-current electrolysis of Bu_3P (11 mmol) in CH_2Cl_2 (35 ml) containing the appropriate 1,3-diketone (22 mmol) and 2,6-lutidinium perchlorate (22 mmol) gave (1a), (1b), and (1c) in yields (isolated) of 70, 58, and 59%, respectively [one-compartment electrolysis cell; graphite plate anode and Pt foil cathode; 30 mA (current density, 1 mA/cm²); 2 F/mol of Bu_3P were consumed at ambient temperature].

Scheme 1. Reagents and conditions: i, Me_3SiCl , toluene, 150°C .

Diels–Alder adducts (**3a**), (**3b**), and (**3c**) were obtained§ in yields of 17, 23, and 5%, respectively. Similarly, the adducts (**4a**) (5%) and (**4b**) (14%) were formed§ by using 1,3-diphenylisobenzofuran as the diene. Since [4 + 2] cycloaddition reactions have often been employed as proof for the intermediacy of highly reactive cycloalkynes,^{1,6} the present results together with the following observation support the formation of cycloalkyn-2-ones (**2**) from the phosphoranes (**1**). Thus, the acetylenic ketone (**6**) (80%) and the β -chloroenone (**7**) (18%) were isolated in the thermal decomposition of the phosphorane (**5**) performed under the conditions described above.

The chloroenone (**8a**) was also formed in 8% yield along with tarry materials by heating a mixture of the phosphorane (**1a**) and Me_3SiCl in toluene under the same conditions. However, the enone (**8a**) cannot be the intermediate in the conversion of (**1a**) into the adduct (**3a**); the formation of (**3a**) was negligible in the thermal reaction of (**8a**) prepared separately⁷ with homophthalic anhydride in the presence of Me_3SiCl , and 60% of (**8a**) was recovered.

Although the optimum conditions for the generation of (**2**) remain to be explored and the number of dienes examined is limited, we believe the phosphorane (**1**) will find potential synthetic utility as cycloalkyn-2-one equivalents.

Received, 10th March 1988; Com. 8/00974K

References

- M. Nakagawa, in 'The Chemistry of the Carbon–Carbon Triple Bond,' ed. S. Patai, Wiley, Chichester, 1978, pp. 636–643.
- F. Kienzle and R. E. Minder, *Helv. Chim. Acta*, 1987, **70**, 1537.
- P. E. Eaton and C. E. Stubbs, *J. Am. Chem. Soc.*, 1967, **89**, 5722.
- D. A. Ben-Efraim, in ref. 1, p. 781.
- Y. Tamura, A. Wada, S. Sasho, and Y. Kita, *Tetrahedron Lett.*, 1981, **22**, 4283.
- D. H. R. Barton, G. Bashiardes, and J.-L. Fourrey, *Tetrahedron*, 1988, **44**, 147.
- R. D. Clark and H. Heathcock, *J. Org. Chem.*, 1976, **41**, 636.
- Y. Tamura, A. Wada, S. Okuyama, S. Fukumori, Y. Hayashi, N. Goda, and Y. Kita, *Chem. Pharm. Bull.*, 1981, **29**, 1312.

§ All the Diels–Alder adducts (**3**) and (**4**) gave the expected analytical results. (**3a**): m.p. $98\text{--}99^\circ\text{C}$ (lit.⁸ $97\text{--}98^\circ\text{C}$); (**3b**): m.p. $93\text{--}94^\circ\text{C}$ (lit.⁸ $93.5\text{--}94^\circ\text{C}$); (**3c**): m.p. 112°C ; ν_{max} 3300br., 1675, 1635 cm^{-1} ; ^1H n.m.r. (CDCl_3) δ 2.60–3.05 (m, 2H, CH_2), 3.10–3.50 (m, 2H, CH_2), 7.20–8.50 (m, 4H, ArH), 8.10–8.50 (m, 1H, ArH); (**4a**): m.p. $236\text{--}238^\circ\text{C}$; ν_{max} 1680 cm^{-1} ; ^1H n.m.r. (CDCl_3) δ 0.98 (s, 6H, CH_3), 2.44 (s, 2H, CH_2), 2.67 (s, 2H, CH_2), 7.00–7.70 (m, 14H, ArH); (**4b**): m.p. $218\text{--}220^\circ\text{C}$; ν_{max} 1680 cm^{-1} ; ^1H n.m.r. (CDCl_3) δ 2.00 (q, J 6 Hz, 2H, CH_2), 2.55 (t, J 6 Hz, 2H, CH_2), 2.72 (t, J 6 Hz, 2H, CH_2), 7.00–7.70 (m, 14H, ArH).