

Kinetics and Mechanism of the Thermal Decomposition of Hexaborane(12) in the Gas Phase

Robert Greatrex, Norman N. Greenwood, and Simon D. Waterworth

School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.

The gas-phase thermolysis of *arachno*-B₆H₁₂ produces predominantly B₅H₉ and B₂H₆ in a molar ratio of 2 : 1 via a first-order reaction having Arrhenius parameters which are essentially identical to those reported for the decomposition of the structurally related B₅H₁₁; these results imply a mechanism involving elimination of BH₃ as the rate-determining initial step in both reactions.

Progress has been slow in elucidating a coherent mechanistic description of the facile thermal interconversions of the simple binary boranes, not only because of the inherent complexity of the reactions but also because of difficulties in acquiring reliable kinetic data for these highly reactive, air-sensitive species.¹ We have developed a quantitative mass spectrometric technique for monitoring these reactions in detail, and have thereby gained new insights into the thermal decompositions and interconversion reactions of *nido*-B₆H₁₀² and the two *arachno* species B₄H₁₀^{3,4a} and B₅H₁₁.⁴ We now report preliminary results of the first quantitative kinetic study on B₆H₁₂. The results are important because they provide the first opportunity to correlate kinetic, mechanistic, and structural patterns of behaviour in a unique series of closely related *arachno* binary boranes.

A typical reaction profile for the thermolysis of B₆H₁₂ at ca. 100 °C is shown in Figure 1 for an initial pressure of 3.14 mmHg. From this and similar profiles recorded over the pressure and temperature ranges 1.8–10.0 mmHg and 75–150 °C, it emerges that this thermolysis is the most straightforward of all borane decompositions studied so far. The overall stoichiometry is very well defined, with at least 80% of the B₆H₁₂ decomposing to B₅H₉ plus ½B₂H₆. From a detailed analysis of initial rates, the consumption of B₆H₁₂ and the production of B₅H₉ and of B₂H₆ were all found to be accurately first-order with respect to the concentration of B₆H₁₂. In addition, a small amount of H₂ is produced, and even less B₆H₁₀; the latter builds up rather slowly but persists in the later stages of the reaction when all the B₆H₁₂ has decomposed. Only traces of higher boranes such as B₁₀H₁₄ are observed, and involatile solid hydrides account for as little as






Figure 1. Reaction profile for the thermolysis of B₆H₁₂ (P_0 3.14 mmHg) at 99.4 °C. ● B₆H₁₂, ◇ B₅H₉, □ B₂H₆, △ B₁₀H₁₄, and ○ H₂.

10% of the boron consumed, compared with typical values of 40–50% in the thermolysis of B₄H₁₀³ and B₅H₁₁.^{4b}

These results can readily be explained by a simple two-step mechanism involving the first order rate-determining elimination of {BH₃} which then rapidly dimerizes to B₂H₆ [reactions (1) and (2)]. Previous qualitative studies of the decomposition of B₆H₁₂ in the gas phase⁵ have yielded no mechanistic information and have sometimes suggested more complex behaviour than observed in the present work.



Our recent kinetic study of B₅H₁₁⁴ has shown that this borane also decomposes via the initial rate-determining elimination of {BH₃} from the cluster [reaction (3)], and it is therefore particularly significant that the activation energy and pre-exponential factor now found for the B₆H₁₂ thermolysis (E_a 75.0 ± 5.8 kJ mol⁻¹; A 3.8 × 10⁷ s⁻¹) are essentially identical to the values obtained for B₅H₁₁ (E_a 72.6 ± 2.4 kJ mol⁻¹; A 1.3 × 10⁷ s⁻¹). This is persuasive additional evidence that the rate-determining steps in the two decompositions do indeed involve very similar processes and this finds a ready interpretation in terms of the detailed molecular structures of gaseous B₅H₁₁⁶ and B₆H₁₂,⁷ as recently determined by electron diffraction. As shown in Figure 2,⁷ these two species bear a close structural relationship to each other and to B₄H₁₀.⁸ Thus, notional replacement of H_{endo} and one H_μ on, say, B(4) in B₄H₁₀ by a BH₃ group yields B₅H₁₁ and




Figure 2. Structural relationship between the *arachno*-boranes (a) B₄H₁₀,⁸ (b) B₅H₁₁,⁶ and (c) B₆H₁₂,⁷ as determined by gas-phase electron diffraction.

repetition of this process on the opposite side of the molecule, *i.e.* at B(2), generates the observed structure of B_6H_{12} having C_2 symmetry. In view of the similarity of the Arrhenius parameters for the decomposition of B_6H_{12} and B_5H_{11} , it seems reasonable to identify these structurally similar BH_3 groups as the fragments involved in the initial steps (1) and (3). In B_4H_{10} , by contrast, these particular incipient BH_3 groups are absent and its decomposition is characterized by quite different Arrhenius parameters (E_a 99.2 ± 0.8 kJ mol⁻¹; A 6.0×10^{11} s⁻¹).³ In this case the initial step is believed to involve elimination of H_2 to give $\{B_4H_8\}$. This type of reaction is clearly not favoured in the case of B_5H_{11} and B_6H_{12} , though small amounts of B_6H_{10} are produced in the thermolysis of B_6H_{12} , and further work is necessary to determine whether this species and dihydrogen arise from a competing, but minor, reaction channel involving direct elimination of H_2 from B_6H_{12} .

We thank Mr. D. Singh for experimental assistance and the S.E.R.C. for a maintenance grant (to S. D. W.).

Received, 22nd March 1988; Com. 8/01156G

References

- 1 N. N. Greenwood and R. Greatrex, *Pure Appl. Chem.*, 1987, **59**, 857, and references therein.
- 2 R. Greatrex, N. N. Greenwood, and G. A. Jump, *J. Chem. Soc., Dalton Trans.*, 1985, 541.
- 3 R. Greatrex, N. N. Greenwood, and C. D. Potter, *J. Chem. Soc., Dalton Trans.*, 1986, 81.
- 4 (a) M. D. Attwood, R. Greatrex, and N. N. Greenwood, *J. Chem. Soc., Dalton Trans.*, submitted for publication; (b) *ibid.*, submitted for publication.
- 5 D. F. Gaines and R. Schaeffer, *Inorg. Chem.*, 1964, **3**, 438; A. L. Collins and R. Schaeffer, *ibid.*, p. 2153; C. A. Lutz, D. A. Phillips, and D. M. Ritter, *ibid.*, p. 1191; S. J. Steck, G. A. Pressley, F. E. Stafford, J. Dobson, and R. Schaeffer, *ibid.*, p. 2452; T. C. Gibb, N. N. Greenwood, T. R. Spalding, and D. Taylorson, *J. Chem. Soc., Dalton Trans.*, 1979, 1392.
- 6 R. Greatrex, N. N. Greenwood, D. W. H. Rankin, and H. E. Robertson, *Polyhedron.*, 1987, **6**, 1849.
- 7 R. Greatrex, N. N. Greenwood, M. B. Millikan, D. W. H. Rankin, and H. E. Robertson, *J. Chem. Soc., Dalton Trans.*, in the press.
- 8 C. J. Dain, A. J. Downs, G. S. Laurensen, and D. W. H. Rankin, *J. Chem. Soc., Dalton Trans.*, 1981, 472.