

Acetylenedicarbalddehyde: Isolation and some Examples of Exclusive Dienophilicity under Neutral Conditions

Dominique Stéphan,^a Alain Gorgues,^{*a,b} Ahmed Belyasmine,^a and André Le Coq^a

^a Laboratoire de Synthèse Organique, U. A. 415 du CNRS, Université de Rennes, Campus de Beaulieu, 35042 Rennes Cédex, France

^b Groupe de Recherche de Chimie Organique et Bio-Organique, Université d'Angers, 2 Bd. Lavoisier, 49045 Angers Cédex, France



Acetylenedicarbalddehyde (1) is isolated in the pure state after acidolysis of its monoacetal (2) with an excess of formic acid followed by dehydration of residual HCO₂H into CO with P₂O₅; under neutral conditions, (1) reacts with conjugated dienes as a dienophile only, in contrast to what occurs in acidic medium.

We have previously shown that acetylenedicarbalddehyde (ADCA) (1) and its monoacetal (2), both attractive synthetic intermediates,^{1,2,3b} were readily generated by the acidolysis of the diacetal (3) with a large excess of formic acid, equation (1) and (2), respectively.⁴

Whereas (2) is easily isolable,^{3,4†} the unstable ADCA could neither be isolated in the pure state, nor conveniently obtained in aprotic solutions.⁴ Although formic acid solutions of (1), see equation (1), can be used in some syntheses,^{1,2,4} residual HCO₂H is often incompatible in many reactions (*e.g.* with carbanionic species, dipolar compounds etc.), or can strongly modify the reactivity of ADCA⁴ (*vide infra*). Here we describe procedures for eliminating HCO₂H, from the formolysis reactions in order to get aprotic solutions of ADCA, and for isolating ADCA in the pure form. They are based on the use of P₂O₅, an electrophilic reagent which promotes the dehydration of formic acid into CO.⁵





After treatment of (2) (2.5 g), under nitrogen and in the dark, HCO₂H (5 ml; dried over anhydrous CuSO₄) and anhydrous CuSO₄ (2 g), for 1.5 h at 20 °C (or 1 h at 25 °C) (see equation 3), and dilution with CH₂Cl₂ (200 ml), the resulting

solution was added dropwise to a 100 ml CH₂Cl₂ suspension of P₂O₅ (17 g) for 10–15 min, at room temperature and in the dark. When CO was evolved (1–1.5 h of stirring), the reaction mixture was filtered and concentrated to 100 ml (rotary evaporation under reduced pressure, bath 20–25 °C); this solution A contains (1) (50–55% yield), ethyl formate, and only a small amount of HCO₂H⁶ (<10% with respect to ADCA).



It was necessary to use a purification procedure consistent with the explosive character of ADCA in the dry state (*vide infra*) which allows the elimination of the remaining HCO₂H. Thus, to the solution A, diethyl phthalate (10 ml) (as a high boiling solvent) and P₂O₅ (0.5 g) were added, and the greater

† Compounds (2), (3), and 5,6-dimethyl-4,7-dihydroisobenzofuran are marketed by Janssen Pharmaceutica.



part of the dichloromethane and ethyl formate slowly distilled off under reduced pressure (bath 15 °C, nitrogen capillary) for 1 h (completion of HCO₂H decomposition). When the temperature of the bath was raised to 50–60 °C, ADCA condensed (column 20–22 °C) in the refrigerant cooled to –40 °C as pale-yellow needles that explode, even under nitrogen, on melting (m.p. –11 to –10 °C). This explosion can be avoided by dissolving the solid at a temperature well below –11 °C in an aprotic solvent (CH₂Cl₂, CHCl₃, CCl₄), the resulting solutions (up to 0.4 mol/l) being fairly stable for a week when stored at –20 °C [30% yield of (1)].‡

For synthetic purposes, HCO₂H-free solutions of (1) can be more readily obtained by a further treatment of the solution A with P₂O₅ (5.7 g) in CH₂Cl₂: thus is obtained solution B [30% yield of (1)]. If ethyl formate and CH₂Cl₂ cannot be used in subsequent reactions, both can be distilled off under reduced pressure (bath 15–20 °C) after dilution of solution B with another higher boiling solvent (e.g. toluene) without any significant decrease in yield.

Whereas under HCO₂H conditions, ADCA shows a variety of reactivity towards conjugated dienes^{1,4} (i.e. Diels–Alder and/or pseudo-Michael adducts), in contrast, under neutral conditions (from freshly prepared solutions of pure ADCA or from solution B), only the dienophilic behaviour is observed irrespective of the aromaticity of the diene.

The most significant example is provided by furan which leads to the exclusive formation of (5) in the presence of formic acid,⁴ and of (4) (pale yellow solid, m.p. 70–73 °C, 98% yield) in the presence of pure ADCA collected into CCl₄.§ It should be noted that the addition of formic acid to a

CHCl₃ solution of (4) does not give rise to (5), which suggests a kinetically controlled formation of the latter under acidic conditions.^{4,7} Similarly, the dienes diphenylfulvene, anthracene, 4,7-ethano-4,7-dihydrobenzofuran,⁸ and 5,6-dimethyl-4,7-dihydroisobenzofuran react to form the cycloadduct analogous to (4) instead of the mixture of products which is formed when the reaction takes place in the presence of HCO₂H.

With non-aromatic, or α,α'-disubstituted aromatic, dienes which are not subject to electrophilic substitution and will therefore form cycloadducts only, cycloadditions take place more slowly, but more cleanly in neutral solutions. For example, the product between cyclopentadiene and (1) was formed in 80% yield and that between cyclohexa-1,3-diene was formed in 75% yield, both from solution B. Also, 2,3-dimethylbuta-1,3-diene forms a cycloaddition product with (1) (pale yellow solid, m.p. 106–107 °C) in a neutral medium but only a tarry material in the presence of HCO₂H. Finally, the non-isolated, sensitive cycloadducts between (1) and 2,5-dimethylfuran and (1) and 1,3,3-trimethyl-2-vinyl cyclohexene⁹ could be characterised only by starting with neutral solutions of ADCA.

We thank the Agence de Valorisation de la Recherche for financial support.

Received, 13th October 1987; Com. 1489

References

- 1 A. Gorgues, *Janssen Chim. Acta*, 1986, **4**, 21 and refs. cited.
- 2 P. J. Dunn and C. W. Rees, *J. Chem. Soc., Perkin Trans. 1*, 1987, 1579.
- 3 In a slight modification of procedures given by (a) A. Gorgues and A. Le Coq, *Tetrahedron Lett.*, 1979, 4825, and (b) A. K. Saksena, M. J. Green, P. Mangiaracina, J. K. Wong, W. Kreutner, and A. R. Gulbenkian, *ibid.*, 1985, **26**, 6427: 23 mol of HCO₂H and 25 mol of CHCl₃ per mol of (3), 12 h at 15–20 °C.
- 4 A. Gorgues, A. Simon, A. Le Coq, A. Hercouet, and F. Corre, *Tetrahedron*, 1986, **42**, 351.
- 5 E. G. Graeber and D. S. Cryder, *Ing. Eng. Chem.*, 1935, **27**, 828.
- 6 Probably as formic anhydride which decomposes into CO and HCO₂H, see G. A. Olah, Y. D. Vankar, M. Arvanaghi, and J. Sommer, *Angew. Chem., Int. Ed. Engl.*, 1979, **18**, 614.
- 7 This is in sharp contrast to the case of acrolein and furan: P. Laszlo and J. Lucchetti, *Tetrahedron Lett.*, 1984, **25**, 4387.
- 8 D. Stephan, A. Gorgues, and A. Le Coq, *Tetrahedron Lett.*, 1986, **27**, 4295.
- 9 M. Jalali-Naini, D. Guillermin, and J. Y. Lallemand, *Tetrahedron*, 1983, **39**, 405.

‡ Estimated by ¹H n.m.r. with an internal standard; i.r. (CCl₄) 1685 cm⁻¹ (C=O), and no band in the C≡C absorption range; ¹H n.m.r. (CDCl₃) δ 9.50 (s, CHO); M⁺, m/z 82.0057 (calc. 82.0055).

§ 1.8 mmol ADCA in CCl₄ (6 ml) reacted with 1.7 mmol furan for 2 h at 20 °C.