

## First Samarium-catalysed Coupling of Aldehydes and Ketones

Elisabeth Léonard, Elisabet Duñach,\* and Jacques Périchon

Laboratoire d'Electrochimie, Catalyse et Synthèse Organique, (U.M. N° 28) C.N.R.S., 2, rue Henry Dunant, 94320 Thiais, France

The SmCl<sub>3</sub> catalysed electrochemical coupling of carbonyl compounds affords 1,2-diols in good yields.

A considerable interest in the use of lanthanide compounds as new reagents in organic transformations has developed in the last decade.<sup>1</sup> Among them, low-valent lanthanide derivatives, mainly Sm(II) compounds, have been used as promoters for various coupling reactions between carbonyl compounds and/or organic halides.<sup>2</sup> Sm(II) derivatives have strong reducing properties and react as one-electron transfer reagents in the C-C bond forming reactions. These coupling reactions require the use of stoichiometric amounts of Sm(II)<sup>†</sup> which is transformed into Sm(III) and no further recycling of these last complexes has been described.

We report here the first samarium-catalysed coupling reaction of aldehydes and ketones (pinacol condensation<sup>2a-c,7,11</sup>, involving the recycling of the active Sm(II) species by electrochemical reduction.

Our one-compartment cell electrolysis procedure, based on the use of sacrificial anodes of magnesium or aluminium,<sup>8</sup> allows the use of only 5–10% of SmCl<sub>3</sub> as catalyst precursor. A further advantage of the electrochemical method is the use of SmCl<sub>3</sub> (a stable and commercially available compound) as starting material, with no need to prepare the air and water sensitive Sm(II) derivatives [usually SmI<sub>2</sub><sup>9</sup> or (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Sm<sup>10</sup>].

The carbonyl condensations take place in dimethylformamide (DMF) or *N*-methylpyrrolidone (NMP) as solvents, under mild conditions. The results of the pinacol coupling for several aldehydes and ketones are presented in Table 1.

Aromatic, as well as aliphatic aldehydes afford good yields of the corresponding 1,2-diols. The competition between the reduction of an aromatic chloride and that of an aldehyde is

shown in example 2, where the chloro group remains unchanged in the diol adduct.

Several examples of ketone condensation (entries 5–9) demonstrate the general applicability of the method. Intramolecular coupling of ketones is presented in examples 8 and 9. The strained diphenyl-substituted cyclopropanediol ring (entry 9) is formed quantitatively.

Table 1. Electrochemically mediated SmCl<sub>3</sub>-catalysed carbonyl condensation to 1,2-diols.<sup>a</sup>

$$2 R^1-C(:O)-R^2 \rightarrow R^1-C(OH)R^2-C(OH)R^2-R^1$$

Conditions: i, SmCl<sub>3</sub> (5–10%) + e, then H<sub>3</sub>O<sup>+</sup>.

Entry	R <sup>1</sup>	R <sup>2</sup>	Solvent, T/°C	Anode	Isolated yield/% <sup>c</sup>
1	Ph	H	DMF, 20	Mg	80
2	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	H	NMP, 20	Al	73
3	Me <sub>3</sub> C	H	NMP, 20	Al	72
4	<i>n</i> -C <sub>6</sub> H <sub>13</sub>	H	NMP, <sup>b</sup> 20	Al	75
5	Ph	Me	DMF, 20	Mg	90
6	Ph	Ph	NMP, 20	Al	85
7	<i>n</i> -C <sub>5</sub> H <sub>11</sub>	Me	NMP, <sup>b</sup> 50	Al	40
8	PhC(O)(CH <sub>2</sub> ) <sub>4</sub>	Ph	NMP, 20	Al	90
9	PhC(O)-CH <sub>2</sub>	Ph	NMP, 50	Al	98

<sup>a</sup> Electrolysis: cathode was nickel or stainless steel; solvent, 30 ml; R<sup>1</sup>COR<sup>2</sup>, 10<sup>-2</sup> mol; apparent current densities 0.4–0.5 A dm<sup>-2</sup>; Faradaic yield: 60–100%; reaction time: 5–18 h. Entries 1,5,8,9: 5% SmCl<sub>3</sub>; entries 2–4, 6,7: 10% SmCl<sub>3</sub> (Alfa, anhydrous). <sup>b</sup> 6 ml Et<sub>2</sub>O were added to the NMP solution during electrolysis. <sup>c</sup> Work up: Acid hydrolysis and diethyl ether extraction. The crude product chromatographed on silica gel or alumina(III) column, with pentane-Et<sub>2</sub>O mixtures as eluent. Pinacols recovered as a mixture of *meso*/(±) diastereoisomers.

<sup>†</sup> The samarium-catalysed examples concern hydrogenation,<sup>3</sup> polymerization,<sup>4</sup> isomerization,<sup>5</sup> or redox<sup>6</sup> type processes.

Samarium ions are essential in our experiments for an effective pinacol condensation; in the absence of the catalyst, heptan-2-one yielded heptan-2-ol and mixtures of aldol-type condensation products and no 1,2-diol was formed. The alcohol reduction is an important side-reaction in other reported electrochemical mediated pinacol condensations (applied only to aromatic derivatives).<sup>11</sup>

From a mechanistic point of view, we propose that  $\text{SmCl}_3$  is electrochemically reduced to  $\text{Sm(II)}$  species [ $E_{\text{aq}}^{\circ} \text{Sm(III)}/\text{Sm(II)} = 1.55$  vs. standard calomel electrode (S.C.E.)<sup>12</sup>]. This reacts with the carbonyl compound to form a  $\text{Sm(III)}$ -pinacolate after the coupling of two ketyl radical ions.<sup>11,13</sup> Evidence that  $\text{Sm(II)}$  is effectively formed in solution was obtained from the electrolysis of a  $\text{SmCl}_3$  solution (NMP, Al anode, 1.1 F  $\text{mol}^{-1}$ ), followed by the addition of excess acetophenone. After stirring for 3 h the expected stoichiometric amount of pinacol was isolated.

The originality of the present method lies in the possibility of the  $\text{Sm(III)}$  alcoholates being involved in a transmetallation reaction  $\text{Sm(III)}/\text{Mg(II)}$  [or  $\text{Sm(III)}/\text{Al(III)}$ ], the magnesium (or aluminium) ions being formed *in situ* by anodic oxidation. By metal-exchange,  $\text{Mg(II)}$  [or  $\text{Al(III)}$ ] alcoholates are formed, liberating the  $\text{Sm(III)}$  ion, able to undergo a further electrochemical reduction and thus to re-enter the catalytic cycle.

In conclusion, the samarium-catalysed pinacol condensation demonstrates that electrosynthesis can be an interesting alternative for adapting  $\text{Sm(II)}$  promoted stoichiometric processes to catalytic reactions, by the *in situ* regeneration of the low-valent active species.

We thank Professor H. Kagan for useful discussions.

Received, 17th October 1988; Com. 8/04103B

## References

- (a) H. B. Kagan and L. J. Namy, *Tetrahedron Report* 213, 1986, 42, 6573; (b) H. B. Kagan, *Inorg. Chim. Acta*, 1987, 140, 3; (c) W. J. Evans and D. K. Drummond, *J. Am. Chem. Soc.*, 1988, 110, 2772.
- (a) J. L. Namy, J. Soupe, and H. B. Kagan, *Tetrahedron Lett.*, 1983, 765; (b) J. Soupe, L. Danon, J. L. Namy, and H. B. Kagan, *J. Organomet. Chem.*, 1983, 250, 227; (c) G. A. Molander and C. Kenny, *J. Org. Chem.*, 1988, 53, 2132, and refs. therein; (d) K. Otsubo, K. Kawamura, J. Inanaga, and M. Yamaguchi, *Chem. Lett.*, 1987, 1487.
- (a) W. J. Evans, *J. Organomet. Chem.*, 1983, 250, 217; (b) P. L. Watson and C. W. Parshall, *Acc. Chem. Res.*, 1985, 18, 51.
- G. Jeske, H. Lauke, H. Mauerman, H. Schumann, and T. J. Marks, *J. Am. Chem. Soc.*, 1985, 107, 8111, and refs. therein.
- J. Prandi, J. L. Namy, G. Menoret, and H. B. Kagan, *J. Organomet. Chem.*, 1985, 285, 499.
- (a) J. L. Namy, J. Soupe, J. Collin, and H. B. Kagan, *J. Org. Chem.*, 1984, 49, 2045; (b) J. Collin, J. L. Namy, and H. B. Kagan, *Nouv. J. Chim.*, 1986, 10, 229.
- For methods of pinacol formation, see A. Furstner, R. Csuk, C. Rohrer, and H. Weidmann, *J. Chem. Soc., Perkin Trans. 1*, 1988, 1729, and refs. therein.
- (a) M. Heintz, O. Sock, C. Saboureau, and J. Périchon, *Tetrahedron*, 1988, 44, 1631; (b) G. Silvestri, S. Gambino, G. Filardo, and A. Gulotta, *Angew. Chem., Int. Ed. Engl.*, 1984, 23, 979.
- P. Girard, J. L. Namy, and H. B. Kagan, *J. Am. Chem. Soc.*, 1984, 102, 2693.
- W. J. Evans, *Inorg. Chim. Acta*, 1987, 139, 169, and refs. therein.
- F. Fourier and M. Fourier, *Can. J. Chem.*, 1986, 64, 881.
- H. B. Kagan and J. L. Namy, in 'Handbook on the Physics and Chemistry of Rare Earth,' eds. K. A. Gschneidner, Jr., and L. Eyring, ch. 50, North Holland Publ. Co., Amsterdam, 1984.
- H. B. Kagan, J. L. Namy, and P. Girard, *Tetrahedron*, 1981, 37, Suppl. 1, 175.