


### *Corrigenda*

#### **Stereospecificity in the Synthesis of C-5 Modified Nucleosides using Radical Chemistry: Furanosidic Chain-lengthening through C-4**

Derek H. R. Barton, S. D. Géro, Béatrice Quiclet-Sire, and Mohammad Samadi

*J. Chem. Soc., Chem. Commun.*, 1988, 1372.


Structures (12)–(14) and (21)–(23) are incorrect. The correct structures are shown below.



(12)  $X = \text{CO}_2\text{H}$

(13)  $X = \text{CH}_2\text{CHZ}-\text{SO}_2\text{Ph}$

(14)  $X = \text{CH} \overset{\cdot}{=} \text{CH}-\text{SO}_2\text{Ph}$



(21)  $X = \text{CO}_2\text{H}$

(22)  $X = \text{CH}_2\text{CHZ}-\text{SO}_2\text{Ph}$


(23)  $X = \text{CH} \overset{\cdot}{=} \text{CH}-\text{SO}_2\text{Ph}$

## Generation of Spirocyclic Quaternary Centres via a Tandem Free Radical Cyclopropylcarbinyl Rearrangement–Cyclisation Strategy

J. D. Harling and W. B. Motherwell

*J. Chem. Soc., Chem. Commun.*, 1988, 1380.

Structures (15) and (16) are incorrect. The correct structures are shown below.



# An Unexpected Rearrangement of 2-(2-Benzofuranyl)-benzonitriles: a New Route to Phenylmethylene-dihydro-isoindolinones

Jean Guillaumel, Nicole Bocvara, Pierre Demerseman, and René Rover

J. Chem. Soc., Chem. Commun., 1988, 1604

Compound nos. (1)–(4) were omitted from Table 1 on p. 1605. The correct Table is as shown below.

Table 1.

Solvent	Time	Temp./°C	% Yield								
			(1)	(2)	(3)	(4)	a	b	a	b	a
EtOH	72 h	80		6				81			
MeOCH <sub>2</sub> CH <sub>2</sub> OH	1 h	125	7	12	78	58	Traces	Traces	3	14	15
	16 h	—			5				6	92	85
HOCH <sub>2</sub> CH <sub>2</sub> OH	5 min	198		31		45			2		3
	15 min	—	3	2	28	25	32	26	25	25	26
	1 h	—			5	Traces	52	39	42		45