

## 1,1-Dimetallagermacyclopent-3-enes: Precursors to Transition Metal Substituted Germylenes (Germanediyls)

Deqing Lei and Mark J. Hampden Smith\*


*Department of Chemistry, University of New Mexico, Albuquerque, NM 87131 U.S.A.*

Transition metal substituted germylenes have been generated by pyrolysis of 1,1-dimetallagermacyclopent-3-enes and by the metathesis reaction between  $\text{GeI}_2$  and  $[\text{Mo}(\text{CO})_3\text{Cp}]^-$  ( $\text{Cp} = \text{C}_5\text{H}_5$ ).


---

The chemistry of silylene and germylene reactive intermediates has been intensely studied in recent years, with particular emphasis on the preparation of suitable generators,<sup>1</sup> trapping reagents,<sup>2</sup> and their rearrangement reactions.<sup>3</sup> However, the work has been confined almost exclusively to

the study of silylenes and germylenes containing electronegative substituents. The sole exception is the recent generation of a transition metal substituted silylene,  $\{\text{Si}[\text{Fe}(\text{CO})_2\text{Cp}]\text{Me}\}$  ( $\text{Cp} = \text{C}_5\text{H}_5$ ), by Marietti-Migrani and West,<sup>4</sup> which was trapped by reaction with benzophenone. Since heavy carbene



Scheme 1



Scheme 2


analogues containing electropositive substituents are likely to exhibit interesting chemistry,<sup>5</sup> we are examining methods to prepare precursors for transition metal substituted germynes,  $\{\text{:Ge}[\text{ML}_n]_2\}$ . 1,1-Disubstituted germacyclopent-3-enes seem to be suitable precursors since dimethylgermylene<sup>6</sup> and dichlorogermylene<sup>7</sup> have been cleanly extruded under relatively mild conditions.

A series of 1,1-ditransition metal substituted germacyclopent-3-enes have been prepared according to the metathesis reaction of Scheme 1. When reactions were carried out over short periods (30 min), mixtures of mono- and di-substituted products were obtained. However, the 1,1-disubstituted species, (1)–(4), were isolated in high yields after a reaction period of 12 h as colourless or pale yellow solids that gave satisfactory spectroscopic and elemental analytical data.†


Thermogravimetric analyses of (1)–(4) indicate that the onset of weight loss occurs at  $\sim 150^\circ\text{C}$  and is complete by  $\sim 250^\circ\text{C}$ .‡ After pyrolysis in the solid state at  $165^\circ\text{C}$  for five minutes in a sealed tube, (1) generated 2,3-dimethylbuta-1,3-diene as shown by  $^1\text{H}$  n.m.r. spectroscopy of the reaction products, implying the possible extrusion of  $\{\text{:Ge}[\text{Mn}(\text{CO})_5]_2\}$  (5). In order to determine whether transition metal substituted germynes are extruded intact, a number of trapping experiments have been carried out. Scheme 2 is typical of these experiments. Compound (4) was pyrolysed in the presence of a large excess of 2,3-dimethylbuta-1,3-diene diluted with toluene solution and sealed in an

† Selected spectroscopic data: (1)  $^1\text{H}$  n.m.r.  $\delta$  2.44 (s, 4H,  $\text{CH}_2$ ), 1.82 (s, 6H, Me); i.r.  $\nu(\text{C}-\text{O})$  ( $\text{cm}^{-1}$ ), 2100m, 2067m, 2032s, br., 1950sh. (2)  $^1\text{H}$  n.m.r.  $\delta$  2.48 (s, 4H,  $\text{CH}_2$ ), 2.06 (s, 6H, Me), 4.40 (s, 5H, Cp); i.r.  $\nu(\text{C}-\text{O})$  ( $\text{cm}^{-1}$ ), 1983s, 1933sh, 1917s, 1900sh. (3)  $^1\text{H}$  n.m.r.  $\delta$  2.47 (s, 4H,  $\text{CH}_2$ ), 1.66 (s, 6H, Me); i.r.  $\nu(\text{C}-\text{O})$  ( $\text{cm}^{-1}$ ) 2100s, 2081s, 2032s, 1944s, br. (4)  $^1\text{H}$  n.m.r.  $\delta$  2.30 (s, 2H,  $\text{CH}_{2a}$ ), 2.46 (s, 2H,  $\text{CH}_{2b}$ ), 1.88 (s, 3H, Me), 5.88 (s, 1H, CH); i.r.  $\nu(\text{C}-\text{O})$  ( $\text{cm}^{-1}$ ) 2075m, 2010s, 2001s, br., 1986s, br.

‡ Data were recorded from 40–400°C with a heating rate of 10°C/min and from 40–900°C with a heating rate of 40°C/min. No weight change was observed above 250°C.



Scheme 3



Scheme 4

ampoule *in vacuo*. After heating to  $185^\circ\text{C}$  for 5 min,§ the volatile components were removed *in vacuo* and the whole reaction mixture dissolved in  $[^2\text{H}_6]$ benzene. Methyl and methylenic  $^1\text{H}$  n.m.r. resonances corresponding to an authentic sample of (1) could be unequivocally identified, together with resonances due to unchanged (4) and a small amount of polydiene. After this short pyrolysis, at least 30% of (5), extruded from (4), had been trapped to form (1).¶ This experiment provides quantitative evidence that is consistent with the extrusion of the transition metal substituted germylene,  $\{\text{:Ge}[\text{Mn}(\text{CO})_5]_2\}$ , upon pyrolysis of (4) and demonstrates that both the Ge–Mn and Mn–CO bonds are kinetically stable to the reaction conditions employed. It has previously been reported<sup>8</sup> that compound (3) extrudes  $\{\text{:Ge}[\text{Co}(\text{CO})_4]_2\}$  between 40 and  $60^\circ\text{C}$ . However, we found that heating (3) for 6 h at  $70^\circ\text{C}$  resulted only in the quantitative extrusion of CO to form (6), according to Scheme 3. Elimination of 2,3-dimethylbuta-1,3-diene occurs at a significant rate only above  $170^\circ\text{C}$ .

Additional evidence for the generation of transition metal substituted germynes has been obtained from trapping reactions according to Scheme 4. Compound (7) was isolated in 30% yield after reaction for 12 h at room temperature in diethyl ether and was characterized by  $^1\text{H}$  n.m.r. spectroscopy.¶ Since the reaction of  $\text{GeI}_2$  with 2,3-dimethylbuta-1,3-diene is negligible over this period at room temperature (significant reaction occurs only  $\geq 50^\circ\text{C}$ ), and the reaction of  $\text{Na}[\text{Mo}(\text{CO})_3\text{Cp}]$  with 1,1-di-iodo-3,4-dimethylgermacyclopent-3-ene is only 20% complete after three days (control experiment using  $\text{Et}_2\text{O}$  as solvent under similar conditions to those of Scheme 4), we conclude that  $\{\text{:GeI}[\text{Mo}(\text{CO})_3\text{Cp}]\}$  (8) must be a relatively long-lived species under the conditions of

§ A twenty-fold excess of trapping reagent was used and the experiment carried out at high dilution [ $\sim 0.1$  mm in (4)] to reduce the possibility of polymerization of the dienes.

¶ This is the minimum extent of conversion as determined by comparison of the  $^1\text{H}$  n.m.r. integration of resonances due to unchanged (4) and (1). However, it is likely that more than 30% of (4) reacted.

|| (6),  $^1\text{H}$  n.m.r.  $\delta$  2.46 (s, 4H,  $\text{CH}_2$ ), 1.46 (s, 6H, Me); i.r.  $\nu(\text{C}-\text{O})$  ( $\text{cm}^{-1}$ ) 1999s, br.; 2041s, 2088s, 1836s. (7)  $^1\text{H}$  n.m.r.  $\delta$  2.57 (d, 2H,  $J_{\text{HH}}$  16 Hz,  $\text{CH}_{2a}$ ), 2.36 (d, 2H,  $\text{CH}_{2b}$ ), 1.68 (s, 6H, Me).

Scheme 4. The isolation of only the mono-substituted product is probably the result of the large steric demands of the  $[\text{Mo}(\text{CO})_3\text{Cp}]^-$  moiety.

Further studies on the thermal rearrangements of this interesting class of compounds and the reactivity of transition metal substituted germylenes are in progress.

We thank Dr. Dan Doughty of Sandia National Laboratories for helpful discussions, Eva Quesnell for assistance, and the Sandia-University Research Program for financial support of this project. Elemental analyses were performed by Guelph Chemical Laboratories.

Received, 15th February 1989, Com. 9/00720B

## References

- 1 P. P. Gaspar, in 'Reactive Intermediates,' eds. M. Jones and R. A. Moss, Wiley, New York, 1985, vol. 3, pp. 333—427, and references therein.
  - 2 D. Lei and P. P. Gaspar, *J. Chem. Soc., Chem. Commun.*, 1985, 1149; H. Vancik, G. Raabe, M. J. Michalczyk, R. West, and J. Michl, *J. Am. Chem. Soc.*, 1985, **107**, 4097.
  - 3 L. Linder, A. Revis, and T. J. Barton, *J. Am. Chem. Soc.*, 1986, **108**, 2742; B. H. Boo and P. P. Gaspar, *Organometallics*, 1986, **5**, 698.
  - 4 A. Marietti-Migrani and R. West, *Organometallics*, 1987, **6**, 141.
  - 5 For example, it has been suggested that silylenes, when substituted with electropositive substituents, may exhibit triplet electronic ground states, M. E. Colin, J. Breulet, and H. F. Schaefer, *Tetrahedron*, 1985, **41**, 1434.
  - 6 D. Lei and P. P. Gaspar, unpublished results.
  - 7 S. P. Kolesnikov, I. S. Royozhin, M. G. Safarov, and O. M. Nefedov, *Izv. Akad. Nauk, SSSR, Ser. Khim*, 1981, **6**, 1423, *Engl. transl.*, 1981, 1143. Dichlorogermylene was extruded at  $\sim 500^\circ\text{C}$ .
  - 8 We have isolated compound (3) as a colourless crystalline solid, according to the reaction of Scheme 1, and recrystallized it from n-pentane at  $-30^\circ\text{C}$ . The previous report of this compound refers to it as a yellow-brown oil that decomposes upon distillation at  $60^\circ\text{C}$  and  $5 \times 10^{-1}$  mmHg. A. Castel, P. Riviere, J. Satge, J. J. E. Moreau, and R. J. P. Corriu, *Organometallics*, 1983, **2**, 1498.
-