

Reaction of Nitro-alkenes with Iodotrimethylsilane: A New Method for the Conversion of Vinyl Nitrosteroids to Ketosteroids

Gireesh M. Singhal, Nalin B. Das, and Ram P. Sharma*

Natural Products Chemistry Division, Regional Research Laboratory, Jorhat-6, Assam, India

Iodotrimethylsilane generated *in situ* from chlorotrimethylsilane and sodium iodide effects the reduction of nitro-alkenes (**1a—e**) and (**5**) at -5 to 0°C to furnish the ketones (**2a—e**) and (**6**) respectively as the major products.

Since the trimethylsilyl cation (Me_3Si^+) is considered to be an equivalent of a proton in many of the reactions of halogenosilanes, it was argued that the reaction of chlorotrimethylsilane with 6-nitrocholest-5-ene (**1a**) would provide 5α -chloro-6-hydroxyiminocholestane as does the reaction of hydrochloric acid with (**1a**).¹ However, it failed to react with (**1a**) in refluxing dichloromethane, and bromotrimethylsilane was also unreactive towards (**1a**). The reaction of (**1a**) with iodotrimethylsilane generated *in situ* from $\text{Me}_3\text{SiCl}/\text{NaI}$ provided a complex mixture of products at room temperature. However, the reaction of (**1a**) with $\text{Me}_3\text{SiCl}/\text{NaI}$ at -5 to 0°C for 10 min followed by aqueous work-up gave one major and two minor products (t.l.c.) which were isolated and identified as (**2a**), (**3a**),² and (**4a**) respectively by direct comparison with authentic samples. The stereochemistry of the oxime (**4a**) was

established as *E* by n.m.r. analysis. This reaction was then studied with 3-substituted- Δ^5 -6-nitrocholestanes (**1b—e**) and in each case the major product (70—85% yield) was the




Table 1. Reductions with $\text{Me}_3\text{SiCl}/\text{NaI}$.^a

Substrate	Product ^b (% Yield)
(1a)	(2a) (75), (3a) (6), (4a) (8)
(1b)	(2b) (70), (3b) (8), (4b) (10)
(1c)	(2c) (80), (3c) (5), (4c) (6)
(1d)	(2d) (85), (3d) (2), (4d) (3)
(1e)	(2e) (77), (3e) (4), (4e) (8)

^a In a typical experiment, a solution of (**1a**) (0.5 mmol) in dry CH_2Cl_2 (4 ml) was treated with NaI (2.0 mmol) and Me_3SiCl (1.0 mmol) with stirring at -5 to 0°C for 10 min; the reaction was monitored by t.l.c. Aqueous work-up followed by t.l.c. on silica gel furnished (**2—4a**).

^b All compounds were characterized by i.r., n.m.r., and mass spectroscopy, and by direct comparison with authentic samples.

a; R = H
b; R = OAc
c; R = Cl
d; R = OMe
e; R = OH

corresponding 6-ketone (see Table 1). Similarly, 4-nitrocholest-4-ene on treatment with $\text{Me}_3\text{SiCl}/\text{NaI}$ gave cholestan-4-one (74%). Reaction of the nitro-alkene (**5**) prepared from the natural product tagitinin A³, with $\text{Me}_3\text{SiCl}/\text{NaI}$ furnished the ketone (**6**) in 78% yield.

Reaction of the oxime of (**2a**) with $\text{Me}_3\text{SiCl}/\text{NaI}$ under the same conditions did not furnish the ketone (**2a**); starting material was recovered unchanged, thus indicating that oximes are not intermediates in the formation of ketones. Since the conditions are mild, this method could be a useful alternative to the standard method (Zn/AcOH)⁴ for reducing the vinyl nitrosteroids to ketosteroids where refluxing conditions are required.

We have earlier reported⁵ that hydrogen iodide generated *in situ* from trifluoroacetic acid and NaI could be an equivalent of Me_3SiI . The reaction of nitro compound (**1a**) with $\text{CF}_3\text{CO}_2\text{H}/\text{NaI}$ did indeed furnish compounds (**2a**), (**3a**), and (**4a**) in 40, 20, and 25% yields respectively.

Olah *et al.*⁶ have reported that the reaction of primary, secondary, and tertiary nitro compounds with Me_3SiI furnishes the corresponding cyanides, oximes, and iodides respectively, and that the reaction of 1-nitrocyclohexane with

Me_3SiI yielded a complex mixture of products. We have obtained similar results.

We thank the Director, Dr. J. N. Baruah, for use of facilities.

Received, 12th April 1989; Com. 9/01516G

References

- 1 Y. Komeichi, S. Tomioka, T. Iwasaki, and K. Watanabe, *Tetrahedron Lett.*, 1970, 4677.
- 2 Hydroxy-ketones are also the byproducts in the nitration of alkenes: see, *e.g.*, C. R. Eck and B. Green, *J. Chem. Soc., Chem. Commun.*, 1972, 537, and references cited therein. Therefore, the purity of the nitro compounds used was fully ensured before use.
- 3 J. C. Sarma, N. C. Barua, R. P. Sharma, and J. N. Baruah, *Tetrahedron*, 1983, **39**, 2843.
- 4 L. F. Fieser and M. Fieser, 'Reagents for Organic Synthesis,' vol. 1, New York, 1967, p. 1276 (J. Mauthner and W. Suida, *Monatschfte*, 1894, **15**, 85; 1903, **24**, 648; A. Windans, *Berichte*, 1903, 36, 3752).
- 5 D. N. Sarma and R. P. Sharma, *Chem. Ind. (London)*, 1984, 712.
- 6 G. A. Olah, S. C. Narang, L. D. Field, and A. F. Fung, *J. Org. Chem.*, 1983, **48**, 2766.