

2,5-Bis(3,5-di-*t*-butyl-4-oxo-cyclohexa-2,5-dien-1-ylidene)-2,5-dihydro-1,4-dithiapentalene: A New Class of Amphoteric Multi-stage Redox Systems consisting of Both the Wurster and the Weitz Types

Kazuko Takahashi* and Masashi Ogiyama

Department of Chemistry, Faculty of Science, Tohoku University, Sendai 980, Japan

The dihydrodithiapentalene-extended diphenoquinone DTP-DQ (1) has been synthesized and characterized as a hitherto unknown amphoteric four-stage one-electron redox system involving both a two-stage one-electron redox reaction of the Weitz type and a two-stage one-electron redox reaction of the Wurster type.


Organic π -donors and π -acceptors undergoing stepwise transfer of single electrons have been classified into 'Weitz' or 'Wurster' types¹ depending on whether the end groups of the redox system are ring members or external groups of a cyclic π -system, respectively. There have appeared several reports on amphoteric hydrocarbons² and amphoteric heteroatom-containing compounds³ which are both anodically oxidized and cathodically reduced in a multi-stage redox fashion. However, there has so far been no report on closed shell amphoteric four-stage redox systems involving different redox-active units, in which a single molecule is both

anodically oxidized in a Weitz type manner and cathodically reduced in a Wurster type. This kind of redox system is of current interest in its possibilities for the creation of novel conducting complexes with three interacting components. 2,5-Bis(4-oxocyclohexa-2,5-dien-1-ylidene)-2,5-dihydro-1,4-dithiapentalene shows promise for such a hitherto unknown redox system since the corresponding radical ions and divalent ions could be significantly stabilized by the chalcogen atoms and the oxygen atoms which are incorporated as ring members (Weitz type) and as outside groups (Wurster type) of a cyclic π -system, respectively. With this in mind, we have now

Table 1. Oxidation and reduction potentials,^a and the difference between E_1 and E_2 of quinones (1), (8), and (9) (V vs. standard calomel electrode, SCE).

Compd.	Solvent	$E_{p2^{ox}}$	$E_{p1^{ox}}$	$E_{1^{ox}}$	$\Delta E_{p^{ox}}$	$E_{1^{red}}$	$E_{2^{red}}$	ΔE^{red}
(1)	PhCN	+1.59	+1.09	+1.04	0.50	-0.42	-0.55	0.13
(1)	CH ₂ Cl ₂	+1.66	+1.14	+1.06	0.52	-0.45 ^b	-0.55 ^b	0.10
(8)	CH ₂ Cl ₂					-0.52	-0.89	0.37
(9)	CH ₂ Cl ₂			+1.20		-0.46	-0.60	0.14

^a Obtained by cyclic voltammetry operated under the conditions given in the caption to Figure 1. ^b Divided from a coalescent peak according to the Myers-Shain method.⁵ E^{ox} and E^{red} values were calculated by averaging peak potentials of the reversible anodic and cathodic segments of the cycle; $E = (E_p^a + E_p^c)/2$.



value§ is far larger than the ΔE^{red} value in (1), and is even larger than the ΔE^{ox} of tetrathiafulvalene (TTF) (0.33 V). This clearly indicates the significantly increased Coulomb repulsion in (1²⁺) whose groups bearing the bulk of the positive charge are close to each other. Thus, the central sulphur atoms should function as the end groups of the Weitz-type oxidation sequence. The presence of the two central sulphur atoms in (1) might be an important prerequisite for the formation of (1²⁺), since the monosulphide (9) failed to give (9²⁺) and gave only (9^{•+}). From the ESR study, the bulk of the spin population in (1^{•+}) resides on the sulphur atoms, since the g value of (1^{•+}) (2.0055) is substantially larger than that of (1^{•-}) (2.0035). The significant difference in the g values between (1^{•+}) and (1^{•-}) can be fully appreciated from the prominent features of the HOMO and the LUMO calculated by the CNDO/2 method: the $2p\pi$ or $3p\pi$ AO coefficients are large at the oxygen atoms and almost vanish at the sulphur atoms in the LUMO; in contrast, they are very large at the sulphur atoms and C-3',6' in the HOMO.

Therefore, the amphoteric four-stage redox reaction of (1)

proceeds according to the sequence shown in Scheme 2, involving both two-stage one-electron reduction of the Wurster type and two-stage one-electron oxidation of the Weitz type. The central dithiapentalene ring may have pseudo 8π , 10π , and 6π electron character in (1), (1²⁻), and (1²⁺), respectively.

We thank Professor Yusaku Ikegami and Dr Kimio Akiyama for obtaining g values and for helpful discussions.

Received, 26th April 1990; Com. 0/01853H

References

- 1 K. Deichert and S. Hunig, *Angew. Chem., Int. Ed. Engl.*, 1978, **17**, 875; S. Hunig and H. Berneth, *Top. Curr. Chem.*, 1980, **92**, 1.
- 2 V. D. Parker, *J. Am. Chem. Soc.*, 1976, **98**, 98; O. Hammerich and V. D. Parker, *ibid.*, 1974, **96**, 4289; B. S. Jensen and V. D. Parker, *ibid.*, 1975, **97**, 5211; S. Hunig, M. Horner, and P. Schilling, *Angew. Chem., Int. Ed. Engl.*, 1975, **14**, 556; B. Hagenbruch, K. Hess, S. Hunig, and G. Klug, *Liebigs Ann. Chem.*, 1981, 256; K. Nakasuji, K. Yoshida, and I. Murata, *J. Am. Chem. Soc.*, 1982, **104**, 1432; 1983, **105**, 5136.
- 3 (a) S. Hunig and H. Putter, *Chem. Ber.*, 1977, **110**, 2524; (b) D. J. Sandman and A. F. Richter, *J. Am. Chem. Soc.*, 1979, **101**, 7079; (c) J. Heinz, *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 202; (d) K. Takahashi and T. Suzuki, *J. Am. Chem. Soc.*, 1989, **111**, 5483.
- 4 J. B. Torrance, *Acc. Chem. Res.*, 1979, **12**, 79.
- 5 R. L. Myers and I. Shain, *Anal. Chem.*, 1969, **41**, 980.

§ The second half-wave oxidation potential, E_2^{ox} , cannot be determined exactly from Figure 1, since the corresponding reduction segment of the cycle is missing. The Coulomb repulsion in (1²⁺) was therefore evaluated from the ΔE_p^{ox} value instead of ΔE^{ox} .