


**NEW
JOURNAL**

MENDELEEV COMMUNICATIONS

Preliminary Accounts of Important New Work
in Chemistry from the USSR and Elsewhere



- A prestigious new journal to be launched in January 1991
- The result of a collaborative venture between The Academy of Sciences of the USSR and The Royal Society of Chemistry
- Similar in scope and content to its highly successful 'sister' publication – Chemical Communications
- Primary publication in English of original Soviet Chemistry
- Two stages of rigorous refereeing to maintain the highest possible standards
- Rapid publication of papers – within three months of receipt in the UK
- News section, containing details of forthcoming international conferences, letters, and information about The Academy of Sciences of the USSR

For further information, please contact:

The Sales and Promotion Department, Royal Society of Chemistry,
Thomas Graham House, Science Park, Milton Road, Cambridge
CB4 4WF, United Kingdom.
Telephone: (0223) 420066. Fax: (0223) 423623. Telex: 818293 ROYAL

ROYAL
SOCIETY OF
CHEMISTRY



Information
Services



Lancaster Catalogue

89/90


UNITED KINGDOM
Lancaster Synthesis Ltd.
 Eastgate, White Lund,
 Morecambe, Lancs. LA3 3DY
 Freephone: 0800-262336
 Telephone: 0524-36101
 Fax: 0524-39727
 Telex: 65151 (LNCSYN G)

U.S.A. AND CANADA
Lancaster Synthesis Ltd.
 P.O. Box 1000,
 Windham,
 New Hampshire 03087
 Toll-free lines: 800-238-2324
 Telephone: 603-889-3306
 Fax: 603-889-3326

FRANCE
Lancaster Synthesis Ltd.
 15 rue de l'Atome Z.I.,
 67800 Bischheim,
 Strasbourg.
 Telephone: 88-62-26-90
 Fax: 88-62-26-81
 Telex: 870551 (LNCSTRM F)


GERMANY
Deutsche MTM Chemie GmbH.
 Lämmerspieler Straße 100A,
 6052 Mühlheim am Main,
 Telephone: 06108 73 019
 Fax: 06108 74 814

JAPAN
Hydrus Chemical Inc.
 Tomitaka Bldg. 8-1,
 Uchikanda 2-chome,
 Chiyoda-ku,
 Tokyo 101.
 Telephone: (03) 258-5031
 Fax: (03) 258-6535
 Telex: 2324032 (HYDRUS J)




Completely revised
 1152 pages
 1000 new items

6500 literature references to some 2000 items
 Illustrated by 1500 reaction flow-charts
 Semi-bulk and bulk quantity indications
 Extensive cross-referencing




Deprotonation by lithium diisopropylamide at low temperature gives the unstable azomethine ylide, which undergoes 1,3-dipolar addition even with unactivated alkenes, to give pyrrolidines: *J.Chem.Soc., Chem. Commun.*, 31 (1983):




Compare also N-methylmorpholine-N-oxide, 5957, p.710.

Nickel acetylacetonate hydrate
 [Nickel(II)2,4-pentanedionate hydrate]
 F.W. 274.94, m.p. ca 285°(dec), [3264-82-2]
HARMFUL / POSSIBLE CARCINOGEN
 Please ask for bulk prices (5Kg to 100Kg+)



Catalyst for a variety of useful coupling reactions, including:
 Conjugate addition of alkynylaluminium reagents to enones: *J.Am.Chem.Soc.*, 100, 2244 (1978):



Conjugate addition of cis-alkenylzirconium reagents, from the hydrozirconation of alkynes, to Michael acceptors, with retention of configuration: *J.Am.Chem.Soc.*, 102, 1333 (1980).

Coupling of Grignard reagents to give biaryls: *J.Org.Chem.*, 41, 2252 (1976).

Coupling of Grignard reagents with silyl enol ethers of both aldehydes and ketones, to give alkenes. In contrast to dichlorobis(triphenylphosphine)nickel, this reagent gives the thermodynamically more stable alkene: *Tetrahedron*, 36, 295 (1980):



SEND NOW
MAKE SURE YOU HAVE
YOUR PERSONAL COPY

LANCASTER SYNTHESIS - PART OF MTM RESEARCH CHEMICALS