


When you're in the STN EXPRESS® lane, you're not restricted to just "10 items or less."



Don't put another package in your cart until you've compared these features of STN Express Version 3.00:


- auto-logout to STN International, Questel, and other vendors
- structure modeling for searching databases on STN and Questel
- scrolling text and graphics
- freehand or template structure drawing
- searching for exact or related substances
- searching for preparations of substances
- searching for role of substance in a reaction
- prepackaged queries
- automatic normalization of aromatic and tautomeric bonds
- VERY friendly look and feel
- easy to use script language
- an "undo" feature

To find out just how easy it is to search in the STN Express lane, please complete and return the following coupon, or contact us as indicated.


Here are more than 10 fresh reasons to try searching STN International® with STN Express.

When you "shop" in the STN International online network you have access to 58 databases relating to chemistry and more than 50 other scientific and technical databases too. One of them, the CAS Registry file, contains CAS Registry Numbers® and more than 10 million structure diagrams!


With STN EXPRESS software for your IBM® PS/2, or Apple Macintosh®, you can build and search these structures more easily.



"Select "aromatics" to display this selection of templates. Simply use your mouse to select the template of interest. Now you can modify the structure to fit your specifications."



"STN Express Version 3.00 is compatible with other software."



"You get high-quality text and graphics display with STN Express."

Yes! I want more information about STN Express software.

Name _____

Job Title _____

Organization _____

Street _____

City _____ Country _____

Date _____ Phone _____

Signature _____

Enquirers from the UK and Eire, please mail to The Secretary, STN International, c/o Royal Society of Chemistry, Thomas Graham House, Science Park, Milton Road, Cambridge CB4 4WF.

STN INTERNATIONAL®



Lancaster Catalogue

89/90

UNITED KINGDOM

Lancaster Synthesis Ltd.
Eastgate, White Lund,
Morecambe, Lancs. LA3 3DY
Freephone: 0800-262336
Telephone: 0524-36101
Fax: 0524-39727
Telex: 65151 (LNC SYN G)

U.S.A. AND CANADA

Lancaster Synthesis Ltd.
P.O. Box 1000,
Windham,
New Hampshire 03087
Toll-free lines: 800-238-2324
Telephone: 603-889-3306
Fax: 603-889-3326

FRANCE

Lancaster Synthesis Ltd.
15 rue de l'Atome Z.I.,
67800 Bischheim,
Strasbourg.
Telephone: 88-62-26-90
Fax: 88-62-26-81
Telex: 870551 (LNCSTRM F)

GERMANY

Deutsche MTM Chemie GmbH.
Laemmerspieler Straße 100A,
6052 Muhlheim am Main.
Telephone: 06108 73 019
Fax: 06108 74 814


JAPAN

Hydrus Chemical Inc.
Tomitaka Bldg. 8-1,
Uchikanda 2-chome,
Chiyoda-ku,
Tokyo 101.
Telephone: (03) 258-5031
Fax: (03) 258-6535
Telex: 2324032 (HYDRUS J)




Completely revised
1152 pages
1000 new items

6500 literature references to some 2000 items
Illustrated by 1500 reaction flow-charts
Semi-bulk and bulk quantity indications
Extensive cross-referencing



... is converted to the Mannich reagent, N,N-dimethylacetamide, an excellent reagent for the α -dimethyl-
Bull.Soc.Chim.Fr., 2707 (1970). Compare dimethyl-
J.Chem.Soc., 0131, p.423:

Deprotonation by lithium diisopropylamide at low temperature gives the unstable azomethine ylide, which undergoes 1,3-dipolar addition even with unactivated alkenes, to give pyrrolidines: *J.Chem.Soc., Chem. Commun.*, 31 (1983):



Compare also N-methylmorpholine-N-oxide, 5957, p.710.

Nickel acetylacetonate hydrate


[Nickel(II)2,4-pentanedionate hydrate]
F.W. 274.94, m.p. ca 285° (dec), [3264-82-2]

HARMFUL / POSSIBLE CARCINOGEN

Please ask for bulk prices (5Kg to 100Kg+)

Catalyst for a variety of useful coupling reactions, including:


Conjugate addition of alkynylaluminium reagents to enones: *J.Am.Chem.Soc.*, 100, 2244 (1978):



Conjugate addition of cis-alkenylzirconium reagents, from the hydrozirconation of alkynes, to Michael acceptors, with retention of configuration: *J.Am.Chem.Soc.*, 102, 1333 (1980).

Coupling of Grignard reagents to give biaryls: *J.Org.Chem.*, 41, 2252 (1976).

Coupling of Grignard reagents with silyl enol ethers of both aldehydes and ketones, to give alkenes. In contrast to dichlorobis(triphenylphosphine)nickel, this reagent gives the thermodynamically more stable alkene: *Tetrahedron*, 36, 295 (1980):



		£
	50g	7.60
	250g	30.40

SEND NOW
MAKE SURE YOU HAVE
YOUR PERSONAL COPY

LANCASTER SYNTHESIS - PART OF MTM RESEARCH CHEMICALS