

## A Convenient Synthesis of (*E*)- $\beta$ -Ethoxycarbonylvinylsilanes by Palladium-catalysed Regio- and Stereo-specific Hydroesterification of Trimethylsilylacetylenes

Ryo Takeuchi,\* Masaharu Sugiura, Naomi Ishii and Nobuhiro Sato

Department of Chemistry, Yokohama City University, Kanazawa-ku, Yokohama 236, Japan

Palladium-catalysed hydroesterification of trimethylsilylacetylenes **1** gives (*E*)- $\beta$ -ethoxycarbonylvinylsilanes **2** exclusively in excellent yields.

Viny silanes have attracted much attention because of their synthetic utility.<sup>1</sup> Their electrophilic substitution reaction provides a useful procedure for the stereoselective synthesis. Viny silanes that bear the alkoxy carbonyl group on one of the vinylic carbons should be useful synthetic intermediates, because they contain both the viny silane and  $\alpha,\beta$ -unsaturated ester functionalities in the same molecule sharing the same carbon-carbon double bond.<sup>2</sup> Zweifel *et al.* reported the high yield and stereodefined synthesis of (*Z*)- $\alpha$ -ethoxycarbonylviny silanes *via* the stereo- and regio-specific hydroalumination of silylacetylene.<sup>2a</sup> Although (*E*)-3-trimethylsilylprop-2-enoic acid with the carboxy group on  $\beta$ -position in viny silane, the generality of the reaction is lacking.<sup>2d</sup> A general synthesis of viny silanes that bear the alkoxy carbonyl or carboxy group on the  $\beta$ -position of the vinylic carbons from easily accessible materials is desired. Transition metal complex catalysed carbonylation is a useful method for introducing a carbonyl functionality into organic compounds.<sup>3</sup> In our previous paper, we demonstrated the selective synthesis of  $\alpha$ - or  $\beta$ -silyl esters by the highly regioselective carbonylation of viny silanes.<sup>4</sup> We now report a general method for the synthesis of (*E*)- $\beta$ -ethoxycarbonylviny silanes **2** by the Pd-catalysed regio- and stereo-specific hydroesterification of easily accessible trimethylsilylacetylenes **1**.

The results are summarized in Table 1. The Pd-catalysed hydroesterification of **1** gave (*E*)- $\beta$ -ethoxycarbonylviny silanes **2** exclusively except for entry 1 [eqn. (1)]. PdCl<sub>2</sub>(dppf) combined with SnCl<sub>2</sub>·2H<sub>2</sub>O was the most effective catalyst






Table 1 Hydroesterification of trimethylsilylacetylenes **1**<sup>a</sup>


Entry	Substrate R	Product	Yield of <b>2</b> (%) <sup>b</sup>
1	<b>1a</b> H	<b>2a</b>	84 <sup>c</sup>
2	<b>1b</b> Bu <sup>n</sup>	<b>2b</b>	78
3	<b>1c</b> Oct <sup>n</sup>	<b>2c</b>	91
4	<b>1d</b> Cy <sup>d</sup>	<b>2d</b>	85
5	<b>1e</b> Ph	<b>2e</b>	88
6	<b>1f</b> 	<b>2f</b>	82
7	<b>1g</b> 	<b>2g</b>	90

<sup>a</sup> For conditions see text. <sup>b</sup> Isolated yield. <sup>c</sup> Determined by GLC. Ethyl 2-trimethylsilylprop-2-enoate was obtained in 1% yield. <sup>d</sup> Cy = cyclohexyl

with regard to the selectivity and the yield of the reaction [dppf = 1,1'-bis(diphenylphosphino)ferrocene];<sup>5</sup> addition of SnCl<sub>2</sub>·2H<sub>2</sub>O as a cocatalyst was essential for the reaction. Dicarboxylation or polymerization of **1** was not observed under the chosen conditions. The stereochemistry of **2** shows that the reaction proceeds in a *syn* manner.<sup>†</sup> A typical procedure is as follows. A mixture of trimethylsilylacetylene **1** (5 mmol), ethanol (10 ml), PdCl<sub>2</sub>(dppf) (0.1 mmol), and SnCl<sub>2</sub>·2H<sub>2</sub>O (0.5 mmol) was heated in a 50 ml stainless steel autoclave and stirred at 90 °C for 15 h under 20 kg cm<sup>-2</sup> of carbon monoxide. Products were isolated by distillations under reduced pressure. The silylacetylenes substituted by a primary or a secondary alkyl group as well as a phenyl group afforded the corresponding viny silanes in excellent yields (entries 2-5). This hydroesterification is also successful with trimethylsilylacetylenes bearing a functional group away from the triple bond (entries 6 and 7). The reduction of the carbon-chloro bond or ethoxycarbonyl group did not occur during the reaction.

The selectivity of the present reaction can be rationalized as follows. The hydropalladation of **1** proceeds in a *syn* manner to give complex **3** or **4**.<sup>6</sup> The steric repulsion between a trimethylsilyl substituent and ligands on palladium would yield **3** in preference to **4**. Insertion of carbon monoxide into the carbon-palladium bond would yield acylpalladium species. Subsequent nucleophilic attack by ethanol would yield the corresponding products.

As shown here, the present reaction provides an efficient route to (*E*)- $\beta$ -ethoxycarbonylviny silanes. Further application and mechanistic studies of the reaction are in progress.



<sup>†</sup> Selected spectroscopic data: Ethyl (*E*)-3-(trimethylsilyl)prop-2-enoate **2a**: <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>)  $\delta$  0.13 (s, 9H), 1.29 (t, *J* 7.25 Hz, 3H), 4.20 (q, *J* 7.25 Hz, 2H), 6.23 (d, *J* 18.80 Hz, 1H), 7.24 (d, *J* 18.80 Hz, 1H); <sup>13</sup>C NMR (67.8 MHz)  $\delta$  -2.03, 14.13, 60.33, 133.94, 149.31, 165.75; IR  $\nu$ /cm<sup>-1</sup> 1720.

Ethyl (*E*)-2-(*n*-butyl)-3-(trimethylsilyl)prop-2-enoate **2b**: <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>)  $\delta$  0.17 (s, 9H), 0.92 (t, *J* 7.26 Hz, 3H), 1.30 (t, *J* 7.26 Hz, 3H), 1.34-1.44 (m, 4H), 2.38 (t, *J* 7.59 Hz, 2H), 4.19 (q, *J* 7.26 Hz, 2H), 6.77 (s, 1H); <sup>13</sup>C NMR (67.8 MHz)  $\delta$  -0.38, 13.89, 14.20, 22.91, 31.90, 32.26, 60.61, 140.23, 148.23, 167.44 (<sup>3</sup>*J*<sub>C=O,H</sub> 9.8 Hz); IR  $\nu$ /cm<sup>-1</sup> 1715.

Ethyl (*E*)-2-(phenyl)-3-(trimethylsilyl)prop-2-enoate **2e**: <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>)  $\delta$  -0.12 (s, 9H), 1.25 (t, *J* 7.26 Hz, 3H), 4.20 (q, *J* 7.26 Hz, 2H), 7.17-7.23 (m, 3H), 7.29-7.34 (m, 3H); <sup>13</sup>C NMR (67.8 MHz)  $\delta$  -0.76, 14.16, 61.10, 127.58, 127.62, 129.34, 138.33, 144.91, 147.87, 166.76 (<sup>3</sup>*J*<sub>C=O,H</sub> 9.8 Hz); IR  $\nu$ /cm<sup>-1</sup> 1720.

Ethyl (*E*)-2-(3-chloropropyl)-3-(trimethylsilyl)prop-2-enoate **2f**: <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>)  $\delta$  0.20 (s, 9H), 1.31 (t, *J* 7.26 Hz, 3H), 1.87-1.98 (m, 2H), 2.54 (m, 2H), 3.57 (t, *J* 6.59 Hz, 2H), 4.20 (q, *J* 7.26 Hz, 2H), 6.89 (s, 1H); <sup>13</sup>C NMR (67.8 MHz)  $\delta$  -0.45, 14.16, 29.65, 32.65, 44.89, 60.77, 142.25, 146.09, 166.86 (<sup>3</sup>*J*<sub>C=O,H</sub> 9.7 Hz); IR  $\nu$ /cm<sup>-1</sup> 1710.

We are grateful for financial support from a Grant-in aid for Science Research (03750621) from the Ministry of Education, Science and Culture of Japan and grants in support of the promotion of research at Yokohama City University.

Received, 27th May 1992; Com. 2/02764J

### References

- 1 E. W. Colvin, *Silicon Reagents in Organic Synthesis*, Academic Press, London, 1988, p. 7; E. W. Colvin, *Silicon in Organic Synthesis*, Butterworths, London, 1981, p. 44; W. P. Weber, *Silicon Reagents for Organic Synthesis*, Springer-Verlag, Berlin, 1983, p. 79; I. Fleming, *Chem. Soc. Rev.*, 1981, **10**, 83; T. H. Chan and I. Fleming, *Synthesis*, 1979, 761.
  - 2 (a) M. R. Najafi, M. Wang and G. Zweifel, *J. Org. Chem.*, 1991, **56**, 2468; (b) M. P. Cooke, *J. Org. Chem.*, 1987, **52**, 5729; (c) R. L. Danheiser and H. Sard, *J. Org. Chem.*, 1980, **45**, 4810; (d) R. F. Cunico and F. J. Clayton, *J. Org. Chem.*, 1976, **41**, 1480.
  - 3 H. M. Colquhoun, J. Holton, D. J. Thompson and M. V. Twigg, *New Pathways for Organic Synthesis*, Plenum, New York, 1984, p. 195.
  - 4 R. Takeuchi, N. Ishii and N. Sato, *J. Chem. Soc., Chem. Commun.*, 1991, 1247; R. Takeuchi, N. Ishii, M. Sugiura and N. Sato, *J. Org. Chem.*, 1992, **57**, 4189.
  - 5 PdCl<sub>2</sub>(dppf) was prepared by the reaction of PdCl<sub>2</sub>(PhCN)<sub>2</sub> and 1,1'-bis(diphenylphosphino)ferrocene, see: T. Hayashi, M. Konishi, Y. Kobori, M. Kumada, T. Higuchi and K. Hirotsu, *J. Am. Chem. Soc.*, 1984, **106**, 158.
  - 6 J. F. Knifton, *J. Mol. Catal.*, 1977, **2**, 293; J. Tsuji, *Acc. Chem. Res.*, 1969, **2**, 144.
-