Corrigenda

Novel Photocurrent Rectification Behaviour for a Photoconductive Cell using the Mesogenic 5,10,15,20-Tetrakis(4-n-pentadecylphenyl)porphyrin

Yo Shimizu, Atsushi Ishikawa, Shigekazu Kusabayashi, Masaru Miya and Akira Nagata

J. Chem. Soc., Chem. Commun., 1993, 656.

The correct structure for C15TPP (n = 15), on page 657, is shown below.





New Classes of Anion Receptor containing Charged and Neutral Transition Metal Lewis Acidic Recognition Sites

Paul D. Beer, Christian A. P. Dickson, Nicholas Fletcher, Alistair J. Goulden, Alan Grieve, Jana Hodacova and Trevor Wear

J. Chem. Soc., Chem. Commun., 1993, 828.

The correct structure for L^7 , in Scheme 5 on page 829, is shown below.



trans Ring-fused Bicyclic Structures by 6-*exo*-Trigonal Radical Closure: Use of the Triethylborane–Stannane–Air System for Intramolecular Radical Addition to Aldehydes Derrick L. J. Clive and Maarten H. D. Postema

J. Chem. Soc., Chem. Commun., 1993, 429.

On page 429, right hand column, line 11:

'... the equatorial alcohol was the major (ca. 65% of total) product.' should read '... the axial alcohol was the major (ca. 65% of total) product.'