

Fluxional Ligand Migrations in Triosmium Clusters containing 2-Pyridylphosphines

Antony J. Deeming* and Martin B. Smith

Department of Chemistry, University College London, 20 Gordon Street, London, UK WC1H 0AJ

The cluster $[\text{Os}_3(\text{CO})_{10}(\mu\text{-L})]$ ($\text{L} = \text{Ph}_2\text{PC}_5\text{H}_4\text{N}$) (X-ray structure reported), containing the diaxially coordinated 2-pyridyldiphenylphosphine bridge, undergoes rapid transfer of the pyridine between two metal atoms while the phosphorus atom remains firmly coordinated to the third.

The ligand 2-pyridyldiphenylphosphine ($\text{Ph}_2\text{PC}_5\text{H}_4\text{N}$, L) has been widely used to bridge metal centres and an extensive coordination and organometallic chemistry based on it has been developed.¹ It is generally reckoned to stabilize dinuclear compounds by being firmly attached to the metal atoms it bridges. We have now shown that the ligand L reacts with $[\text{Os}_3(\text{CO})_{10}(\text{MeCN})_2]$ in dichloromethane solution at room temperature to give orange crystals of the cluster $[\text{Os}_3(\text{CO})_{10}(\mu\text{-L})]$ **1** (45%) and yellow crystals of $[\text{Os}_3(\text{CO})_{11}\text{L}]$ **2** (21%) which were separated by TLC on silica from other products including $[\text{Os}_3(\text{CO})_{10}\text{L}_2]$. The ^1H NMR spectrum† shows that **2** contains the ligand L coordinated

through the phosphorus atom only [6-H pyridyl resonance at δ 8.87 compared with δ 8.72 for the free ligand] and its IR spectrum† is very similar to that of $[\text{Os}_3(\text{CO})_{11}(\text{PPh}_3)]$ indicating that they are isostructural.²

The cluster **1** is very interesting because of its crystal structure and fluxional behaviour in solution. Its single-crystal X-ray structure‡ (Fig. 1) establishes that the N and P atoms are both coordinated axially, consistent with the non-equivalence of the Ph groups as observed in the ^1H NMR spectra† between -20 and -90 °C, a result that is inconsistent with diequatorial coordination. This stereochemistry contrasts with that of $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-dppm})]$ and $[\text{Os}_3(\text{CO})_9(\mu\text{-dppm})(\eta^1\text{-dppm})]$ (dppm = $\text{Ph}_2\text{PCH}_2\text{PPh}_2$), e.g. in which bridging dppm ligands are diequatorially coordinated.^{3,4} Normally ligands bulkier than hydride⁵ or acetontirile⁶ are equatorially coordinated and we believe that the small span of L ($\text{N}\cdots\text{P} = 2.650$ Å)⁷ compared with that of dppm ($\text{P}\cdots\text{P} = 2.970$ Å)⁸ favours diaxial coordination. The $\text{N}\cdots\text{P}$ distance in **1** is 2.716 Å whereas the $\text{P}\cdots\text{P}$ distance in $[\text{Ru}_3(\text{CO})_{10}(\mu\text{-dppm})]$ is 3.127 Å.³ These distances might be compared with the

† Spectroscopic data for compound **1**: IR $\nu(\text{CO})/\text{cm}^{-1}$ (cyclohexane): 2083s, 2032vs, 2017vs, 2004s, 1977s, 1969s, 1955m, 1938w; ^1H NMR (400 MHz, CD_2Cl_2 , -60 °C): 2-pyridyl: δ 9.42 (6-H), 7.55 (4-H), 7.30 (5-H), 7.25 (3-H); phenyl: δ 7.50, 7.45, 7.09; (1,1,2,2- $\text{C}_2\text{D}_2\text{Cl}_4$, 80 °C): 2-pyridyl: δ 9.41 (6-H), 7.51 (4-H), 7.22 (5-H, 3-H); phenyl: δ 7.45, 7.33 (signals at δ 7.61 and 7.09 at 20 °C have coalesced); $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, CD_2Cl_2 , 20 °C), δ 5.6 (s).

Compound **2**, IR $\nu(\text{CO})/\text{cm}^{-1}$ (cyclohexane): 2108m, 2053s, 2042m, 2020vs, 1999m, 1987m, 1971m; ^1H NMR (400 MHz, CDCl_3 , 20 °C): 2-pyridyl: δ 8.87 (6-H), 7.57 (5-H), 7.27 (4-H), 7.09 (3-H); phenyl: δ 7.40–7.50; $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, CDCl_3 , 20 °C): 4.2(s).

Compound **3**: IR $\nu(\text{CO})/\text{cm}^{-1}$ (cyclohexane): 2083s, 2034vs, 2018vs, 2003m, 1977s, 1968s, 1952w, 1939w; ^1H NMR (400 MHz, CD_2Cl_2 , -20 °C): N-bonded 2-pyridyl: δ 9.40 (6-H), 7.91 (3-H), 7.59 (4-H), 7.30 (5-H), N-non-bonded 2-pyridyl: δ 8.70, 8.59 (6-H), 8.03, 7.88 (5-H), 7.74, 7.39, 7.29, 7.19 (3-H, 4-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, CD_2Cl_2 , 20 °C): δ 12.7(s).

Compound **4**: IR $\nu(\text{CO})/\text{cm}^{-1}$ (cyclohexane): 2083s, 2033vs, 2018vs, 2003m, 1978s, 1969s, 1952w, 1938w; ^1H NMR (400 MHz, CD_2Cl_2 , 20 °C): two isomers: N-bonded 2-pyridyl: δ 9.49 and 9.37 (6-H); N-non-bonded 2-pyridyl: δ 8.72 and 8.67 (6-H); other resonances at δ 8.12, 8.03, 7.86, 7.81, 7.62, 7.51, 7.41, 7.35, 7.27, 7.19, 7.01; $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, CD_2Cl_2 , 20 °C): δ 11.0 (s) and 8.4 (s) (ca. 1:1 ratio).

‡ Structure determination: Nicolet R3v/m diffractometer, Mo-radiation ($\lambda = 0.71073$ Å), room temp. data corrected empirically for absorption. Orange crystal, $\text{C}_{27}\text{H}_{14}\text{NO}_{10}\text{POs}_3$, $M = 1113.99$ g mol⁻¹, size = $0.40 \times 0.35 \times 0.30$ mm³, monoclinic, $P2_1/n$, $a = 9.278(4)$, $b = 25.508(7)$, $c = 13.096(6)$ Å, $\beta = 107.31(4)^\circ$, $U = 2959(2)$ Å³, $Z = 4$, $D_c = 2.50$ g cm⁻³, $\mu(\text{Mo-K}\alpha) = 129.7$ cm⁻¹, $F(000) = 2024$. Direct methods, $R = 0.0556$, $R' = 0.0545$, where $R' = [\sum w(|F_o| - |F_c|)^2 / \sum |F_o|^2]$, after refinement 379 parameters using 4107 data with $I_o > 1.5\sigma(I_o)$ in the range $5 \leq 2\theta \leq 50^\circ$. All non-H atoms were refined anisotropically with H-atoms included in idealised positions ($\text{C-H } 0.96$ Å, $U = 0.08$ Å³). Atomic coordinates, bond lengths and angles, and thermal parameters for both structures have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.




Fig. 1 Molecular structure of $[\text{Os}_3(\text{CO})_{10}(\mu\text{-Ph}_2\text{PC}_5\text{H}_4\text{N})]$, cluster **1**. Selected bond lengths (Å) and angles ($^\circ$): Os(1)–Os(2), 2.827(2); Os(2)–Os(3), 2.872(2); Os(1)–Os(3), 2.929(2); Os(1)–P, 2.380(4); Os(2)–N, 2.22(1); P–C(41), 1.85(2); N–C(41), 1.32(2); Os(1)–C(11), 1.92(2); Os(2)–C(21), 1.87(2); Os(3)–C(31), 2.01(2); Os(3)–C(34), 1.94(2); av. Os–C (equatorial), 1.894; Os(1)–Os(2)–N, 93.2(4); Os(2)–Os(1)–P, 81.9(1); Os(1)–P–C(41), 117.3(5); Os(2)–N–C(41), 125(1); N–C(41)–P, 117(1).





Fig. 2 400 MHz ^1H NMR spectra of the cluster $[\text{Os}_3(\text{CO})_{10}\{\mu\text{-P}(\text{C}_5\text{H}_4\text{N})_3\}]$ recorded in CD_2Cl_2 (-20 and 20 $^\circ\text{C}$) or in $1,1,2,2\text{-C}_2\text{D}_2\text{Cl}_4$ (80 $^\circ\text{C}$). Signals for the 6-H protons of the pyridyl rings are indicated for the free and coordinated groups; * indicates peaks for decomposition products at 80 $^\circ\text{C}$.

shortest axial–axial C...C distances (av = 2.875 Å) and the shortest equatorial–equatorial distances (av = 3.424 Å) in $[\text{Os}_3(\text{CO})_{12}]$ (calculated from data given in ref. 5). Dppm is a more flexible ligand and has been found to span metal–metal distances from 2.1 to 4.4 Å,⁹ while $\text{Ph}_2\text{PC}_5\text{H}_4\text{N}$ has never been found associated with a metal–metal distance greater than about 2.9 Å.

The ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra† of **1** are temperature dependent. At 20 $^\circ\text{C}$ sharp 2-pyridyl ^1H NMR resonances are observed while the phenyl resonances are very broad (line-widths up to 30 Hz). Lower temperatures lead to better resolution whereas higher temperatures lead to coalescence of peaks at δ 7.61 and 7.09 to a new broad resonance at δ 7.33. We have not been able to assign these phenyl signals with confidence but the observed changes are consistent with dynamic exchange of the two non-equivalent phenyl groups leading to a time-averaged plane of symmetry.



Scheme 1 Possible mechanisms for $\text{R}^a\text{-R}^b$ exchange in compounds **1**, **3** and **4**

The corresponding tri(2-pyridyl)phosphine complex $[\text{Os}_3(\text{CO})_{10}\{\mu\text{-P}(\text{C}_5\text{H}_4\text{N})_3\}]$ **3**, prepared similarly, gives more easily interpreted ^1H NMR spectra. At -20 $^\circ\text{C}$ there are three sharp sets of 2-pyridyl resonances [6-H resonances: δ 8.70 and 8.59 for the non-coordinated groups and δ 9.40 for the coordinated group] (Fig. 2). The pairwise coalescence of the non-coordinated 2-pyridyl resonances without the coordinated 2-pyridyl resonances changing very much shows that the exchange process does not involve the formation of monodentate *P*-bonded tri(2-pyridyl)phosphine which would lead to the mutual exchange of all three groups. Therefore we propose that R^a exchanges with R^b while the N and P atoms remain coordinated (Scheme 1). Processes A, B and C could account for the coalescences but we feel that process C is less likely than the others. Process A corresponds to a classical CO rearrangement process but with the pyridine and a CO ligand migrating *via* bridging positions instead of two CO ligands. However, a phosphine migration is an alternative and has been observed in triplatinum systems.¹⁰

We examined ^{13}C NMR spectra of cluster **1**, hoping to discriminate between these processes. A set of ten ^{13}C carbonyl resonances consistent with the static structure is observed for a CD_2Cl_2 solution at -85 $^\circ\text{C}$ but there are fast processes leading to CO exchange which prevent our specifically identifying the CO exchanges corresponding to $\text{R}^a\text{-R}^b$ exchange. At room temperature nine of the ten resonances have coalesced to a broad resonance at δ 183.5 while that for the CO ligand *trans* to the PPh_2 group remains as a doublet [δ 184.2 (d, $J = 4.5$ Hz)]. We suggest that process A together with other processes including a merry-go-round process in the M_3 plane over all three metal atoms as identified for $[\text{M}_3(\text{CO})_{10}(\mu\text{-}1,2\text{-diazine})]$ ($\text{M} = \text{Ru}$ or Os),^{11,12} also containing diaxially coordinated ligands, would lead to just the one CO ligand not exchanging. Process B or C could not account for this observation. Final confirmation that the P-atom remains bonded to the same Os atom throughout came from the observation of sharp ^{187}Os satellites^{13,14} to the $^{31}\text{P}\{^1\text{H}\}$ NMR singlet [$J(^{187}\text{Os}^{31}\text{P})$ 165 Hz] at temperatures from -40 $^\circ\text{C}$ up to 50 $^\circ\text{C}$. Any process involving the migration of the P-atom between the Os sites would have led to the collapse of the ^{187}Os satellites. Only process A, or the very unlikely process C, would be consistent with the observed $^{31}\text{P}\{^1\text{H}\}$ NMR spectra.

The corresponding cluster $[\text{Os}_3(\text{CO})_{10}\{\mu\text{-PhP}(\text{C}_5\text{H}_4\text{N})_2\}]$ **4** from di(2-pyridyl)phenylphosphine shows two isomers both in

the ^1H and ^{31}P NMR spectra as expected but a rapid equilibrium between them at room temperature leads to broad resonances. This is completely consistent with what we have concluded for the clusters **1** and **3**.

We thank the SERC for support and for funds towards the purchase of a diffractometer and the University of London Central Research Fund for support.

Received, 11th February 1993; Com. 3/00852E

References

- 1 C. G. Arena, E. Rotondo, F. Faraone, M. Lanfranchi and A. Tiripicchio, *Organometallics*, 1991, **10**, 3877; S. Lo Schiavo, E. Rotondo, G. Bruno and F. Faraone, *Organometallics*, 1991, **10**, 1613; E. Rotondo, S. Lo Schiavo, G. Bruno, C. G. Arena, R. Gobetto and F. Faraone, *Inorg. Chem.*, 1989, **28**, 2944; J. L. Roustan, N. Ansari and F. R. Ahmed, *Inorg. Chim. Acta*, 1987, **129**, L11; Z.-Z. Zhang, H.-K. Wang, H.-G. Wang and R.-J. Wang, *J. Organomet. Chem.*, 1986, **314**, 357; T. J. Barder, F. A. Cotton, G. L. Powell, S. M. Tetrick and R. A. Walton, *J. Am. Chem. Soc.*, 1984, **106**, 1323; J. P. Farr, M. M. Olmstead, N. M. Rutherford, F. E. Wood and A. L. Balch, *Organometallics*, 1983, **2**, 1758; J. P. Farr, M. M. Olmstead, F. E. Wood and A. L. Balch, *J. Am. Chem. Soc.*, 1983, **105**, 792.
 - 2 M. I. Bruce, D. C. Kehoe, J. G. Matison, B. K. Nicholson, P. H. Rieger and M. L. Williams, *J. Chem. Soc., Chem. Commun.*, 1982, 442; B. F. G. Johnson, J. Lewis and D. A. Pippard, *J. Chem. Soc., Dalton Trans.*, 1981, 407.
 - 3 A. W. Coleman, D. F. Jones, P. H. Dixneuf, C. Brisson, J.-J. Bonnet and G. Lavigne, *Inorg. Chem.*, 1984, **23**, 952.
 - 4 S. Cartwright, J. A. Clucas, R. H. Dawson, D. F. Foster, M. M. Harding and A. K. Smith, *J. Organomet. Chem.*, 1986, **302**, 403.
 - 5 M. R. Churchill and B. G. DeBoer, *Inorg. Chem.*, 1977, **16**, 878.
 - 6 P. A. Dawson, B. F. G. Johnson, J. Lewis, J. Puga, P. R. Raithby and M. J. Rosales, *J. Chem. Soc., Dalton Trans.*, 1982, 233.
 - 7 Calculated from data given by J.-P. Charland, J.-L. Roustan and N. Ansari, *Acta Crystallogr., Sect. C*, 1989, **45**, 680.
 - 8 Calculated from data given by H. Schmidbaur, G. Reber, A. Schier, F. E. Wagner and G. Müller, *Inorg. Chim. Acta*, 1988, **147**, 143.
 - 9 A. L. Balch, R. R. Guimerans, J. Linehan and F. E. Wood, *Inorg. Chem.*, 1985, **24**, 2021.
 - 10 A. M. Bradford, G. Douglas, L. Manojlović-Muir, K. W. Muir and R. J. Puddephatt, *Organometallics*, 1990, **9**, 409.
 - 11 F. A. Cotton, B. E. Hanson and J. D. Jamerson, *J. Am. Chem. Soc.*, 1977, **99**, 6588.
 - 12 F. A. Cotton and B. E. Hanson, *Inorg. Chem.*, 1977, **16**, 2820.
 - 13 M. A. Gallop, B. F. G. Johnson and J. Lewis, *J. Chem. Soc., Chem. Commun.*, 1987, 1831.
 - 14 R. Benn, H. Brenneke, E. Jousen, H. Lehmkuhl and F. López Ortiz, *Organometallics*, 1990, **9**, 756.
-