


Synthesis and Reactivity of Allenylvinyl Cycloalk-2-enyl Ethers

Jean-Pierre Dulcère, Valérie Agati and Jean Rodriguez

Faculté des Sciences et Techniques de St Jérôme, Laboratoire de Synthèse Organique URA CNRS 1411, av. Esc. Normandie-Niemen, D 12, F-13397 Marseille Cedex 20, France

Cohalogenation of alkenes **6** by *N*-bromosuccinimide (NBS) in 3-methylpent-2-en-4-yn-1-ol provides β -bromo-3-methylpent-2-en-4-ynyl ether **7** precursors of 3-(penta-1,3,4-trienyloxy)cycloalkenes **9** which react chemo- and regio-selectively with NBS in methanol to give bromovinyl acetals **10**.

In connection with our continuing studies on the reactivity and synthetic application of allenyl ethers,¹ we required to prepare compounds incorporating the structure of penta-1,3,4-trienyl ethers of type **5** (Scheme 1). Although extensive work on the synthesis of the corresponding isomeric penta-1,2,4-trienyl ethers has been reported,² the preparation of compound **5** has received only a little attention. Allenyl silyl enol ethers³ or trifluoromethylsulfonylethers⁴ can be obtained by 1,6-addition of lithium dimethylcuprate to acceptor-substituted enynes followed by treatment with the appropriate electrophile. Other procedures involve the ring cleavage of 2-furfuryl carbanions⁵ or the [2 + 2] cycloaddition of ynamines with silyl vinyl ketenes.⁶ However, as yet, general methods for the synthesis of these derivatives are not available. In the course of mechanistic studies, we observed that β -allenic aldehydes **4** can easily be obtained by treatment of pent-2-en-4-yn-1-ol **1** with excess of BuLi (4 equiv.), quenching with ClSiMe₃ (4 equiv.) and then acidic work up. We showed that this transformation evolved via the allenyl silyl enol ether **3**, since treating 1-methoxy-3-methylpent-2-en-4-yne **2** with 0.5 equiv. of Bu^tOK in dimethyl sulfoxide (DMSO) provided 1-methoxy-3-methylpenta-1,3,4-triene **5**[†] in 92% yield. The easy base-catalysed isomerization of **2** prompted us to prepare and study the reactivity of β -bromo-pent-2-en-4-ynyl ethers **7** (Scheme 2). These compounds were obtained in 60–72% yields by halogenation of alkenes **6** by *N*-bromosuccinimide (NBS) in 3-methylpent-2-en-4-yn-1-ol at –30 °C.‡ Selective dehydrohalogenation of **7** in diazabicycloundecene (DBU) led to allyl pent-2-en-4-ynyl ethers **8**, which under isomerization conditions (Bu^tOK, 0.5 equiv.) gave allyl vinyl ethers **9** (60–94%). Interestingly, halogenation of **9** by NBS in methanol in the presence of K₂CO₃ (1 equiv.) proceeded chemo- and regio-selectively by 1,4-addition to afford the sensitive bromovinyl acetals **10**. These compounds were obtained in good yields, providing the HBr was neutralized as




Scheme 1 Reagents and conditions: i, BuLi, ClSiMe₃ (4 equiv.); ii, 5% HCl; iii, Bu^tOK (0.5 equiv., DMSO, 15 °C, 0.5 h)

[†] **5**: IR, ν/cm^{-1} 2940, 1945, 1635 and 1140; NMR (200 MHz, CDCl₃): ¹H, δ 6.40 (d, 1H, *J* 12.9 Hz), 5.36 (d, 1H, *J* 12.9 Hz), 4.75 (q, d, 2H, *J* 2.9 Hz), 3.56 (s, 3H) and 1.75 (t, 3H, *J* 2.9 Hz); ¹³C δ 209.6, 146.7, 102.5, 96.6, 74.1, 55.6 and 15.4.

‡ Satisfactory analytical and spectral data were obtained for all compounds.

soon as it was generated. Furthermore, Bu^tOK promoted dehydrohalogenation of **10** resulted in the formation of allyl 2-en-4-ynyl mixed acetals **11**.§ The overall sequence isomerization–halogenation–dehydrohalogenation provides an efficient conversion of ethers **8** into the corresponding acetals **11**.

An alternative route to **11b–c** involves the base-catalysed isomerization of **7b–c** (Bu^tOK, 0.3 equiv.) into β -bromo



Scheme 2 Reagents and conditions: i, NBS (–30 °C, 3 h); ii, DBU, (110 °C, 3 h); iii, Bu^tOK (0.5 equiv., benzene, 60 °C, 1 h); iv, NBS, MeOH, K₂CO₃ (1 equiv., –40 °C, 1 h); v, Bu^tOK (1.2 equiv., 20 °C, 12 h); vi, Bu^tOK (0.3 equiv., pentane, 25 °C, 12 h)

§ **10b** (mixture of isomers): IR ν/cm^{-1} 3300, 3040, 2940, 2100, 1645, 1130 and 1040; NMR (200 MHz, CDCl₃): ¹H δ 5.85 (d, 1H, *J* 6.4 Hz), 5.78 (m, 2H), 5.30, 5.21 (2d, 1H, *J* 6.4 Hz), 4.11 (m, 1H), 3.25, 3.24 (2s, 3H), 2.80 (s, 1H), 1.83 (s, 3H) and 1.92–1.18 (m, 6H); ¹³C, 135.5, 135.4, 131.0, 130.8, 128.1, 127.5, 86.0, 76.0, 69.0, 51.7, 51.3, 29.9, 28.8, 24.93, 24.89, 19.1, 19.0 and 17.8.

allenyl vinyl ethers **12b-c**. 1,4-Halogenation of **12b-c** by NBS in methanol provided **13b-c**, precursors of β -bromo mixed acetals **14b-c** by selective dehydrohalogenation. Bis allyl mixed acetals **11b-c** were isolated subsequently upon treatment with DBU.

Our results show that the combination of the halogenation of alkenes and the base-catalysed isomerization of pent-2-en-4-ynyl ethers provide a novel, general and efficient access to a new class of polyunsaturated derivatives which can be used as building blocks in organic synthesis.⁷

Received, 29th March 1993; Com. 3/01784B

References

- 1 J. P. Dulcère, M. N. Mihoubi and J. Rodriguez, *J. Chem. Soc., Chem. Commun.*, 1988, 237; J. P. Dulcère and J. Crandall, *J. Chem. Soc., Chem. Commun.*, 1990, 561; J. P. Dulcère, V. Agati and R. Faure, *J. Chem. Soc., Chem. Commun.*, 1993, 270.
- 2 H. A. Selling, J. A. Rompes, P. P. Montijn, S. Hoff, J. H. Van Boom, L. Brandsma and J. P. Arens, *Rec. Trav. Chim. Pays-Bas*, 1969, **88**, 119; J. P. C. M. Van Dongen, A. J. De Jong, H. A. Selling, P. P. Montijn, J. M. Van Boom and L. Brandsma, *Recl. Trav. Chim. Pays-Bas*, 1967, **86**, 1077. A. J. Bartlett, T. Laird and W. D. Ollis, *J. Chem. Soc., Perkin Trans. 1*, 1975, 1315; *J. Chem. Soc., Chem. Commun.*, 1974, 496; L. Brandsma and H. D. Verkruijine, in *Studies in Organic Chemistry, Synthesis of Acetylenes, Alkenes and Cumulenes*, Elsevier, Amsterdam, 1981, p. 82.
- 3 S. Arndt, G. Handke and N. Krause, *Chem. Ber.*, 1993, **126**, 251.
- 4 S. H. Lee, M. J. Shih and M. Hulce, *Tetrahedron Lett.*, 1992, **33**, 185; M. Cheng and M. Hulce, *J. Org. Chem.*, 1990, **55**, 964.
- 5 K. Atsumi and I. Kuwajima, *J. Am. Chem. Soc.*, 1979, **101**, 2208; K. Takanischi, H. Urabe and I. Kuwajima, *Tetrahedron Lett.*, 1987, **28**, 2281.
- 6 K. H. Dotz, J. Muhlemeier and B. Trenkle, *J. Organomet. Chem.*, 1985, **289**, 257.
- 7 R. Zimmer, *Synthesis*, 1993, 165.