

A Novel Synthesis of (Oxodimethylenemethane)-palladium and -platinum Complexes

Akihiro Ohsuka, Takanobu Fujimori, Toshikazu Hirao, Hideo Kurosawa* and Isao Ikeda*

Department of Applied Chemistry, Faculty of Engineering, Osaka University, Suita, Osaka 565, Japan

Treatment of π -allyl-palladium and -platinum complexes bearing a methoxymethoxy group at the 2-position with base give the corresponding oxodimethylenemethane complexes, in contrast to the formation of 2-hydroxy-substituted π -allyl complexes *via* acidic hydrolysis.

Functionalization of π -allylpalladium intermediates is envisaged to widen their synthetic utility. The π -allyl complex **2a** bearing an acetal function, methoxymethoxy group, at the 2-position seems to possess versatile reactive sites. This complex has been considered to be involved as a reaction intermediate in the catalytic allylation step in the acetylation process using **1**.¹ We describe herein a novel and efficient synthesis of the oxodimethylenemethane complexes **3** from the palladium and platinum complexes **2**. Complex **3** is known to be the oxygen analogue of the trimethylenemethane complexes,² which have a high synthetic potential.


The π -allylpalladium complex **2a** was readily prepared as shown in Scheme 1. Treatment of this palladium complex **2a** with KOH–propan-2-ol at room temperature resulted in unexpected bond cleavage of the acetal moiety to give the oxodimethylenemethane complex **3a** (44% yield). Variable temperature ¹H NMR (400 MHz, CDCl₃) of **3a** showed the coalescence of methylene proton resonances at a higher temperature, 29 °C, compared with –20 °C for **3b**,[†] showing that the interconversion of the metallacyclobutanone ring of **3a** is slower than that of **3b**. Thus, palladacyclobutanone **3a**

should be considered to have the higher contribution of the π -allylic canonical structure **3'** than the platinum analogue.³


It should be noted that this is the first case of the preparation of the unsubstituted parent (oxodimethylenemethane)palladium complexes.⁴ The isolated complex **3a** reacted with an equivalent amount of norbornadiene to afford 3-acetyltricyclo[3.2.1.0^{2,4}]oct-6-ene quantitatively, supporting the fact that **3a** is proposed to be a reaction intermediate in the Pd(0)-catalysed cyclization of 1-acetoxy-3-(trimethylsilyl)propan-2-one and norbornadiene.⁵

A similar reaction of **2b** with NaOH(aq.) led to the formation of the (oxodimethylenemethane)platinum complex **3b** in 93% yield.³

Complex **3a** in moistened organic solvent showed basicity *viz.* proton acceptability. The basicity of **3a**, pH = 9.7 (0.01 mmol in H₂O–acetone = 0.1 ml:0.9 ml), was stronger than that of **3b** (pH = 8.6) under the same conditions. This result may support the above-mentioned assumption concerning the higher contribution of the allylic canonical structure in **3a** than in **3b**. Protonation of **3** with trifluoroacetic acid resulted in the formation of the novel 2-hydroxy-substituted π -allyl complexes **4e** and **f**.



[†] Coalesced at –78 °C in [D₂H₈]THF (400 MHz), see ref. 3.



Scheme 1 Reagents and conditions: for **2a**: **1** (1 equiv.), Pd₂(dba)₃ (0.4 equiv.), PPh₃ (2.1 equiv.) CH₂Cl₂-acetone (*in situ*), 20 °C; for **2b**: Pt(PPh₃)₄ (0.8 equiv.), C₆H₆, 20 °C; for **2c,d**: **2a,b** (1 equiv.), NH₄PF₆ (1.1 equiv.), CH₂Cl₂-acetone, 20 °C; for **3a**: **2a** (1 equiv.), 0.1 mol dm⁻³ KOH-PrⁱOH (1.1 equiv.), CH₂Cl₂-acetone, 20 °C; for **3b**: **2b** (1 equiv.), 0.1 mol dm⁻³ NaOH (aq.) (1.1 equiv.), tetrahydrofuran (THF), 20 °C; for **4c,d**: **2c,d** (1 equiv.), 60% HPF₆ (aq.) (excess), CH₂Cl₂, 20 °C; for **4e,f**: **3a,b** (1 equiv.), CF₃CO₂H (excess), CD₂Cl₂, 20 °C; (H₄dba = dibenzylideneacetone)

Complexes **4c** and **d** were prepared by acidic hydrolysis of the π-allyl complexes **2c** and **d**, respectively. Complexes **3** and **4** are intertransferable because the regeneration of **3** was achieved by treatment of **4c** and **d** with pyridine.

In summary, complexes **2** showed high reactivity under both basic and acidic conditions to provide an efficient route to

unsubstituted oxodimethylenemethane and 2-hydroxy π-allyl complexes.‡

Received, 12th February 1993; Com. 3100878I

References

- I. Ikeda, X.-P. Gu, T. Okuhara and M. Okahara, *Synthesis*, 1990, 32; X.-P. Gu, N. Nisida, I. Ikeda and M. Okahara, *J. Org. Chem.*, 1987, **52**, 3192; X.-P. Gu, Y. Kirito, I. Ikeda and M. Okahara, *J. Org. Chem.*, 1990, **55**, 3391.
- P. Binger and U. Schuchardt, *Chem. Ber.*, 1981, **114**, 3313.
- J. Fawcett, W. Henderson, M. D. Jones, R. D. W. Kemmitt, D. R. Russell, B. Lam, S. K. Kang and T. A. Albright, *Organometallics*, 1989, **8**, 1991.
- R. D. W. Kemmitt, P. McKenna, D. R. Russell and L. J. S. Sherry, *J. Chem. Soc., Dalton Trans.*, 1985, 259.
- B. M. Trost and S. Schneider, *J. Am. Chem. Soc.*, 1989, **111**, 4430.

‡ Satisfactory elemental analysis (C, H, Cl) for **2c**, **3a**, **3b** and **4c** was obtained.

Melting point (°C) and ¹H NMR data [600 MHz, solvent CDCl₃ (unless stated otherwise), 25 °C] are as follows: **2b**: m.p. 228–232 (dec.); δ 3.03 (br, 2H, *syn*-CH₂C), 3.30 (s, 2H, *anti*-CH₂C), 3.42 (s, 3H, OCH₃), 4.94 (s, 2H, OCH₂O), 7.28 (m, 24H, *o,m*-Ph), 7.38 (t, 6H, *p*-Ph). **2c**: m.p. 200–204 (dec.); δ 3.22 (m, 2H, *syn*-CH₂C), 3.44 (s, 2H, *anti*-CH₂C), 3.38 (s, 3H, OCH₃), 4.88 (s, 2H, OCH₂O), 7.24 (m, 24H, *o,m*-Ph), 7.38 (t, 6H, *p*-Ph). **2d**: m.p. 196–201 (dec.); δ 2.75 (dd, 2H, ²J_{PH} 42, ³J_{PH} 9.1, ²J_{HH} 4.8 Hz, *syn*-CH₂C), 3.27 (d, 2H, ²J_{HH} 4.8 Hz, *anti*-CH₂C), 3.42 (s, 3H, OCH₃), 4.92 (s, 2H, OCH₂O), 7.28 (m, 24H, *o,m*-Ph), 7.38 (t, 6H, *p*-Ph). **3a**: m.p. 165–175 (dec.); δ 2.40 (br, 2H, *syn*-CH₂), 2.80 (br, 2H, *anti*-CH₂), 7.20 (m, 24H, *o,m*-Ph), 7.28 (t, 6H, *p*-Ph); δ (30 °C) 2.58 (br, 4H, CH₂). **3b**: m.p. 222–226 (dec.); δ 2.35 (d, 4H, ²J_{PH} 48, ³J_{PH} 3.0 Hz, CH₂), 7.18 (t, 12H, *m*-Ph), 7.27 (m, 18H, *o,p*-Ph). **4c**: m.p. 173–193 (dec.); δ (CD₂Cl₂) 2.68 (m, 2H, *syn*-CH₂), 3.40 (s, 2H, *anti*-CH₂), 7.20 (dd, 12H, *o*-Ph), 7.26 (t, 12H, *m*-Ph), 7.38 (t, 6H, *p*-Ph). **4d**: m.p. 180–185 (dec.); δ (CD₂Cl₂) 2.30 (dd, 2H, ²J_{PH} 46, ³J_{PH} 10, ²J_{HH} 4.3 Hz, *syn*-CH₂), 3.24 (d, 2H, ²J_{HH} 4.3 Hz, *anti*-CH₂), 7.24 (m, 24H, *o,m*-Ph), 7.36 (t, 6H, *p*-Ph). **4e**: δ (CD₂Cl₂) 2.62 (m, 2H, *syn*-CH₂), 3.45 (s, 2H, *anti*-CH₂), 7.20 (dd, 12H, *o*-Ph), 7.28 (t, 12H, *m*-Ph), 7.38 (t, 6H, *p*-Ph). **4f**: δ (CD₂Cl₂) 2.45 (dd, 2H, ²J_{PH} 46, ³J_{PH} 10, ²J_{HH} 4.3 Hz, *syn*-CH₂), 3.15 (d, 2H, ²J_{HH} 4.3 Hz, *anti*-CH₂), 7.22 (dd, 12H, *o*-Ph), 7.26 (t, 12H, *m*-Ph), 7.40 (t, 6H, *p*-Ph).