

A Novel Synthesis of 4-Dioxy-2-enones from 1,3-Dienes using Pyridinium Dichromate-*tert*-Butyl Hydroperoxide

Shridhar Bhat, Nallaperumal Chidambaram and Srinivasan Chandrasekaran*

Department of Organic Chemistry, Indian Institute of Science, Bangalore-560 012, India

















Treatment of 1,3-dienes with the pyridinium dichromate (PDC)-*Bu*^tOOH reagent system leads directly to the formation of 4-*tert*-butyldioxy-2-enones in good yield under mild reaction conditions.

The addition reaction of alkyl hydroperoxide and cyclopentadiene in the presence of Fe^{II} and Cu^{II} salts yields among other products, 4-alkoxycyclopent-2-ene.¹ Oxidative cleavage of endoperoxides of 1,3-dienes using acidic or basic reagents,^{2a} Fe^{II}, Ti^{III}, Co^{II}, Rh^I and Pd⁰ salts or complexes^{2b}

gives rise to the corresponding 4-hydroxyenones as one of the products; however, the reactions generally lead to low yields and difficult separations.

The PDC-*Bu*^tOOH system has been shown to be very effective in a variety of oxidative transformations.³ Interest-

Table 1 Reaction of 1,3-dienes with PDC–Bu^tOOH

Entry	Substrate	Product ^a	Yield (%) ^b
1			76
2			55
3			52
4			24 ^c
5			60
6			42 ^c
7			86
8			73

^a All the products gave satisfactory spectral and analytical data.
^b Yields refer to isolated products taking into account the recovery of starting materials (6–20%). ^c Formation of some unidentified polymeric material lowers the yield in these cases.

ingly, when 1,3-dienes were treated with a PDC and Bu^tOOH mixture supported on silica gel (substrate : PDC : Bu^tOOH = 1 : 2 : 3) for 8–13 h,[†] the only products that could be isolated in

[†] *General procedure:* Silica gel (1.505 g) and PDC (4 mmol, 1.505 g) were suspended in dichloromethane and cooled to 0 °C. This suspension was treated with Bu^tOOH (6 mmol, 0.8 ml, 70% solution in water) to give a crimson coloured mixture. After 10 min, a 1,3-diene (2 mmol) was added and the reaction was monitored by TLC. The reaction mixture was filtered using a sintered-glass funnel, through a pad of silica gel. The solvent was removed and the product was purified by column chromatography (silica gel, eluent 2% ethyl acetate in hexane).

a reasonable yield were the 4-*tert*-butyldioxy-2-enones.[‡] The results are shown in Table 1.

Both cyclic and acyclic dienes reacted with PDC–Bu^tOOH and this transformation is the first report of its type in the literature. Although substrates **13** and **15** initially gave the corresponding dioxyenones, only the diones could be isolated successfully. Surprisingly the diene moiety of the triene **11** did react enabling the isolation of compound **12**. In spite of the possibility of multiple reaction pathways open to the dienes **7** and **11** only **8** and **12** respectively, were obtained. Diene **9** with two tertiary centres at the diene termini gave **10** as expected.

A set of experiments was carried out with cyclopentadiene illustrating the use of Cr^{VI} reagent in this reaction. Replacing PDC with chromium trioxide and 4 equiv. of Bu^tOOH instead of 3 equiv., gave **2** in 74% yield. A catalytic amount of PDC (10 mol%) or chromium trioxide (10 mol%) and 6 equiv. of Bu^tOOH (at –5 °C, 24 h) in the reaction provided **2** in 65 and 70% yields respectively. Compounds **2**, **4**, **6** and **8** could be further converted to the corresponding 4-hydroxy-2-enones in quantitative yield following a closely related literature procedure for hydrogenolysis.⁴

The direct conversion of 1,3-dienes to the 4-dioxy-2-enones and their subsequent transformation to 4-hydroxy-2-enones could be useful in organic synthesis. *E.g.*, 4-hydroxycyclopent-2-enone which could be easily accessible from readily available and inexpensive cyclopentadiene, is a key intermediate in the synthesis of prostaglandins.⁵ The mechanism of this reaction is being studied in our laboratory.

The authors thank DST and UGC New Delhi for financial support.

Received, 11th January 1993; Com. 3/00142C

References

- H. Haubenstock, P. G. Mennitt and P. E. Butler, *J. Org. Chem.*, 1970, **35**, 3208; G. Stork and M. Isobe, *J. Am. Chem. Soc.*, 1975, **97**, 6260.
- (a) J. W. Cornforth, B. V. Milborrow and G. Ryback, *Nature*, 1975, **205**, 715; A. Murai, M. Ono, A. Akibo and T. Masamune, *J. Am. Chem. Soc.*, 1978, **100**, 7751; (b) M. Suzuki, Y. Oda and R. Noyori, *Tetrahedron Lett.*, 1981, **22**, 4413 and references cited therein.
- N. Chidambaram and S. Chandrasekaran, *J. Org. Chem.*, 1987, **52**, 5048; N. Chidambaram, K. Satyanarayana and S. Chandrasekaran, *Tetrahedron Lett.*, 1989, **30**, 2429; N. Chidambaram, S. Bhat and S. Chandrasekaran, *J. Org. Chem.*, 1992, **57**, 5013.
- S.-I. Murahashi, T. Naota and T. Kuwabara, *Synlett*, 1989, 62.
- M. Suzuki, A. Yanagisawa and R. Noyori, *J. Am. Chem. Soc.*, 1985, **107**, 3348.

[‡] 6–20% of the starting material was recovered. A small amount (5–10%) of a highly polar product formed in all these reactions has not been characterized.