

## Synthesis and Properties of a Novel Redox System containing Fullerene and *p*-Benzoquinone

Masahiko Iyoda,\* Fatema Sultana, Shigeru Sasaki and Masato Yoshida

Department of Chemistry, Faculty of Science, Tokyo Metropolitan University, Hachioji, Tokyo 192-03, Japan

Synthesis of a fullerene derivative of *p*-benzoquinone was carried out using the [4 + 2]cycloaddition of C<sub>60</sub> to 3,6-dihydroxybenzocyclobutene, followed by DDQ oxidation.

Buckminsterfullerene, C<sub>60</sub>, is known to possess considerably high electron affinity and low ionization potential on the basis of both experimental measurements and theoretical calculations.<sup>1</sup> Thus, C<sub>60</sub> can play the role as a  $\pi$ -acceptor and  $\pi$ -donor. Especially, the abilities of C<sub>60</sub> as the  $\pi$ -acceptor are very important properties,<sup>2</sup> and the chemical and electrochemical reductions of C<sub>60</sub> and its derivatives have been extensively studied.<sup>3,4</sup>

We are interested in the title compound **1**, because of a potential donor-acceptor interaction between fullerene and

*p*-benzoquinone parts in **1**, and because of the opposite acceptor-donor interaction between C<sub>60</sub> part and semiquinone radical or hydroquinone dianion in **2** and **3**. We report here the first synthesis of the fullerene derivative of quinones.

To construct the novel redox system composed of fullerene and 1,4-hydroquinone or *p*-benzoquinone, C<sub>60</sub> was allowed to react with 3,6-dimethoxybenzocyclobutene **4**<sup>5</sup> in *o*-dichlorobenzene at 220 °C for 24 h to give the mono-adduct **6**<sup>†</sup> in 44% yield, together with a mixture of isomers of the bis-adduct (34%) and the tris-adduct (11%). The removal of methyl groups in **6** was satisfactorily carried out using BBr<sub>3</sub>. Thus, the reaction of **6** with an excess of BBr<sub>3</sub> in benzene at 5 °C–room temp. for 24 h to give the hydroquinone derivative **7**<sup>†</sup> in 80% yield. Furthermore, the reaction of C<sub>60</sub> with 3,6-dihydroxybenzocyclobutene **5**<sup>6</sup> in *o*-dichlorobenzene at 230–235 °C for 24 h produced the mono-adduct **7** in one step in 32% yield. Oxidation of **7** with an excess of DDQ in benzene at room temp. for 15 min afforded the desired *p*-benzoquinone derivative **1**<sup>†</sup> in 98% yield. The compound **1** is stable and can be stored under air at room temp.

It is known that the adducts of C<sub>60</sub> to *o*-quinodimethane usually show the inversion of six-membered ring, and the activation energy for the ring inversion is estimated to be  $\Delta G^\ddagger = 13.2\text{--}14.6 \text{ kcal mol}^{-1}$  (1 cal = 4.184 J).<sup>7</sup> Thus, **6** and **7** show similar activation energies for the ring inversion (**6**:  $\Delta G^\ddagger = 13.7 \text{ kcal mol}^{-1}$  at 25 °C; **7**:  $\Delta G^\ddagger = 14.2 \text{ kcal mol}^{-1}$  at 40 °C). Interestingly, **1** shows the much lower activation energy ( $\Delta G^\ddagger = 11.0 \text{ kcal mol}^{-1}$  at –30 °C), although the quinone part of **1** may reduce the torsional and angular constraints. We now expect that the low barrier of inversion in **1** might be attributable to an electronic repulsion between the fullerene  $\pi$ -electron and the quinone part (acceptor-acceptor interaction).

In order to estimate the  $\pi$ - $\pi$  interaction between the fullerene and quinone parts in **1**, redox potentials of **1**, **6** and C<sub>60</sub> were measured by cyclic voltammetry. As shown in Table 1 and Fig. 1, **1** shows interesting redox properties. Thus, **6** shows typical three reduction waves which are cathodically shifted by 0.12 to 0.17 V as compared with those of C<sub>60</sub>.<sup>8</sup> In contrast, four reduction waves were observed in the case of **1** (Fig. 1). The first reduction is assigned to the reduction of *p*-benzoquinone part to the corresponding semiquinone radical **2**. The second reduction ( $E_{\text{red}}^1$ ) corresponds to the reduction of the fullerene part to the fullerene anion radical. The third reduction ( $E_{\text{red}}^2$ ) seems to be two electron reduction to form the tetraanion, *i.e.* C<sub>60</sub><sup>2-</sup>–(CH<sub>2</sub>)<sub>2</sub>–C<sub>6</sub>H<sub>2</sub>O<sub>2</sub><sup>2-</sup>. Therefore, some interactions between the fullerene and *p*-benzoquinone parts can be expected in the anionic species.

This work was partly supported by the Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Science, and Culture, Japan (05233224).

Received, 29th April 1994; Com. 4/02554G





Fig. 1 Cyclic voltammogram of **1** in benzonitrile: sample,  $2.5 \times 10^{-4} \text{ mol dm}^{-3}$ ; supporting electrolyte (Bu<sub>4</sub>NClO<sub>4</sub>)  $0.1 \text{ mol dm}^{-3}$ ; scan rate  $0.1 \text{ V s}^{-1}$ , vs. ferrocene/ferrocinium

Table 1 Redox potentials of C<sub>60</sub>, **6**, and **1**<sup>a</sup>

Compound	$E_{\text{red}}^{\text{quinone}}$	$E_{\text{red}}^1$	$E_{\text{red}}^2$	$E_{\text{red}}^3$
C <sub>60</sub>		–0.95	–1.35	–1.83
<b>6</b>		–1.07	–1.48	–2.00
<b>1</b>	–0.92	–1.10	–1.50	–2.07

<sup>a</sup> All potentials in V vs. ferrocene/ferrocinium; benzonitrile;  $0.1 \text{ mol dm}^{-3}$  Bu<sub>4</sub>NPF<sub>6</sub>;  $0.1 \text{ V s}^{-1}$ .

### Footnote

<sup>†</sup> Selected data for new fullerene derivatives. **6**: mp 350–360 °C (decomp.); FAB–MS *m/z* 885 (*M* + 1); <sup>1</sup>H NMR (CDCl<sub>3</sub>, –50 °C)  $\delta$  3.96 (s, 6H, OCH<sub>3</sub>), 4.33 (d, 2H, *J* 13.9 Hz, CH<sub>2</sub>), 4.93 (d, 2H, *J* 13.9 Hz, CH<sub>2</sub>) and 7.05 (s, 2H, =CH). **7**: mp 280–300 °C (decomp.);

FAB-MS  $m/z$  857 ( $M + 1$ );  $^1\text{H NMR}$  ( $[\text{2H}_8]$ toluene,  $-15^\circ\text{C}$ )  $\delta$  4.00 (d, 2H,  $J$  14.2 Hz,  $\text{CH}_2$ ), 4.86 (d, 2H,  $J$  14.2 Hz,  $\text{CH}_2$ ), 4.14 (s, 2H, OH) and 6.25 (s, 2H, =CH). **1**: mp 330–340 °C (decomp.); FAB-MS  $m/z$  856 ( $M + 2$ );  $^1\text{H NMR}$  ( $\text{CD}_2\text{Cl}_2$ ,  $-70^\circ\text{C}$ )  $\delta$  4.11 (d, 2H,  $J$  15.1 Hz,  $\text{CH}_2$ ), 4.80 (d, 2H,  $J$  15.1 Hz,  $\text{CH}_2$ ), 7.09 (s, 2H, =CH);  $^{13}\text{C NMR}$  ( $\text{CS}_2/[\text{2H}_6]$ acetone, room temp.)  $\delta$  38.5 ( $\text{CH}_2$ ), 65.8 ( $\text{C}_{60}$ ), 136.3 ( $\text{CH}=\text{CH}$ ), 137.3 ( $\text{C}=\text{C}$ ), 184.0 ( $\text{C}=\text{O}$ ), 141.0, 142.5, 142.9, 143.4, 143.9, 144.6, 145.4, 146.0, 146.3, 146.4, 146.6, 147.1, 147.3, 148.5 and 159.7 (15 signals of  $\text{C}_{60}$ ).

## References

- 1 W. Weltner, Jr. and R. V. Van Zee, *Chem. Rev.*, 1989, **89**, 1713.
- 2 P.-M. Allemand, K. C. Khemani, A. Koch, F. Wudl, K. Holczer, S. Donovan, G. Grüner and J. D. Thompson, *Science*, 1991, **253**, 301; P. W. Stephens, D. Cox, J. W. Lauher, L. Mihaly, J. B. Wiley, P.-M. Allemand, A. Hirsch, K. Holczer, Q. Li, J. D. Thompson and F. Wudl, *Nature*, 1992, **355**, 331.
- 3 P.-M. Allemand, A. Koch, F. Wudl, Y. Rubin, F. Diederich, M. M. Alvarez and R. L. Whetten, *J. Am. Chem. Soc.*, 1991, **113**, 1050; P.-M. Allemand, G. Srdanov, A. Koch, K. Khemani and F. Wudl, *J. Am. Chem. Soc.*, 1991, **113**, 2780; D. M. Cox, S. Behal, M. Disko, S. M. Gorun, M. Greaney, C. S. Hsu, E. B. Kollin, J. Millar, J. Robbins, W. Robbins, R. D. Sherwood and P. Tindall, *J. Am. Chem. Soc.*, 1991, **113**, 2940; D. Dubois, K. M. Kadish, S. Flanagan, R. E. Haufler, L. P. F. Chibante and L. J. Wilson, *J. Am. Chem. Soc.*, 1991, **113**, 4364; D. Dubois, K. M. Kadish, S. Flanagan and L. J. Wilson, *J. Am. Chem. Soc.*, 1991, **113**, 7773; Q. Xie, E. Pérez-Cordero and L. Echegoyen, *J. Am. Chem. Soc.*, 1992, **114**, 3978; W. K. Fullagar, I. R. Gentle, G. A. Heath and W. J. White, *J. Chem. Soc., Chem. Commun.*, 1993, 525.
- 4 Recently, a cyclic voltammetry study of some fullerene-ferrocene derivatives has been reported: M. Maggini, A. Karlsson, G. Scorrano, G. Sandonà G. Farnia and M. Prato, *J. Chem. Soc., Chem. Commun.*, 1994, 589.
- 5 T. Watabe, Y. Hosoda, K. Okada and M. Oda, *Bull. Chem. Soc. Jpn.*, 1987, **60**, 3801; M. Iyoda, S. Kitami, T. Yamauchi and M. Oda, *Chem. Lett.*, 1986, 2113.
- 6 M. Oda and Y. Kanao, *Chem. Lett.*, 1981, 37; M. Iyoda, M. Sakaitani, A. Kojima and M. Oda, *Tetrahedron Lett.*, 1985, **26**, 3719.
- 7 Y. Rubin, S. Khan, D. I. Freedberg and C. Yeretizian, *J. Am. Chem. Soc.*, 1993, **115**, 344; T. Tago, T. Minowa, Y. Okada and J. Nishimura, *Tetrahedron Lett.*, 1993, **34**, 8461.
- 8 T. Suzuki, Y. Maruyama, T. Akasaka, W. Ando, K. Kobayashi and S. Nagase, *J. Am. Chem. Soc.*, 1994, **116**, 1359.