

A New P-Chiral Aminophosphine Ligand Containing a 2,2'-Coupled Pyrrolidine–Phospholane Ring System. Synthesis and Coordination Properties with Rhodium(I) and Iridium(I) Fragments

Claudio Bianchini,^{a*} Stefano Cicchi,^a Maurizio Peruzzini,^a K. Michal Pietrusiewicz^b and Alberto Brandi^{c*}

^a Istituto per lo Studio della Stereochimica ed Energetica dei Composti di Coordinazione del CNR, Via J. Nardi 39, 50132 Firenze, Italy

^b Department of Organic Chemistry, Maria Curie-Skłodowska University, ul. Gliniana 33, 20-614 Lublin, Poland

^c Dipartimento di Chimica Organica 'U. Schiff', Università di Firenze, Via G. Capponi 9, I-50121 Firenze, Italy

A new approach to the synthesis of chiral P,N-ligands containing a stereogenic phosphorus atom is reported; the synthesis, characterization and coordination properties of (*PR*, 2*R*, 3*R*, 2'*R*, 3'*S*, 4'*S*) 2[2'(3',4'-di-*tert*-butoxy-1'-methyltetrahydropyrrolyl)]-3-methoxy-1-phenyltetrahydrophosphole (**P*-N-1**), which contains a stereogenic phosphorus in addition to five contiguous stereocentres in the ligand backbone; compound **P*-N-1** represents the first example of a new class of chiral P,N-ligands with the heteroatoms incorporated in two 2,2'-coupled pyrrolidine–phospholane rings.

The design and development of chiral aminophosphine ligands is being actively pursued for applications in transition metal asymmetric catalysis.¹ So far, however, very few P,N-ligands are known which contain a stereogenic phosphorus atom in addition to stereocentres in the ligand backbone.² This report details our endeavours to design, synthesize and characterize a new class of optically pure P,N-ligands in which the phosphorus donor is a stereocentre and the heteroatoms are incorporated in two 2,2'-connected five-membered rings.


A structural arrangement of this type is expected to favour the formation of metal complexes in which the five-membered chelate ring is forced into a specific conformation, as is often the case for some widely used chiral diphosphine ligands, such as those adopting binaphthyl-like chelate rings.

We also report some preliminary results on the bonding capabilities of **P*-N-1** with selected organorhodium(I) and organoiridium(I) fragments, which are widely used in a variety of homogeneous catalytic reactions.


The synthesis of **P*-N-1** was carried out as illustrated in Scheme 1. The isoxazolidine **4**,³ obtained as the major isomer of the 1,3-dipolar cycloaddition of the nitron **2**⁴ with the phosphole **3**,⁵ was treated with MeI in CH₂Cl₂ to give the isoxazolinium salt **5** in quantitative yield. The salt **5** was

reduced to the amino alcohol **6** with Zn–MeCO₂H,⁶ and after treatment with NaH in THF, was alkylated with MeI to afford the protected phosphine oxide **7**. The key step of the synthesis of **P*-N-1** is thus the stereoselective reduction of **7**. This reaction was carried out in refluxing benzene for 12 h with HSiCl₃ which, in the presence of NEt₃, is known to bring about the reduction of the P=O functionality with retention of configuration.^{2,7†} In this way, the optically pure aminophosphine ligand (*PR*, 2*R*, 3*R*, 2'*R*, 3'*S*, 4'*S*) 2[2'(3',4'-di-*tert*-butoxy-1'-methyltetrahydropyrrolyl)]-3-methoxy-1-phenyltetrahydrophosphole (**P*-N-1**)[‡] was obtained in 55% yield (based on **4**) as a single stereoisomer [³¹P{¹H}] NMR (CDCl₃) δ –15.22 (s)] with a known absolute configuration at all six stereogenic centres. This methodology is suitable for the synthesis of a class of P,N-ligands bearing different substituents and in either enantiomeric form.

The ligand **P*-N-1** reacts in deaerated CH₂Cl₂ with an equivalent amount of [Rh(CO)₂Cl]₂, [Rh(cod)Cl]₂ or [Ir(cod)Cl]₂ (cod = 1,5-cyclooctadiene) to give brown to orange microcrystals of the square-planar complexes (**P*-N-1**)Rh(CO)Cl **8**,[‡] (η¹-*P*-**P*-N-1**)Rh(cod)Cl **9**,[‡] and (η¹-*P*-**P*-N-1**)Ir(cod)Cl **10**,[‡] respectively (Scheme 2). The ³¹P{¹H} NMR spectra of **8** and **9** show the expected doublet multiplicity with coupling constants within the range for Rh^I square-planar complexes containing phosphorus donor ligands⁸ [**8**, δ 76.51 (d,



Scheme 1 Reagents and conditions: i, MeI; ii, MeCO₂H, Zn; iii, MeI, NaH; iv, HSiCl₃



Scheme 2

J_{PRh} 171.2 Hz); **9**, δ 25.82 (d, J_{PRh} 146.2 Hz); **10**, δ 19.26 (s)]. The coordination of the nitrogen donor, demonstrated by the low-field shift of the N–Me signal,⁹ occurs only for the carbonyl complex **8**, which thus contains a bidentate **P*–N-1** ligand. The carbonyl ligand in **8** lies *trans* to the nitrogen atom as inferred from the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum, which shows the CO resonance as a doublet of doublets at δ 189.12 (J_{CRh} 70.9, J_{CP} 19.0 Hz).¹⁰

Unlike **8**, the two diene complexes **9** and **10** do not exhibit a bonding interaction of the pyrrolidine N atom with the metal as shown by the chemical shifts of both the N–Me group⁹ (**9**, δ 2.51; **10**, δ 2.75) and the P nucleus (**9**, δ 29.49; **10**, δ 19.26). Thus, a chloride, a bidentate cod ligand and a phosphorus atom complete the square-planar geometry about the metal centre in both **9** and **10**. The coordination of the nitrogen donor atoms can readily be achieved by treatment of **9** and **10** in THF with a chloride scavenger such as NaBPh₄ in ethanol. As a result, the complexes [(**P*–N-1**)Rh(cod)]BPh₄ **11**‡ and [(**P*–N-1**)Ir(cod)]BPh₄ **12**‡ are quantitatively obtained, in which the presence of a chelating P,N-ligand ring is shown by the highfield shift of the phosphorus resonance (**11**, δ 58.50; **12**, δ 54.14).⁸

The potential of **P*–N-1** and related ligands bearing different substituents on the pyrrolidine and phospholane rings in asymmetric catalysis is currently under investigation in our laboratories.

C. B. thanks Progetto Strategico 'Tecnologie Chimiche Innovative', CNR, Rome (Italy), and the EC (contract CI1*-CT93-0329), A. B. thanks M.U.R.S.T. (Ministero della Ricerca Scientifica e Tecnologica) for financial support.

Received, 21st December 1994; Com. 4/07801B

Footnotes

‡ The retention of configuration at the phosphorus atom has been confirmed by reoxidation of the **P*–N-1** ligand to the starting product **7** by reaction

with hydrogen peroxide, which is known to proceed with complete retention of configuration (L. Horner, *Pure Appl. Chem.*, 1964, **9**, 225).

‡ Satisfactory spectroscopic and analytical data were obtained for all compounds.

References

- H. Brunner and W. Zettlemeier, *Handbook of Enantioselective Catalysis with Transition Metal Compounds*, VCH, Weinheim, 1993, vol. II; M. J. Burk, J. E. Feaster and R. L. Harlow, *Tetrahedron Asymmetry*, 1991, **2**, 569; K. Mashima, T. Agutawa, X. Zhang, T. Taketomi, H. Kumobayashi and S. Agutawa, *J. Am. Chem. Soc.*, 1993, **115**, 3318; A. Albinati, F. Lianza, H. Berger, P. S. Pregosin, H. Ruegger and R. W. Kunz, *Inorg. Chem.*, 1993, **32**, 478; C. Bianchini, E. Farnetti, L. Glendenning, M. Graziani, G. Nardin, M. Peruzzini, E. Rocchini and F. Zanobini, *Organometallics*, in the press; C. G. Arena, F. Nicolò, D. Drommi, G. Bruno and F. Faraone, *J. Chem. Soc., Chem. Commun.*, 1994, 2251; J. M. Brown, D. I. Hulmes and T. P. Layzell, *J. Chem. Soc., Chem. Commun.*, 1993, 1329.
- K. M. Pietrusiewicz and M. Zablocka, *Chem. Rev.*, 1994, **94**, 1375.
- A. Brandi, S. Cicchi, A. Goti, M. Koprowski and K. M. Pietrusiewicz, *J. Org. Chem.*, 1994, **59**, 1315; K. M. Pietrusiewicz, W. Wisniewski, S. Cicchi and A. Brandi, *Phosphorus Sulfur Silicon Relat. Elem.*, in the press.
- S. Cicchi, I. Hold and A. Brandi, *J. Org. Chem.*, 1993, **58**, 5274; R. Ballini, E. Marcantoni and M. Petrini, *J. Org. Chem.*, 1992, **57**, 1316.
- L. D. Quinn, J. P. Gratz and P. T. Barket, *J. Org. Chem.*, 1968, **33**, 1034.
- J. J. Tufariello and J. M. Puglis, *Tetrahedron Lett.*, 1986, **27**, 1265.
- S. E. Cremer and R. J. Chorvat, *J. Org. Chem.*, 1967, **32**, 4066.
- P. S. Pregosin and R. W. Kuntz, *^{31}P and ^{13}C NMR of Transition Metal Complexes*, ed. P. Diehl, E. Fluck and R. Kosfeld, Springer Verlag, Berlin, 1979.
- C. Bianchini, P. Innocenti, D. Masi, M. Peruzzini and F. Zanobini, *Gazz. Chim. Ital.*, 1992, **122**, 461.
- J. C. M. Wehman-Oovevaar, G. M. Kaptejin, D. M. Grove, W. J. J. Smeets, A. L. Spek and G. van Koten, *J. Chem. Soc., Dalton Trans.*, 1994, 703.