

A novel one-pot route to cyclopenta[*b*]indole via the palladium-catalysed carbonylative cross-coupling reaction of indolylborate with prop-2-ynyl carbonate

Minoru Ishikura,* Yukinori Matsuzaki and Isao Agata*

Faculty of Pharmaceutical Sciences, Health Sciences University of Hokkaido, Ishikari-Tobetsu, Hokkaido 061-02, Japan

A novel one-pot protocol for the formation of cyclopenta[*b*]indole is developed via the palladium catalysed carbonylative cross-coupling reaction.

The marked facility with which palladium complexes undergo an oxidative addition to prop-2-ynyl halides and esters in a S_N2' manner to form allenylpalladium complexes has been documented, as has the susceptibility of the resulting palladium complexes to a wide variety of chemical transformations.¹ As part of our ongoing studies, we have previously demonstrated

the use of the allenylpalladium complexes for the construction of 2-allenylindoles via the palladium-catalysed cross-coupling reaction of indolylborate **1** with prop-2-ynyl carbonates.² Despite the known reactivity of the allenylpalladium complexes under carbonylation conditions in the presence of a nucleophile producing allenic esters, amides, and ketones,³ its use for the formation of allenyl ketones via the carbonylative cross-coupling reaction is unknown. This posed the interesting question of whether the palladium-catalysed reaction of borate **1** with prop-2-ynyl carbonate can be effective under carbonylation conditions.


The feasibility of the palladium-catalysed carbonylation was first tested with borate **1** and carbonate **2** (Table 1); thus, cyclopenta[*b*]indole **4**,⁴ a basic framework of indole alkaloids such as yuehchukene, pasparine and paxilline, could be isolated in a one-pot manner (Scheme 1).

In general, the present carbonylation reaction can be effected by simply stirring a mixture of borate **1** (generated *in situ* from


Table 1 Cyclopenta[*b*]indoles **4** from borate **1** and carbonates **2**

Carbonate	PdL _n	Yield of 4 ^a (%)
2a	PdCl ₂ (PPh ₃) ₂	58
2a	Pd(OAc) ₂ + 2PPh ₃	60
2a	Pd(PPh ₃) ₄	63
2b	PdCl ₂ (PPh ₃) ₂	60
2c	PdCl ₂ (PPh ₃) ₂	61
2d	PdCl ₂ (PPh ₃) ₂	44 (14) ^b
2e	PdCl ₂ (PPh ₃) ₂	— ^c
2f	Pd(OAc) ₂ + 2PPh ₃	— (56) ^d
2f	Pd(PPh ₃) ₄	— (34) ^d
2f	PdCl ₂ (PPh ₃) ₂	— (38) ^d

^a All yields based on 1-methylindole. ^b Yield of allene **8b** (%). ^c No cross-coupling products. ^d Yield of allene **8a** (%).



Scheme 1



1-methylindole and Bu^tLi, followed by treatment with triethylborane in THF) and 1.5 equiv. of prop-2-ynyl carbonate (**2**, **3**, **6**) in the presence of palladium complex (5 mol%) under 10 atm of carbon monoxide pressure at 60 °C for 12 h.

The carbonylative cross-coupling reaction proceeded to produce cyclopenta[*b*]indole **4** when acetylenic carbon of **2** bears a substituent (such as phenyl and alkyl); otherwise, any cross-coupling product could not be seen in attempted reactions with **2e** bearing the sterically demanding trimethylsilyl group. Allenylindole **8a** was formed solely on the reaction of **1** with terminal alkyne **2f**. Furthermore, carbonates **3** could also be subjected to the carbonylation reaction, providing cyclopenta[*b*]indoles (**5a** in 62%; **5b** in 44%). This one-pot process should involve the intermediary formation of allenyl ketone **A**, and subsequent nucleophilic addition of the C-3 carbon of indole to the internal allenic carbon of **A** (Scheme 1). On treating **1** with **6a** and **6b**, geometrical mixtures of *E*-**7a** and *Z*-**7a** (in 38%) and *E*-**7b** and *Z*-**7b** (in 59% yield) were obtained, respectively, in the ratio 1 : 1, where the axially dissymmetric features of the allene unit of **A** allowed the generation of the geometrical isomers.

In summary, the feasibility of the palladium-catalysed carbonylative cross-coupling reaction of borate **1** with prop-2-ynyl carbonates, accompanied by subsequent cyclisation, is

apparent, which provides a new one-pot procedure for the generation of cyclopenta[*b*]indoles.

References

- 1 T. Jeffery-Luong and G. Linstumelle, *Tetrahedron Lett.*, 1980, **21**, 5019; K. Ruitenbergh, H. Kleijn, C. J. Elsevier, J. Meijer and P. Vermeer, *Tetrahedron Lett.*, 1981, **22**, 1451; H. Kleijn, J. Meijer, G. C. Overbeek and P. Vermeer, *Recl. Trav. Chim. Pays-Bas*, 1982, **101**, 97; C. J. Elsevier, H. Kleijn, K. Ruitenbergh and P. Vermeer, *J. Chem. Soc., Chem. Commun.*, 1983, 1529; C. J. Elsevier, P. M. Stehouwer, H. Westmijze and P. Vermeer, *J. Org. Chem.*, 1983, **48**, 1103; E. Keinan and E. Bosch, *J. Org. Chem.*, 1986, **51**, 4006; J. Tsuji and T. Mandai, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 2589.
- 2 M. Ishikura and I. Agata, *Heterocycles*, 1996, **43**, 1591.
- 3 N. D. Trieu, C. J. Elsevier and K. Vrieze, *J. Organomet. Chem.*, 1987, **325**, C23; T. Mandai, K. Ryoden, M. Kawada and J. Tsuji, *Tetrahedron Lett.*, 1991, **32**, 7683; T. Mandai, S. Suzuki, A. Alkawa, T. Murakami, M. Kawada and J. Tsuji, *Tetrahedron Lett.*, 1991, **32**, 7687; T. Mandai, H. Kunitomi, K. Higashi, M. Kawada and J. Tsuji, *Synlett.*, 1991, 697; J. A. Marshall and M. A. Wolf, *J. Org. Chem.*, 1996, **61**, 3238.
- 4 C.-A. Harrison, R. Leineweber, C. J. Moody and J. M. J. Williams, *J. Chem. Soc., Perkin Trans. 1*, 1995, 1127; M. Ishikura, *Heterocycles*, 1995, **41**, 1385.

Received, 5th August 1996; Com. 6/05438B