

Efficient coupling reactions of allyl amines with soft nucleophiles using nickel-based catalysts


Hervé Bricout, Jean-François Carpentier and André Mortreux*

Laboratoire de Catalyse Hétérogène et Homogène, Groupe de Chimie Organique Appliquée, URA CNRS 402, ENSCL, BP 108-59652 Villeneuve d'Ascq Cedex, France

Substitution reactions of *N,N*-diethylallylamine **1** with soft nucleophiles such as active methylene compounds **2a–c** and piperidine **5** proceed much more rapidly in the presence of Ni(dppb)₂ [dppb = 1,4-bis(diphenylphosphino)butane] as catalyst than with comparable palladium systems.

Recently we reported the effectiveness of preformed Ni⁰-bisphosphine catalysts for the rearrangement of dialkyl allylmalonate derivatives,^{1a} a recently discovered case of nucleophilic allylic substitution which proceeds *via* C–C bond cleavage.^{1b} In this special reaction, nickel catalysts proved to be more active than comparable palladium systems, suggesting the superior ability of the former to activate allylic substrates having a poor leaving group. Here we report that such nickel-based systems are much more efficient than palladium catalysts for coupling of allyl amines² with so-called 'soft' nucleophiles.³

The comparative study was performed using the simple *N,N*-diethylallylamine **1** and active methylene compounds **2a–c** as shown in eqn. (1). The results are summarized in Table 1. In all



cases, the reaction was totally selective for allylation products, although a mixture of monoallylation (**3a–c**) and diallylation (**4a–c**) products was obtained. Using 2 mol% of Ni(dppb)₂ generated from Ni(COD)₂ and 2 mol equiv. of the diphosphine gave allylation products of dimethyl malonate **2a** in virtually quantitative yield with an initial activity of 50 h⁻¹ (turnover frequency, TOF) at 80 °C in THF (entry 1). Additional base such as BSA [BSA = *N,O*-bis(trimethylsilyl)acetamide] did not affect the reaction (entry 2). On the other hand, a significant increase in the catalytic activity arose upon adding tetrabutylammonium perchlorate as a promoter (5 equiv. with respect to Ni) (entry 3), in direct line with our previous observations on nickel-catalysed coupling reactions of allylic alcohol derivatives.⁴ Changing the solvent to DMF further increased the catalytic activity, so that the completion time decreased to 20 min at 80 °C (entry 4), allowing the reaction to be performed at 50 °C (entry 5). At this temperature, the best catalytic activity was observed by combining the solvent effect of DMF and the promoting effect of [NBu₄][ClO₄] (entry 6). Surprisingly, in light of palladium literature,^{2b–h} the *in situ* formation of the allyl ammonium salt *via* addition of acetic acid proved somewhat detrimental (entry 7). The use of palladium catalysts led to poorer results. As a matter of fact, no reaction at all occurred using the *in situ* combination Pd(OAc)₂-PPh₃ (1 : 5) in THF with BSA as a base (entry 8). Modest activity was obtained with this catalytic system in DMF, which was slightly improved upon adding acetic acid (entries 9 and 10). The *in situ* combination Pd(OAc)₂-dppb (1 : 3) showed the best performance of the palladium catalysts tested (entry 11), but the latter is still at least ten times less active than the best nickel-based

Table 1 Catalysed coupling reactions of *N,N*-diethylallylamine **1** with methylene active compounds **2a–c**

Entry	2	Catalytic system ^a	Solvent	Additive (equiv.)	T/°C	Conv. (%)	t/h ^b	3:4	TOF ^c /h ⁻¹
1	2a	Ni ⁰ /dppb	THF	—	80	100	12	87:13	35
2	2a	Ni ⁰ /dppb	THF	BSA (75)	80	100	12	88:12	10
3	2a	Ni ⁰ /dppb	THF	[NBu ₄][ClO ₄] (5)	80	100	2	90:10	200
4	2a	Ni ⁰ /dppb	DMF	—	80	100	0.3	87:13	≥ 170
5	2a	Ni ⁰ /dppb	DMF	—	50	95	3.5	87:13	85
6	2a	Ni ⁰ /dppb	DMF	[NBu ₄][ClO ₄] (5)	50	95	2.8	88:12	100
7	2a	Ni ⁰ /dppb	DMF	AcOH (10)	80	100	3	88:12	40
8	2a	Pd ^{II} /PPh ₃	THF	BSA (75)	80	0	15	—	0
9	2a	Pd ^{II} /PPh ₃	DMF	—	80	54	17	90:10	6
10	2a	Pd ^{II} /PPh ₃	DMF	AcOH (10)	80	64	15	94:6	8
11	2a	Pd ^{II} /dppb	DMF	—	80	100	5	90:10	45
12	2b	Ni ⁰ /dppb	DMF	—	80	100	< 0.1	73:27	≥ 600
13	2b	Ni ⁰ /dppb	DMF	—	50	100	0.5	72:28	350
14	2b	Pd ^{II} /PPh ₃	DMF	—	80	93	9	77:23	25
15	2b	Pd ^{II} /dppb	DMF	—	80	100	1	75:25	215
16	2b	Pd ⁰ /dppb	DMF	—	80	95	1.5	81:19	110
17	2b	Pd ⁰ /dppb	THF	—	80	80	15	76:24	5
18	2c	Ni ⁰ /dppb	DMF	—	80	10	0.25 ^d	97:3	20
19	2c	Pd ^{II} /dppb	DMF	—	80	100	1	78:22	150

^a [Metal]/[**1**]/[**2**] = 1:50:75; Ni⁰/dppb: prepared from Ni(COD)₂ and 2 equiv. of the phosphine ligand; Pd^{II}/PPh₃: prepared *in situ* from Pd(OAc)₂ and 5 equiv. of PPh₃; Pd^{II}/dppb: prepared *in situ* from Pd(OAc)₂ and 3 equiv. of dppb. ^b Optimized time for conversion of **1** into **3** and **4** as indicated by quantitative GLC; No side products were observed. ^c Initial catalytic activity (< 10% conversion). ^d 10% yield after 15 min as after 16 h.

Table 2 Catalysed exchange reactions of *N,N*-diethylallylamine **1** with piperidine **5**^a

Entry	Catalytic system	Solvent	Additive (equiv.)	Conv. (%) ^b	t/min	TOF/h ⁻¹
1	Ni ⁰ /dppb	DMF	—	100	360	35
2	Ni ⁰ /dppb	DMF	[NBu ₄][ClO ₄] (5)	100	150	60
3	Ni ⁰ /dppb	DMF	AcOH (5)	100	40	400
4	Ni ⁰ /dppb	THF	AcOH (5)	45	1020	4
5	Ni ⁰ /dppb	DMF	AcOH (50)	100	5	≥ 600
6	Pd ^{II} /dppb	DMF	AcOH (50)	92	120	60

^a [Metal]/[**1**]/[**5**] = 1:50:500; *T* = 50 °C. ^b Conversion of **1** into *N*-allylpiperidine **6** as indicated by quantitative GLC.





Fig. 1 Average turnover frequencies of nickel and palladium catalysts for the coupling of **1** with **2a–c** (see Table 1 for reaction conditions)

system for this reaction. A similar trend, with even more marked differences between nickel and palladium catalysts, was observed for the alkylation of **1** with methyl acetoacetate **2b**. The simple system using Ni(dppb)₂ as the catalyst and DMF as the solvent allowed the reaction carried out with 2 mol% to Ni to be completed within 5 min at 80 °C and with an average activity of 100 h⁻¹ at 50 °C (entries 12 and 13).[†] The *in situ* combination Pd(OAc)₂-dppb proved once again to be the most effective palladium catalyst (entries 14–17), but its performance at 80 °C was still lower than that of the above nickel system at 50 °C. However, nickel catalysts proved inefficient for the coupling of **1** with acetylacetone **2c** (entry 18), because of a rapid catalyst poisoning evidenced by the gradual shift in the colour of the solution from orange–yellow [typical colour of Ni(dppb)₂] to pale green [Ni^{II} species, possibly Ni(acac)₂]. Although it must be pointed out that **2c** is the most acidic compound among active methylene compounds **2a–c**, there is no definitive evidence that this catalyst decay occurs *via* an oxidative protolysis of nickel intermediates.[‡] Anyway, the catalyst poisoning was still observed in the presence of an excess of **1** ([Ni]:[**1**]:[**2c**] = 1:75:50). No comparable phenomenon was observed with palladium catalysts, such as the *in situ* combination Pd(OAc)₂-dppb, which afforded alkylation products **3c/4c** with performance similar to that observed for the coupling of **1** with **2b** (entry 19). Fig. 1 summarises the overall performance of best Ni–dppb and Pd–dppb systems.

The results of the exchange reaction between **1** and piperidine **5** to give *N*-allylpiperidine **6** (eqn. 2) bring further evidence of



the general ability of nickel catalysts to promote nucleophilic substitution of allylamines (Table 2). We found that the aforementioned trends for the nickel-catalysed coupling of **1** with active methylene compounds **2a, b** proved also to be true for this exchange reaction. A notable exception arose from the

beneficial effect of added acetic acid as expected from the literature.^{2b–h} Namely, the formation of **6** was best conducted by using Ni(dppb)₂ as the catalyst, DMF as the solvent and by adding 1 equiv. of AcOH with respect to **1** (Table 2, entry 5). The most efficient palladium catalyst found [*in situ* combination Pd(OAc)₂-dppb, 1:3] took at least 20 times longer for the exchange reaction to go to completion than the latter nickel-based system under the same conditions (entry 6).


In conclusion we have demonstrated that Ni(dppb)₂ easily generated from Ni(COD)₂ and 2 equiv. of the bisphosphine is generally a much more active catalyst than comparable palladium systems for substitution of allylamines with ‘soft’ nucleophiles. Two explanations can be proposed for this trend: (i) the higher ability of nickel catalysts to activate allylic substrates having a poor leaving group, *i.e.* here the C–N bond, and/or (ii) the lower propensity of Ni to be coordinated by amines, thus resulting in larger amounts of active Ni–phosphine species. The limitation of these nickel-based systems lies in the sensitivity of nickel intermediates towards acetylacetone.

Footnotes

* E-mail: mortreux@ensc-lille.fr

[†] In a typical experiment (Table 1, entry 13), to Ni(COD)₂ (36 mg, 0.13 mmol) in a 50 ml glass reactor equipped with a Teflon cap was added a degassed solution of dppb (111 mg, 0.26 mmol) in DMF (12.5 ml) under nitrogen. After 15 min of magnetic stirring, **1** (0.74 g, 6.5 mmol), **2b** (1.13 g, 9.75 mmol) and heptane (1.00 g, 10 mmol, GLC internal standard) were added. The solution was stirred at 50 °C and the reaction was monitored by quantitative GLC analysis of aliquot samples. Coupling reactions involving a co-reagent and exchange reactions of **1** with piperidine were carried out in a similar manner. *In situ* palladium catalysts were prepared by mixing Pd(OAc)₂ with a degassed solution (THF or DMF) of the phosphine ligand. The resulting yellow solution was stirred for 15 min at room temperature before use.

[‡] Such a degradation of nickel intermediates could be rationalised as follows:



References

- (a) H. Bricout, J.-F. Carpentier and A. Mortreux, *Tetrahedron Lett.*, 1997, **38**, 1053; (b) Y. I. M. Nilsson, P. G. Andersson and J.-E. Bäckvall, *J. Am. Chem. Soc.*, 1993, **115**, 6609.
- Palladium-catalysed nucleophilic substitution of allylamines and related derivatives: (a) K. E. Atkins, W. E. Walker and R. M. Manyik, *Tetrahedron Lett.*, 1970, **43**, 3821; (b) B. M. Trost and E. Keinan, *J. Org. Chem.*, 1980, **45**, 2741; (c) A. Hosomi, K. Hoashi, S. Kohra, Y. Tominaga, K. Otaka and H. Sakurai, *J. Chem. Soc., Chem. Commun.*, 1987, 570; (d) F. Garro-Helion, A. Merzouk and F. Guibé, *J. Org. Chem.*, 1993, **58**, 6109; (e) S. Murahashi, Y. Imada and K. Nishimura, *Tetrahedron*, 1994, **50**, 453; (f) M. Grellier, M. Pfeffer and G. Van Koten, *Tetrahedron Lett.*, 1994, **35**, 2877; (g) S. Lemaire-Audoire, M. Savignac, J.-P. Genêt and J.-M. Bernard, *Tetrahedron Lett.*, 1995, **36**, 1267; (h) S. Lemaire-Audoire, M. Savignac, C. Dupuis and J.-P. Genêt, *Bull. Soc. Chim. Fr.*, 1995, **132**, 1157.
- To the best of our knowledge, the cross-coupling of allylamines with boronic acids (‘hard’ nucleophiles) is the sole example of the use of nickel catalysts: B. M. Trost and M. D. Spagnol, *J. Chem. Soc., Perkin Trans. 1*, 1995, 2083.
- H. Bricout, J.-F. Carpentier and A. Mortreux, *J. Chem. Soc., Chem. Commun.*, 1995, 1863.

Received in Cambridge, UK, 30th April 1997; 7/02954C