

Phosphorus monoxide as a terminal ligand

Marc J. A. Johnson, Aaron L. Odom and Christopher C. Cummins*†

Department of Chemistry, 2-227, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139-4307, USA

Dimethyldioxirane oxidation of phosphide $\text{PMo}[\text{N}(\text{R})\text{Ar}]_3$ [$\text{R} = \text{C}(\text{CD}_3)_2\text{Me}$, $\text{Ar} = \text{C}_6\text{H}_3\text{Me}_2\text{-3,5}$] efficiently provides $(\text{OP})\text{Mo}[\text{N}(\text{R})\text{Ar}]_3$, a complex shown by X-ray crystallography to contain a terminal phosphorus monoxide ligand, and which reacts with dimethylzirconocene to give addition of a Zr-Me bond across the phosphoryl moiety.

Known phosphorus monoxide complexes^{1–3} are those in which $\text{PO}^{4–10}$ serves as a triply bridging ligand. These have been prepared by oxo transfer to a $\text{P}_2\text{Ni}_2\text{W}$ cluster,¹ and by hydrolysis of a $\mu_3\text{-P}(\text{NPr}^i_2)$ functional group.³ Recent reports of terminal phosphide (P^{3-}) complexes^{11–15} hint at the possibility of preparing terminal PO complexes by oxygen-atom transfer to the terminal P ligand. Accordingly, dimethyldioxirane oxidation^{16,17} has been found to effect the desired conversion of $\text{PMo}[\text{N}(\text{R})\text{Ar}]_3$ **1** to $(\text{OP})\text{Mo}[\text{N}(\text{R})\text{Ar}]_3$ **2**. Synthetic, structural, and initial reactivity data pertaining to **2** are detailed herein.

Treatment of a -78°C gold-colored solution of **1** (1.0009 g, 1.4853 mmol) in dichloromethane (50 ml) with dimethyldioxirane (20 ml of a 0.075 M solution in acetone¹⁸) elicited a rapid color change to purple, the color of **2**.[‡] Workup consisted of cold-filtration, precipitation of **2** by addition of acetonitrile (75 ml, pre-chilled to just above its melting point), and collection of solid **2** (73.8% yield) by filtration. Once thus obtained in pure form, solid **2** can be manipulated and maintained at 25°C for at least *ca.* 5 days without noticeable decomposition. When dissolved in various solvents (*e.g.* benzene, diethyl ether, heptane and dichloromethane) **2** is more sensitive, undergoing gradual reversion to **1** according to ^1H and ^{31}P NMR spectroscopic monitoring.

Diamagnetic **2** exhibits a single ^{31}P NMR signal at δ 269.8, shifted dramatically upfield relative to **1** but in a range typical of

two-coordinate phosphorus compounds.¹⁹ Proton NMR data for **2** are indicative of a single environment for the $\text{N}(\text{R})\text{Ar}$ ligands, consistent with the threefold (propellor) symmetry typical of complexes containing three $\text{N}(\text{R})\text{Ar}$ ligands.^{20–25}

Single crystals of **2** suitable for an X-ray diffraction study[§] were obtained by slow cooling to -35°C of a diisopropyl ether solution. Molecular **2** evinces pseudo-threefold symmetry about the near-linear [$177.6(8)^\circ$] Mo-P-O moiety, which is flanked by the three *tert*-butyl substituents (Fig. 1). An intriguing feature of the structure is the molybdenum–phosphorus distance $d(\text{MoP}) = 2.079(5) \text{ \AA}$, which is shorter by 0.04 \AA than $d(\text{MoP})$ for terminal phosphide **1**.¹¹ Contraction of the molybdenum–phosphorus distance in this manner is explicable in terms of rehybridization at P upon addition of a terminal substituent.²⁶ The phosphorus–oxygen distance [$1.49(2) \text{ \AA}$] in **2** is comparable to that measured by IR diode laser spectroscopy for gaseous PO [$1.476370(15) \text{ \AA}$],^{27,28} and to $d(\text{PO})$ values for $\mu_3\text{-PO}$ complexes: $1.509(3)$,³ $1.462(9)$ and $1.480(10)$.¹ Phosphoryl halide species such as OPCl_3 contrastingly exhibit longer phosphorus–oxygen distances, on the order of $1.54\text{--}1.58 \text{ \AA}$,²⁹ while $d(\text{PO})$ for $\text{OP}(\text{C}_6\text{H}_{11})_3$ is $1.490(2) \text{ \AA}$.³⁰ Other metrical parameters pertaining to the structure of **2** are unexceptional.

Previously reported $(\text{SP})\text{Mo}[\text{N}(\text{R})\text{Ar}]_3$ **3**¹¹ was shown recently by X-ray crystallography to resemble **2** in terms of its monomeric nature and gross structural characteristics.³¹ The rough similarity of **2** and **3** was used to advantage to obtain assignment of the ν_{PO} (1232 cm^{-1}) and ν_{PS} (782 cm^{-1}) IR stretching frequencies for the two complexes, by digital subtraction of solution FTIR spectra. The ν_{PO} value for **2** is similar to that reported for gaseous PO (1220.25 cm^{-1}), and also is comparable to values reported for $\mu_3\text{-PO}$ complexes ($1260, 1169 \text{ cm}^{-1}$).

Initial reactivity studies indicate that the phosphoryl phosphorus in **2** is susceptible to nucleophilic attack in a manner reminiscent of the Fischer carbene synthesis.³² Accordingly, treatment of **2** (282.0 mg, 0.4088 mmol) with dimethylzirconocene³³ (103.3 mg, 0.4108 mmol) in benzene (7.5 ml) led to formation of $[(\text{R}^1)(\text{R}^2\text{O})\text{P}]\text{Mo}[\text{N}(\text{R})\text{Ar}]_3$ **4**, $\text{R}^1 = \text{Me}$, $\text{R}^2 = \text{Zr}(\text{Me})(\eta\text{-C}_5\text{H}_5)_2$, which was isolated by crystallization





Fig. 1 ORTEP drawing of the molecular structure of **2** with 35% probability ellipsoids. Selected bond lengths (Å) and angles ($^\circ$): Mo–P 2.079(5), Mo–N(1) 1.95(2), Mo–N(2) 1.94(2), Mo–N(3) 1.994(13), P–O 1.49(2); Mo–P–O 177.6(8), P–Mo–N(1) 101.7(4), P–Mo–N(2) 101.4(5), P–Mo–N(3) 101.6(5), N(1)–Mo–N(2) 113.1(6), N(1)–Mo–N(3) 117.2(6), N(2)–Mo–N(3) 118.0(6).



Scheme 1




Fig. 2 ORTEP drawing of the molecular structure of **4** with 35% probability ellipsoids. Selected bond lengths (Å) and angles (°): Mo–P 2.169(2), Mo–N(1) 1.995(5), Mo–N(2) 2.011(5), Mo–N(3) 1.993(5), P–O 1.613(5), P–C(1) 1.832(8), Zr–O 2.000(4), Zr–C(2) 2.253(11); Mo–P–O 127.7(2), Mo–P–C(1) 133.9(3), O–P–C(1) 97.6(3), Zr–O–P 163.6(3), O–Zr–C(2) 97.1(4), P–Mo–N(1) 101.6(2), P–Mo–N(2) 103.2(2), P–Mo–N(3) 101.7(2), N(1)–Mo–N(2) 109.1(2), N(1)–Mo–N(3) 112.6(2), N(2)–Mo–N(3) 125.0(2).

from a 1:1 diethyl ether–heptane mixture as orange–brown plates in 75.0% yield (Scheme 1). Diamagnetic **4** exhibits a single ^{31}P NMR signal at δ 284.4 [cf. δ 312 for $\text{Mo}[\text{P}(\text{C}_6\text{H}_{11})_2]_4$].³⁴ A reasonable oxidation state assignment for the Mo centre in **4** is +4. Proton NMR data for **4** are indicative of a single N(R)Ar ligand environment on the NMR timescale, although the solid-state structure of **4** reveals a low-symmetry conformation. A doublet ($^2J_{\text{PH}}$ 14 Hz) at δ 1.535 is attributed to the PMe moiety.

The three-coordinate phosphorus in **4** was shown by X-ray diffraction§ to be essentially planar [sum of angles = 359.2(3)°], consistent with Mo–P π bonding in the complex (Fig. 2). The observed $d(\text{MoP})$ of 2.169(2) Å is also consistent with a molybdenum–phosphorus multiple bond [cf. $d(\text{MoP})$ of 2.265(2) Å for $\text{Mo}[\text{P}(\text{C}_6\text{H}_{11})_2]_4$].³⁴ The conformation of **4** is noteworthy in that one of the N(R)Ar ligands is oriented with its *tert*-butyl group distal from phosphorus, presenting an Ar π cloud to the methyl substituent on phosphorus. The Zr–O–P(Me)–Mo core in **4** is reminiscent of Zr–O–C(R′)–M cores in zirconoxy carbenes prepared by addition of Zr–R′ (R′ = H, alkyl) bonds across the CO unit in metal carbonyl complexes.³⁵

We thank Monsanto Company for support of this work, and Michael K. Stern for helpful discussions. M. J. A. J. thanks NSERC for a predoctoral fellowship.

Footnotes and References

* E-mail: ccummins@mit.edu

† Alfred P. Sloan Fellow, 1997–2000.

‡ All manipulations were carried out under an atmosphere of dry nitrogen using solvents purified by standard methods. The new compounds **2** and **4** gave satisfactory elemental analyses (C, H and N).

§ Crystallographic data: For **2**: triclinic, space group $P\bar{1}$, $a = 10.7615(8)$, $b = 11.4942(9)$, $c = 15.1899(11)$ Å, $\alpha = 89.865(2)$, $\beta = 80.838(2)$, $\gamma = 89.4900(10)$ °, $U = 1854.9(2)$ Å³, $Z = 2$, 2988 unique reflections, data/

parameter ratio = 7.9, $R_1 = 0.1207$, $wR_2 = 0.2683$, GOF 1.050, residuals based on $I > 2\sigma(I)$.

For **4**: orthorhombic, space group $Pbca$, $a = 11.3996(2)$, $b = 23.2075(4)$, $c = 39.2443(7)$ Å, $U = 10382.3(3)$ Å³, $Z = 8$, 7456 unique reflections, data/parameter ratio 15, $R_1 = 0.0669$, $wR_2 = 0.2023$, GOF = 1.269, residuals based on $I > 2\sigma(I)$. CCDC 182/527.

- O. J. Scherer, J. Braun, P. Walther, G. Heckmann and G. Wolmershäuser, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 852.
- W. A. Herrmann, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 818.
- J. F. Corrigan, S. Doherty, N. J. Taylor and A. J. Carty, *J. Am. Chem. Soc.*, 1994, **116**, 9799.
- R. M. Atalla and P. D. Singh, *Astrophys. Space Sci.*, 1987, **133**, 267.
- G. A. Blake, E. C. Sutton, C. R. Masson and T. G. Phillips, *Astrophys. J.*, 1987, **315**, 621.
- H. E. Matthews, P. A. Feldman and P. F. Bernath, *Astrophys. J.*, 1987, **312**, 358.
- Z. Mielke, M. McCluskey and L. Andrews, *Chem. Phys. Lett.*, 1990, **165**, 146.
- H.-B. Qian, P. B. Davies and P. A. Hamilton, *J. Chem. Soc., Faraday Trans.*, 1995, **91**, 2993.
- L. R. Thorne, V. G. Anicich, S. S. Prasad and W. T. Huntress Jr., *Astrophys. J.*, 1984, **280**, 139.
- R. J. Vanzee and A. U. Khan, *Chem. Phys. Lett.*, 1975, **36**, 123.
- C. E. Laplaza, W. M. Davis and C. C. Cummins, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 2042.
- N. C. Zanetti, R. R. Schrock and W. M. Davis, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 2044.
- G. Wu, D. Rovnyak, M. J. A. Johnson, N. C. Zanetti, D. G. Musaev, K. Morokuma, R. R. Schrock, R. G. Griffin and C. C. Cummins, *J. Am. Chem. Soc.*, 1996, **118**, 10 654.
- M. Scheer, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 1997.
- M. Scheer, J. Müller and M. Häser, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 2492.
- R. W. Murray, *Chem. Rev.*, 1989, **89**, 1187.
- W. Adam, R. Curci and J. O. Edwards, *Acc. Chem. Res.*, 1989, **22**, 205.
- W. Adam, J. Bialas and L. Hadjarapoglou, *Chem. Ber.*, 1991, **124**, 2377.
- M. Regitz and O. J. Scherer, *Multiple Bonds and Low Coordination in Phosphorus Chemistry*, Georg Thieme Verlag, Stuttgart, 1990.
- A. R. Johnson, P. W. Wanandi, C. C. Cummins and W. M. Davis, *Organometallics*, 1994, **13**, 2907.
- M. J. A. Johnson, P. M. Lee, A. L. Odom, W. M. Davis and C. C. Cummins, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 87.
- C. E. Laplaza, W. M. Davis and C. C. Cummins, *Organometallics*, 1995, **14**, 577.
- C. E. Laplaza, A. L. Odom, W. M. Davis, C. C. Cummins and J. D. Protasiewicz, *J. Am. Chem. Soc.*, 1995, **117**, 4999.
- C. E. Laplaza, M. J. A. Johnson, J. C. Peters, A. L. Odom, E. Kim, C. C. Cummins, G. N. George and I. J. Pickering, *J. Am. Chem. Soc.*, 1996, **118**, 8623.
- J. C. Peters, A. R. Johnson, A. L. Odom, P. W. Wanandi, W. M. Davis and C. C. Cummins, *J. Am. Chem. Soc.*, 1996, **118**, 10 175.
- W. Kutzelnigg, *Angew. Chem., Int. Ed. Engl.*, 1984, **23**, 272.
- J. E. Butler, K. Kawaguchi and E. Hirota, *J. Mol. Spectrosc.*, 1983, **101**, 161.
- H.-B. Qian, *J. Mol. Spectrosc.*, 1995, **174**, 599.
- N. N. Greenwood and A. Earnshaw, *Chemistry of the Elements*, Pergamon, Oxford, 1984.
- J. A. Davies, S. Dutremez and A. A. Pinkerton, *Inorg. Chem.*, 1991, **30**, 2380.
- M. J. A. Johnson, A. L. Odom and C. C. Cummins, manuscript in preparation.
- E. O. Fischer, in *Nobel Lectures in Chemistry*, ed. T. Frängsmyr, World Scientific, Singapore, 1993, p. 105.
- E. Samuel and M. D. Rausch, *J. Am. Chem. Soc.*, 1973, **95**, 6263.
- R. T. Baker, P. J. Krusic, T. H. Tulip, J. C. Calabrese and S. S. Wreford, *J. Am. Chem. Soc.*, 1983, **105**, 6763.
- P. T. Wolczanski and J. E. Bercaw, *Acc. Chem. Res.*, 1980, **13**, 121.

Received in Bloomington, IN, USA, 7th May 1997; 7/03105J