

Chemoenzymatic synthesis of a novel ligand for rhodium-catalysed asymmetric hydrogenation

Brian Adger,^a Ulrich Berens,^a Matthew J. Griffiths,^b Michael J. Kelly,^c Ray McCague,^a John A. Miller,^c Christopher F. Palmer,^a Stanley M. Roberts,^{*b} Rüdiger Selke,^d Ute Vitinius^b and Guy Ward^c

^a Chiroscience plc, Cambridge, UK CB4 4WE

^b Department of Chemistry, University of Liverpool, Liverpool, UK L69 7ZD


^c Department of Chemistry, University of Exeter, Exeter, UK EX4 4QD

^d Max Planck-Gesellschaft Zur Förderung der Wissenschaften e.v., University of Rostock, D-18055 Rostock, Germany

The hydrogenation of alkenes **7a–g** using a chiral rhodium catalyst **6** (based on a bicyclo[3.2.0]heptane framework) takes place to give the phenylalanine derivatives **8a–g** with remarkably high stereoselectivity (59–92% ee).


The relatively inflexible, angular shape of the bicyclo[3.2.0]heptane ring system means those substituents on the concave (*endo*) face of the molecule are positioned close together in space. Molecular modelling studies suggested that the hydroxy groups in bicyclo[3.2.0]heptan-3-*endo*, 6-*endo*-diol would be positioned in such a way that derivitization to form bis(diarylphosphinyl) groups may provide a novel chiral ligand for organorhodium-catalysed hydrogenation reactions.¹

Optically pure diol **1** may be prepared from (–)-bicyclo[3.2.0]hept-2-en-6-one² **2** by the route described in Scheme 1. Thus, formation of the bromohydrin **3**, hydrodebromination (furnishing **4**) and stereocontrolled reduction by delivery of hydride ion from the more exposed *exo*-face afforded the diol **1** in 50% overall yield.



Scheme 1 Reagents and conditions: i, *N*-Bromosuccinimide, acetone, H₂O, 6 h, 80%; ii, Bu₃SnH, AIBN, toluene, heat, 82%; iii, NaBH₄, MeOH, –78 °C, 76%

However, since both enantiomers of the diol **1** were required as intermediates to the chiral ligands it seemed more sensible to perform a kinetic resolution further along the synthetic sequence. Kinetic resolution of the racemic alcohols **1** and **4** using lipase-catalysed acylation reactions were unsuccessful. The bromohydrin (±)-**3** was resolved by employing *Pseudomonas fluorescens* lipase in vinyl acetate⁴ but a more satisfactory



Scheme 2 Reagents and conditions: i, Yeast, H₂O, 6 h, 100% crude yield; ii, Bu₃SnH, AIBN, toluene, heat, 86%





Fig. 1 Delivery of hydride to **3**

and efficient biocatalytic method utilised a yeast-catalysed highly enantio- and stereo-selective reduction of racemic ketone **3** (Scheme 2).

It is noteworthy that reduction takes place to give only the (6*S*)-bromodiol **5**, unlike the corresponding non-enantioselective bioreduction of the parent compound **2** which gave (1*S*,5*R*,6*S*)-bicyclo[3.2.0]hept-2-en-6-*endo*-ol and (1*R*,5*S*,6*S*)-bicyclo[3.2.0]hept-2-en-6-*exo*-ol in almost equal quantities.⁵ Presumably delivery of the hydride ion from the concave face of (+)-**3** is prohibited by unfavourable steric interactions within the active site of the dehydrogenase (Fig. 1).

The diol (–)-**1** was converted into the rhodium complex **6** using standard methodology, and this complex was used to catalyse the asymmetric hydrogenation of compounds **7a–g** to afford the (*R*)-phenylalanine derivatives **8a–g** in excellent yield and moderate to good enantiomeric excess (Table 1). Indeed,

Table 1 Asymmetric hydrogenation of some enamide esters, **7a–g**



Starting material	Product	Enantiomeric excess ^a (%)	Yield (%)
7a	8a	81	> 95
7b	8b	73	> 95
7c	8c	76	> 95
7d	8d	59	> 95
7e	8e	61	> 95
7f	8f	66	> 95
7g	8g	77	> 95

^a Estimation by HPLC using a Chirasil-Val column.

Table 2 Asymmetric hydrogenation of enamide ester **7a**

<i>T</i> /°C	Volume of solvent/ml	Enantiomeric excess of product (%)
25	15	81
12	15	83.5
2	15	87.5
25	10	88
25	5	91.5
2	5	92.5

given the subtlety of the chirality of the bicyclo[3.2.0]heptane framework, the observed level of enantioselectivity is surprisingly high. The reaction rates were perfectly satisfactory (the reaction was completed in 2–10 min) using 1 mmol substrate and 0.01 mmol catalyst in MeOH (15 ml) under an atmosphere of hydrogen. Using similar conditions, itaconic acid was reduced to (2*S*)-methylsuccinate in quantitative yield and 80.5% ee.

The reaction conditions for the asymmetric reduction of substrate **7a** were investigated further. Lowering the temperature of the reaction and increasing the concentration of the reactants both proved to be beneficial, with a satisfactory 92.5% ee for product **8a** being obtained at low temperature and high concentration (Table 2).[†] The concentration effect may be explained by the effect on the rate of delivery of hydrogen to the catalyst, which will be slower in the reactions using small amounts of solvent. Thus, in the more concentrated solution the catalyst may experience a lower 'effective' hydrogen concentration.⁶

We thank the BBSRC for a Fellowship (to U. V.) and a CASE award (to M. J. G.), Chiroscience, and EPSRC and the University of Exeter for studentships (to J. M. and G. W.) and Donna Blackmond for enlightening discussions.

Footnotes and References

* E-mail: sj11@liverpool.ac.uk

[†] Experimental procedure for the asymmetric hydrogenation of compound **7a**. A mixture of complex **6** (8 mg, 0.01 mmol) and α -acetamidocinnamic acid **7a** (205 mg, 1 mmol) was cooled (2 °C) under a nitrogen atmosphere. Dry, degassed MeOH (5 ml) was added to the mixture, which was stirred vigorously. The flask was evacuated, filled with hydrogen and stirred for 10 min. A 0.5 ml aliquot was removed from the reaction vessel and treated with Me₃SiCHN₂ (2.0 M solution in hexane) in order to convert the acid to the methyl ester. The solvent was evaporated and the residue was treated with diethyl ether; filtration and evaporation gave a crude residue (21 mg). The enantiomeric excess of the product **8a** was determined to be 92.5% by chiral gas chromatography, using a Chirasil-Val-III column.

- 1 Review: R. Noyori, *Asymmetric Catalysts in Organic Synthesis*, Wiley, New York, 1994, ch. 2; recent publications: G. Zhu, P. Cao, Q. Jiang and X. Zhang, *J. Am. Chem. Soc.*, 1997, **119**, 1799; R. Selke, M. Ohff and A. Riepe, *Tetrahedron*, 1996, **52**, 15079; M. J. Burk, Y. M. Wang and J. R. Lee, *J. Am. Chem. Soc.*, 1996, **118**, 5142; H.-J. Kreuzfeld, U. Schmidt, C. Döbler and H. W. Krause, *Tetrahedron: Asymmetry*, 1996, **7**, 1011; C. Dobler, H.-J. Kreuzfeld, M. Michalik and H. W. Krause, *Tetrahedron: Asymmetry*, 1996, **7**, 117; J. Albrech and U. Nagel, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 407.
- 2 K. Tanaka, O. Kakinoki and F. Toda, *Tetrahedron: Asymmetry*, 1992, **3**, 517.
- 3 Z. Grudzinski and S. M. Roberts, *J. Chem. Soc., Perkin Trans. 1*, 1975, 1767.
- 4 For enzymatic resolution of related bromohydrins, see I. C. Cotterill, A. G. Sutherland, S. M. Roberts, R. Grobbauer, J. Spreitz and K. Faber, *J. Chem. Soc., Perkin Trans. 1*, 1991, 1365; I. C. Cotterill and S. M. Roberts, *J. Chem. Soc., Perkin Trans. 1*, 1992, 2585.
- 5 R. F. Newton, J. Paton, D. P. Reynolds, S. Young and S. M. Roberts, *J. Chem. Soc., Chem. Commun.*, 1979, 908; J. Davies, S. M. Roberts, D. P. Reynolds and R. F. Newton, *J. Chem. Soc., Perkin Trans. 1*, 1981, 1317; See also G. Fantin, M. Fogagnolo, A. Medici, P. Pedrini, E. Marotta, M. Monti and R. Righi, *Tetrahedron: Asymmetry*, 1996, **7**, 277.
- 6 Y. Sun, R. N. Landau, J. Wang, C. LeBlond and D. G. Blackmond, *J. Am. Chem. Soc.*, 1996, **118**, 1348.

Received in Cambridge, UK, 13th June 1997; 7/04136E