

Photopromoted oxidative cyclization of an *o*-phenylene-bridged Schiff base via a manganese(III) complex, leading to a fluorescent compound, 2-(2-hydroxyphenyl)benzimidazole

Takashi Fukuda,^a Fuminori Sakamoto,^a Minoru Sato,^b Yoshiharu Nakano,^a Xiang Shi Tan^a and Yuki Fujii^{*a†}

^a Department of Chemistry, Faculty of Science, Ibaraki University, Mito, 310-8512, Japan

^b Department of Chemistry and Engineering, Ibaraki National College of Technology, Hitachinaka 344-1200, Japan

Visible light photolysis of [N,N'-*o*-phenylenebis(salicylideneamino)]diaquamanganese(III) resulted in the fluorescent compound 2-(2-hydroxyphenyl)benzimidazole by a one electron redox reaction between Mn^{III} and the Schiff base ligand, followed by the base-hydrolysis of the oxidized Schiff base, and then the cyclization of the base-hydrolyzed product.

The photoactivation of manganese complexes is an attractive theme both in the modeling of PSII and in the development of a new photoreaction.^{1–4} It has been known that, upon visible light irradiation, in the presence of *p*-quinone, manganese(III) complexes with tetradentate Schiff base ligands having the salen-type skeleton generate molecular oxygen;^{5–7} in the absence of *p*-quinone, rearrangement of the coordinated Schiff base ligand occurs.⁸ Recently, we examined the photolysis of [N,N'-*o*-phenylenebis(salicylideneamino)]diaquamanganese(III) **1** and observed a photopromoted oxidative cyclization of the coordinated Schiff base ligand, and the effective formation of the fluorescent compound 2-(2-hydroxyphenyl)benzimidazole **2**, which is useful as a laser dye, high-energy radiation detector, molecular energy storage system, and fluorescent probe.^{9–11} **2** is usually prepared from salicylic acid and *o*-phenylenediamine in polyphosphoric acid at 190 °C in 14% yield.¹² This photoreaction thus provides a new and effective synthetic method for the fluorescent compound **2** and its derivatives.[‡]

Cation **1** was prepared by a literature method,¹³ and isolated as its perchlorate salt. § Absence of Mn^{II} was confirmed by EPR spectroscopy. Slow evaporation of methanol solution of the complex produced crystals suitable for single crystal X-ray structure analysis. ¶ An ORTEP drawing of **1** is shown in Fig. 1. The structure of the complex cation comprises a planar tetradentate Schiff base ligand tightly bound to the Mn^{III} center via two Mn–N bonds [1.981(2), 1.978(2) Å] and two Mn–O bonds [1.859(1), 1.875(1) Å], and by axial methanol and water molecules which complete an octahedron around the Mn^{III} ion. Longer Mn–O distances to the apical ligands [2.272(2) Å for Mn–H₂O and 2.291(2) Å for MeOH] may be partly attributed to the Jahn–Teller effect for the d⁴ ion, although it is common for




Fig. 1 An ORTEP drawing of cation **1** with thermal ellipsoids at 50% probability. Hydrogen atoms are omitted for clarity.

solvated Mn–O distances to be longer than chelated ones.¹⁴ No hydrogen bond between the phenolic oxygen atoms of the Schiff base ligand and apical water or methanol was observed.⁶

Fig. 2 shows the time courses of UV–VIS (320 nm), fluorescence (432 nm), and EPR spectra of **1** in water at initial pH = 6.4 under irradiation with visible light (150 W tungsten halogen lamp with UV cut filter L-39). The UV–VIS spectral change exhibited an isosbestic point at 223 nm, and the half life ($t_{1/2}$) of **1** was 11–15 h at pH 6.4. || A similar spectral change also occurred in the dark but was very slow ($t_{1/2} \approx 55$ h). Our preliminary experiments revealed that the photopromotion occurs by the irradiation into the CT band (400 ± 20 nm), but hardly for the $\pi \rightarrow \pi^*$ (335 ± 20 nm) and $d \rightarrow d$ (540 ± 20 nm) bands. Further, the reaction becomes faster with increase of solution pH ($t_{1/2} \approx 9$ h at pH 7.5, ≈ 5 h at pH 8.6).

On the other hand, the fluorescence and EPR intensities of the photoreaction solution increased with the decay of **1**, indicating that the fluorescent compound, assigned as 2-(2-hydroxyphenyl)benzimidazole **2**, was formed at early stages of the reaction. The EPR spectra show typical six-line Mn^{II} signals ($g = 2.013$), which proves that the Mn^{III} was reduced to Mn^{II} during this reaction. Since neither molecular oxygen nor hydrogen peroxide were detected in the photoreaction solution, it is strongly suggested that intramolecular one electron redox occurred between Mn^{III} and the coordinated Schiff base ligand. A similar photoreaction also occurred under vacuum, however, no EPR signal corresponding to an organic radical was observed.

From the photoreaction solution, **2** and salicylaldehyde **3** could be separated both in about 40% yield by extraction with ethyl acetate, followed by silica-gel column separation using ethyl acetate–*n*-hexane (1 : 2) as eluent ($R_f = 0.6$ for **2**, $R_f = 0.8$ for **3**).** In addition to **2** and **3**, under argon, N,N'-





Fig. 2 Time courses of UV–VIS, fluorescence and X-band EPR spectra of the photolysis solution of **1** at pH = 6.4 and 25 °C. The complex concentration is 2.0×10^{-5} mol dm⁻³ for UV–VIS (●) and fluorescence (○) and 1.0×10^{-3} mol dm⁻³ for EPR (□). Fluorescence: $\lambda_{\text{ex}} = 325$ nm, $\lambda_{\text{em}} = 432$ nm, EPR: microwave power = 5 mW, microwave frequency = 9.238 GHz, 0.5 mT modulation amplitude, 25 °C.



Scheme 1

o-phenylenebis(salicylideneimine) **4** was isolated in *ca.* 10% yield as a photoreaction product of **1**.

On the basis of these facts, the reaction mechanism (Scheme 1) is suggested as follows: (i) Mn^{III}-Schiff base complex **A** is reduced to Mn^{II} complex **B** with partially oxidized Schiff base ligand, for which visible light promotes the reaction (the reaction has been shown to be enhanced by excitation at the CT band of **1**, probably by LMCT). **B** is thought to be a transient radical and a similar radical has been proposed by Floriani and coworkers.¹⁵ (ii) **B** is base-hydrolyzed to form radical **C**, followed by cyclization of the partially oxidized Schiff base ligand to produce radical **D** and Mn^{II} complex **E**, **D** is oxidized by O₂ to produce **F**. These processes are very fast, since final products **F** and Mn^{II} appear from the start of the reaction with no organic radical being detected in the timescale of EPR measurement. No detection of radical **D** under vacuum may be due to a redox reaction between **A** and **D**, which results in the formation of the corresponding Mn^{II}-Schiff base complex and **F**. This is partly supported by the fact that the Schiff base ligand **4** is detected under argon but not under air. Evidence for base-

hydrolysis of **B** is the formation of salicylaldehyde **G** and rate enhancement with increase of solution pH. A study of the detailed reaction mechanism is under way.

This work was partly supported by a Grant-in-Aid for Science Research (No. 09440224 and No. 97370) from the Ministry of Education, Science and Culture of Japan.

Notes and References

† E-mail: yuki@mito.ipc.ibaraki.ac.jp

‡ This method is also applicable for the preparation of the cationic fluorescent compound 2-(2-hydroxy-5-trimethylammoniummethylphenyl)-benzimidazole chloride.

§ This complex contains methanol as apical ligand in the crystal state, however the methanol is thought to be substituted by water in aqueous solution.

¶ *Crystal data:* C₂₁H₂₀N₂O₈MnCl, *M* = 518.79, monoclinic, space group *P*2₁/*c*, *Z* = 4, *a* = 11.804(2), *b* = 13.435(2), *c* = 14.358(2) Å, β = 109.108(9)°, *U* = 2151.5(5) Å³, *D_c* = 1.601 g cm⁻³, Mo-Kα radiation (λ = 0.710 73 Å), *R* = 0.033 and *R_w* = 0.032 for 3585 observed reflections with *I* > 3σ(*I*). CCDC 182/878.

|| Since the reaction occurs with irregular induction time, *t*_{1/2} is variable.
** 2: ¹H NMR [(CD₃)₂SO, TMS standard], δ 13.18 (br, 1 H), 8.057 (d, 1 H), 7.664 (d, 2 H), 7.400 (t, 1 H), 7.303 (br, 2 H), 7.074–7.019 (m, 2 H). ¹³C NMR [(CD₃)₂SO], δ 158.07, 151.68, 140.93, 133.14, 126.19, 123.27, 122.38, 119.11, 117.93, 112.58, 111.51. EI-MS: *m/z* 210 (*M*⁺)

- 1 A. Harriman, *Coord. Chem. Rev.*, 1979, **28**, 147.
- 2 V. L. Pecoraro, *Photochem. Photobiol.*, 1988, **48**, 247.
- 3 K. Wieghardt, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 1153.
- 4 V. L. Pecoraro, M. J. Baldwin and A. Gelasco, *Chem. Rev.*, 1994, **94**, 807.
- 5 F. M. Ashmawy, C. A. McAuliffe, R. V. Parish and J. Tames, *J. Chem. Soc., Dalton Trans.*, 1985, 1391.
- 6 N. Aurangzeb, C. E. Hulme, C. A. McAuliffe, R. G. Pritchard, M. Watkinson, M. R. Bermejo, A. Garcia-Deibe, M. Rey, J. Sanmartin and A. Sousa, *J. Chem. Soc., Chem. Commun.*, 1994, 1153.
- 7 M. Watkinson, A. Whiting and C. A. McAuliffe, *J. Chem. Soc., Chem. Commun.*, 1994, 2141.
- 8 A. Garcia-Deibe, A. Sousa, M. R. Bermejo, P. P. MacRory, C. A. McAuliffe, R. G. Pritchard and M. Helliwell, *J. Chem. Soc., Chem. Commun.*, 1991, 728.
- 9 D. L. Williams and A. Heller, *J. Phys. Chem.*, 1970, **74**, 4473.
- 10 A. U. Acuna, F. Amat, J. Catalan, A. Costela, J. M. Figuera and J. M. Munoz, *Chem. Phys. Lett.*, 1986, **132**, 567.
- 11 E. L. Roberts, J. Dey and I. M. Warner, *J. Phys. Chem., A*, 1997, **101**, 5296.
- 12 C. M. Oelando Jr., J. G. Wirth and D. R. Heath, *J. Org. Chem.*, 1970, **35**, 3147.
- 13 L. J. Boucher and C. G. Coe, *Inorg. Chem.*, 1975, **14**, 1289.
- 14 C. A. McAuliffe, A. Nabhan, R. G. Pritchard, M. Watkinson, M. Bermejo and A. Sousa, *Acta Crystallogr., Sect. C*, 1994, **50**, 1676.
- 15 E. Gallo, E. Solari, N. Re, C. Floriani, A. Chiesi-Villa and C. Rizzoli, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1981.

Received in Cambridge, UK, 30th March 1998; 8/02405G