

# Self-assembly of a bis-tridentate $\text{Py}_2\text{S}_4$ ligand and cadmium cation into 1- and 2-D coordination networks†

Marielle Loi,<sup>a</sup> Ernest Graf,<sup>a</sup> Mir Wais Hosseini,<sup>\*a</sup> André De Cian<sup>b</sup> and Jean Fischer<sup>b</sup>

<sup>a</sup> Laboratoire de Chimie de Coordination Organique and <sup>b</sup> Laboratoire de Cristallographie et Chimie Structurale, Université Louis Pasteur, F-67000 Strasbourg, UMR CNRS 7513, France. E-mail: hosseini@chimie.u-strasbg.fr

Received (in Basel, Switzerland) 18th January 1999, Accepted 16th February 1999

Using a  $\text{Py}_2\text{S}_4$  bis-tridentate ligand and  $\text{Cd(II)}$ , 1- and 2-D coordination networks were obtained and characterised in the solid state; whereas a 1-D network was obtained using the cadmium(II) dichloride salt, a 2-D network was obtained using a mixed chloride and tetrafluoroborate cadmium(II) salt; in the latter case, the inclusion of two  $\text{CHCl}_3$  molecules within cavities formed by the organic and inorganic fragments was observed.


For some time now, intensive research on the design and preparation of coordination networks or polymers has been performed.<sup>1</sup> Although, nearly all cases reported deal with bis- or tris-monodentate systems,<sup>2</sup> examples of coordination polymers using bis-bidentate ligands have also been published.<sup>3,4</sup> On the other hand, only a few examples of characterised systems based on bis-tridentate ligands are known.<sup>5,6</sup>

The formation of coordination networks in the crystalline phase takes place under self-assembly conditions which require, on one hand, the reversible formation of the assembling core by complexation processes and its translation and, on the other hand, the conformity to the packing forces within the crystalline phase. Thus, for simple coordination networks, the choice of the two partners (metal, ligand) is the dominant feature. For porous coordination networks, in addition to the two partners mentioned before, one should also take into account the guest molecules initially occupying the pores. Owing to the fact that tridentate centres form rather stable and non-labile complexes with transition metals, the choice of the metal and the ligand is not obvious and therefore, not many characterised examples are available so far.

Here, we report the synthesis of a new bis-tridentate ligand and its self-assembly into 1- and 2-D coordination polymers using  $\text{Cd(II)}$  cation.

The design of the linear bis-tridentate ligand **7** (Scheme 1) is based on a combination of two pyridine moieties and four thioether groups. The two tridentate ligands are interconnected by a phenyl ring. Examples of other bis-tridentate ligands have been reported.<sup>7</sup>

Dealing with the metal cation, it has been shown previously that **1** forms a binuclear  $\text{Cd(II)}$  complex in which the two metal centres are bridged by two  $\text{Cl}^-$  anions.<sup>8</sup> This type of doubly bridged metal system has been reported for  $\text{Co(II)}$ <sup>6b</sup> and for  $\text{Ni(II)}$  complexes.<sup>9</sup>



Scheme 1

The synthesis of **7** was achieved starting from chelidamic acid **2**, which was first transformed into **3**.<sup>10</sup> The latter was reduced to **4**.<sup>11</sup>  $\text{Pd(PPh}_3)_4$  catalysed coupling reaction of **4** with 1,4-phenyldiboronic acid<sup>12</sup> in dry THF gave **5** in 65% yield. Bromination of the latter using 33%  $\text{HBr-AcOH}$  afforded **6** in 61% yield. Finally, the treatment of **6** with  $\text{NaSMe}$  in dry THF afforded **7** in 71% yield. The latter, in addition to spectroscopic methods, was characterised by X-ray crystallography.


Upon slow diffusion at room temp. of a MeOH solution containing  $\text{CdCl}_2 \cdot x\text{H}_2\text{O}$  into a  $\text{CHCl}_3$  solution of **7** (1 : 1 ratio of Cd : **7**), colourless crystals were deposited after 24 h. The X-ray study<sup>‡</sup> showed the following features (Fig. 1): the crystal **I** (monoclinic) was composed of **7**,  $\text{Cd(II)}$ ,  $\text{Cl}^-$  anions,  $\text{CHCl}_3$  and  $\text{H}_2\text{O}$  molecules. **7**,  $\text{Cd(II)}$  and  $\text{Cl}^-$  anions formed a neutral, infinite 1-D coordination network (Fig. 2).

Depending on the definition of the assembling core, the 1-D network  $(7\text{Cd}_2\text{Cl}_4)_n$  thus obtained may be described in two different ways. Considering the metallic core as a binuclear  $\text{Cd}_2\text{Cl}_4$  unit (Fig. 2), the network may be regarded as resulting from the binding of two such units by the two  $\text{PyS}_2$  fragments of **7** leading thus to the exo-tetranuclear  $\text{Cd(II)}$  complex and the repetition of the binding process leads to the 1-D coordination network. The same system may be also described as resulting from the bridging of consecutive exo-binuclear  $7\text{-Cd}_2$  units by two  $\text{Cl}^-$  anions, and the remaining other two  $\text{Cl}^-$  anions behaving as ancillary ligands. For the ligand part, the pyridine rings were tilted by  $-37.6$  and  $38.3^\circ$  with respect to the phenyl group. The CS and CN distances were roughly the same as those observed for free **7**. The coordination sphere around the Cd cations was composed of one N atom ( $d_{\text{NCd}} = 2.426 \text{ \AA}$ ), two S atoms ( $d_{\text{SCd}} = 2.730, 2.747 \text{ \AA}$ ) and three  $\text{Cl}^-$  anions. Among the three  $\text{Cl}^-$  present, two of them were shared between two Cd centres ( $d_{\text{ClCd}} = 2.591, 2.651 \text{ \AA}$ ) whereas the third one was acting as an ancillary ligand ( $d_{\text{ClCd}} = 2.511 \text{ \AA}$ ). The coordination geometry around the metal centre was distorted octahedral with S–Cd–S, Cl–Cd–Cl and N–Cd–Cl angles of  $146.5$ ,  $172.0$  and  $175.2^\circ$ , respectively. Within the  $\text{Cd}_2 \cdots \text{Cl}_4$  unit, the Cd  $\cdots$  Cd distance was found to be  $3.946 \text{ \AA}$ . The  $\text{CHCl}_3$  and  $\text{H}_2\text{O}$  molecules were localised between the 1-D chains.


Interestingly, when instead of using  $\text{CdCl}_2$ , a mixed  $\text{Cl}^-$  and  $\text{BF}_4^-$  salt  $[\text{CdCl}(\text{BF}_4)]$  was used, the same diffusion method afforded after two days another type of colourless crystal. The solid state analysis (Fig. 3)<sup>‡</sup> showed that the crystal **II** was composed of **7**,  $\text{Cd(II)}$  cations,  $\text{Cl}^-$  and  $\text{BF}_4^-$  anions,  $\text{CHCl}_3$  and MeOH molecules. **7**,  $\text{Cd(II)}$  and  $\text{Cl}^-$  anions formed a cationic infinite 2-D coordination network (Fig. 2). The latter may be described as the result of mutual interconnection of the 1-D coordination polymers described above through  $\text{Cl}^-$  anions acting now as bridging ligands. In a sense, the poor coordination ability of  $\text{BF}_4^-$  has created a  $\text{Cl}^-$  deficiency in the coordination sphere of  $\text{Cd(II)}$ , made up for by an additional  $\text{Cl}^-$  bridging interaction.

Thus, the 2-D network may be regarded as the translation into two different space directions of an assembling core which may be defined  $[\text{7}_4\text{Cd}_4\text{Cl}_7]^+$  (Fig. 2). The charge neutrality being achieved by  $\text{BF}_4^-$  anions present in the lattice. Owing to the aromatic nature of the ligand used, the 2-D network was composed of cavities delimited in an alternating fashion by


† Dedicated to Professor John A. Osborn on the occasion of his 60th birthday.



**Fig. 1** A section of the X-ray structure of the 1-D coordination network **I** formed between **7** and  $\text{CdCl}_2$ . H atoms, anions and solvent molecules are not presented for clarity.



**Fig. 2** Partial representation of the 1- and 2-D coordination polymers **I** and **II**.



**Fig. 3** A section of the X-ray structure of the 2-D coordination network **II** (see text). H atoms, anions and MeOH molecules are not presented for clarity.

organic and inorganic fragments. For **7**, the pyridine rings were tilted by  $-30.7$  and  $32.2^\circ$  with respect to the phenyl group. The C–S and C–N distances were roughly the same as those observed for the free **7** and for the above-mentioned 1-D network. The coordination sphere around the Cd cations was again composed of one N atom (average  $d_{\text{NCd}} = 2.385 \text{ \AA}$ ), two S atoms (average  $d_{\text{SCd}} = 2.697 \text{ \AA}$ ) and three  $\text{Cl}^-$  anions. Although all three  $\text{Cl}^-$ , present in the coordination sphere of Cd cation, adopted a bridging role between the metallic centres, two of them were shared between two Cd centres (average  $d_{\text{ClCd}} = 2.608 \text{ \AA}$ ) within the 1-D network, whereas the third one bridged two consecutive linear networks ( $d_{\text{ClCd}} = 2.559 \text{ \AA}$ ) thus leading to a two-dimensional polymer. The coordination geometry around the metal centre was again distorted octahedral with S–Cd–S, Cl–Cd–Cl and N–Cd–Cl angles of  $147.6$ ,  $169.0$  and  $163.6^\circ$ , respectively. With the  $\text{Cd}_4\text{Cl}_7$  unit, the Cd...Cd distances varied from  $3.921 \text{ \AA}$  for the doubly bridged cations to  $4.625 \text{ \AA}$  for the singly bridged centres. Interestingly, whereas MeOH and  $\text{BF}_4^-$  anions were localised between the

2-D sheets, the  $\text{CHCl}_3$  molecules were included within the cyclophane type cavities composing the 2-D network.

In conclusion, using the  $\text{N}_2\text{S}_4$  bis-tridentate ligand and  $\text{Cd(II)}$ , 1- and 2-D coordination networks were obtained and characterised in the solid state. For the 2-D network the inclusion of two  $\text{CHCl}_3$  molecules within cavities formed by the organic and inorganic fragments was observed. Based on the same strategy, the formation of magnetic networks using paramagnetic metal cations is currently under investigation.

## Notes and references

‡ *Crystal data*: **I** (colorless, 294 K),  $\text{C}_{12}\text{H}_{14}\text{CdCl}_2\text{NS}_2 \cdot \text{CHCl}_3 \cdot 2\text{H}_2\text{O}$ ,  $M = 575.09$ , monoclinic,  $a = 29.037(1)$ ,  $b = 8.4340(2)$ ,  $c = 22.5310(8) \text{ \AA}$ ,  $\beta = 126.63(1)$ ,  $U = 4428(2) \text{ \AA}^3$ ,  $Z = 8$ , space group  $C2/c$ ,  $D_c = 1.73 \text{ g cm}^{-3}$ , Nonius Kappa CCD, Mo-K $\alpha$ ,  $\mu = 1.785 \text{ mm}^{-1}$ , 5315 data with  $I > 3\sigma(I)$ ,  $R = 0.056$ ,  $R_w = 0.076$ .

**II** (colorless, 173 K),  $\text{C}_{24}\text{H}_{28}\text{Cd}_2\text{Cl}_3\text{N}_2\text{S}_4 \cdot \text{BF}_4 \cdot 2\text{CHCl}_3 \cdot \text{CH}_3\text{OH}$ ,  $M = 1161.52$ , orthorhombic,  $a = 16.3181(4)$ ,  $b = 14.8660(3)$ ,  $c = 35.9037(8) \text{ \AA}$ ,  $U = 8709.7(6) \text{ \AA}^3$ ,  $Z = 8$ , space group  $Pbcn$ ,  $D_c = 1.77 \text{ g cm}^{-3}$ , Nonius Kappa CCD, Mo-K $\alpha$ ,  $\mu = 1.764 \text{ mm}^{-1}$ , 4036 data with  $I > 3\sigma(I)$ ,  $R = 0.051$ ,  $R_w = 0.061$ . CCDC 182/1173.

- 1 R. Robson, in *Comprehensive Supramolecular Chemistry*, ed. J. L. Atwood, J. E. D. Davies, D. D. Macnicol and F. Vögtle, Pergamon, vol. 6, ed. D. D. Macnicol, F. Toda and R. Bishop, 1996, pp. 733; O. M. Yaghi, H. Li, C. Davis, D. Richardson and T. L. Groy, *Acc. Chem. Res.*, 1998, **31**, 474.
- 2 T. L. Hennigar, D. C. MacQuarrie, P. Losier, R. D. Rogers and M. J. Zaworotko, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 972 and references therein; M. Loi, M. W. Hosseini, A. Jouaiti, A. De Cian and J. Fischer, *Chem. Commun.*, 1999, submitted.
- 3 O. J. Gelling, F. van Bolhuis and B. L. Faringa, *J. Chem. Soc., Chem. Commun.*, 1991, 917; O. Kahn, Y. Journaux and C. Mathoniere, in *Magnetism: A Supramolecular Function*, ed. O. Kahn, Kluwer, Dordrecht, Netherlands, 1996, vol. C484, pp. 531; S. Decurtins, R. Pellaux, A. Hauser and M. E. von Arx, *ibid.*, p. 487; U. Velten and M. Rehahn, *Chem. Commun.*, 1996, 2639.
- 4 C. Kaes, M. W. Hosseini, C. E. F. Rickard, B. W. Skelton and A. White, *Angew. Chem., Int. Ed.*, 1998, **37**, 920; G. Mislin, E. Graf, M. W. Hosseini, A. De Cian, N. Kyritsakas and J. Fischer, *Chem. Commun.*, 1998, 2545.
- 5 S. J. Loeb and G. K. H. Shimizu, *J. Chem. Soc., Chem. Commun.*, 1993, 1395; A. Neels, B. Mathez Neels and H. Stoeckli-Evans, *Inorg. Chem.*, 1997, **36**, 3402.
- 6 E. C. Constable, A. J. Edwards, D. Philips and P. R. Raithby, *Supramol. Chem.*, 1995, **5**, 93.
- 7 H. A. Goodwin and F. Lions, *J. Am. Chem. Soc.*, 1959, **81**, 6415; J.-P. Sauvage, J.-P. Collin, J.-C. Chambron, S. Guillerez, C. Coudret, V. Balzani, F. Barigelletti, L. De Cola and L. Flamigni, *Chem. Rev.*, 1994, **94**, 993; P. Steenwinkel, H. Kooijman, W. J. J. Smeets, A. L. Spek, D. M. Grove and G. van Koten, *Organometallics*, 1998, **17**, 5411 and references therein.
- 8 F. Teixidor, L. Escriche, I. Rodriguez and J. Casabo, *J. Chem. Soc., Dalton Trans.*, 1989, 1381.
- 9 A. J. Blake, F. Demartin, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis, M. Schröder and G. Verani, *J. Chem. Soc., Dalton Trans.*, 1996, 3705.
- 10 H. Takalo and J. Kankare, *Acta Chem. Scand.*, 1987, **41**, 219.
- 11 H. Takalo, *Acta Chem. Scand.*, 1988, **42**, 373.
- 12 O. C. Musgrave, *J. Chem. Soc. C*, 1970, 488.

Communication 9/00501C