

'Zeofen', a user-friendly oxidizing reagent

Majid M. Heravi,* Dariush Ajami, Kuoumars Aghapoor and Mitra Ghassemzadeh

Chemistry and Chemical Engineering Research Center of Iran, PO Box 14335-186, Tehran, Iran.
E-mail: ccerci@neda.net

Received (in Cambridge, UK) 4th January 1999, Accepted 26th March 1999

A variety of alcohols are oxidized to the corresponding carbonyl compounds in excellent yields by zeofen under classical heating and microwave irradiation in solventless system.

Oxidation of alcohols to carbonyl compounds is a fundamental transformation in organic synthesis and, because of its significant role, the development of newer methods is attracting much current interest^{1–4} in spite of the availability of numerous methods reported in the literature.^{5,6} The use of heavy metals and peracids is very common.^{7,8} However, a drawback of such oxidants and their use in multistage organic synthesis in spite of their power is their lack of selectivity. For instance, over-oxidation of aldehydes to carboxylic acids⁹ and degradation of unsaturated substrates are often unavoidable side reactions.¹⁰ In addition to their lack of selectivity, oxidants based on chromium¹¹ and manganese¹² are corrosive, irritants and toxic.¹³

In view of economic, practical and recent environmental demands,¹⁴ we saw significant advantage in using a non-toxic and inexpensive metal salts such as iron(III) nitrate. Some early success in the use of clayfen for oxidation of alcohols¹⁵ supported this idea.

Reagents impregnated on mineral supports have gained popularity in organic synthesis because of their selectivity and ease of manipulation.^{16,17} Recently we reported the use of zeolite catalyzed¹⁸ and microwave enhanced chemical reactions¹⁹ on organic solid supports.²⁰ We report here a facile oxidation of alcohols to carbonyl compounds using zeofen [short for zeolite HZSM-5 supported iron(III) nitrate] in CH₂Cl₂ at room temperature and under microwave irradiation in a solventless system.

To prepare zeofen, Fe(NO₃)₃·9H₂O (0.80 g, 2 mmol) and a weight equivalent of HZSM-5 zeolite²¹ were crushed together so as to form an intimate mixture (zeofen). This new reagent can be stored in a dark brown bottle and is stable without suffering from the problem of slow decomposition that clayfen does.²² In a typical reaction 1–1.2 equiv. of zeofen was added to a stirring solution of an alcohol in dry CH₂Cl₂ at room temperature. After

the completion of the reactions (monitored by TLC) the reagent was filtered off and washed with CH₂Cl₂. Evaporation of solvent gave the corresponding carbonyl compounds. It should be pointed out that in the absence of zeolite, the reaction was sluggish and even at reflux temperature a considerable amount of starting material was recovered unchanged. Different types of alcohols were converted to their corresponding carbonyl compounds by this procedure in high yields. The results are presented in Table 1.

Although there has been growing interest in applying microwave dielectric heating to accelerate organic reactions,²³ oxidation reactions are less often considered as candidates for microwave irradiation due to their unsafe and uncontrollable experimental conditions.²⁴ It has been shown that polar reactants adsorbed on the surface of solid supports absorb microwaves.²⁵ In connection with our interest in using microwaves to increase the rates of reactions,¹⁹ we have investigated the above oxidation under microwave irradiation. The oxidation reactions were carried out simply by mixing 1–1.2 equiv. of zeofen with 1 equiv. of an alcohol in a beaker under microwave irradiation (900 W) for 2 min. For example, benzyl alcohol was oxidized to benzaldehyde in almost quantitative yields after 10 s. It is noteworthy to mention that in the absence of zeolite, the reaction was slow and, more seriously, molten ferric nitrate and/or its degradation product(s) adhered to the walls of the reaction vessel, forming an intractable solid mass which made the isolation of carbonyl compounds difficult and led to erratic results. Indeed when iron(III) nitrate was used without zeolite the conversion of benzyl alcohol to benzaldehyde was only 50% under conditions otherwise similar to those described above. Clearly zeolite aided the reproducibility and high activity of iron(III) nitrate. To assess the generality of this oxidation a variety of alcohols were reacted under these conditions, which yielded the corresponding carbonyl compounds in excellent yields. Table 2 summarizes the data for oxidation of alcohols under microwave irradiation in a solventless system.

In conclusion, zeofen can be easily prepared from inexpensive and non-toxic materials and serves as an excellent oxidizing agent for various types of alcohols at room tem-

Table 1 Oxidation of alcohols to carbonyl compounds by zeofen in CH₂Cl₂^a

$\begin{array}{c} \text{R}^1 \\ \\ \text{R}^2 - \text{C} - \text{OR} \\ \\ \text{H} \end{array} \xrightarrow[\text{CH}_2\text{Cl}_2 \text{ or microwave}]{\text{zeofen}} \begin{array}{c} \text{R}^1 \\ \diagdown \\ \text{C} = \text{O} \\ \diagup \\ \text{R}^2 \end{array}$			
Alcohol	<i>t</i> /min	Product	Yield ^b (%)
1 PhCH ₂ OH	20	PhCHO	99 (90)
2 4-MeC ₆ H ₄ CH ₂ OH	20	4-MeC ₆ H ₄ CHO	93 (88) ^c
3 2-NO ₂ -5-MeC ₆ H ₃ CH ₂ OH	20	2-NO ₂ -5-MeC ₆ H ₃ CHO	92 (85) ^c
4 (DL)-PhCH(Me)OH	30	PhCO(Me)	89 (80)
5 Ph ₂ CHOH	40	Ph ₂ CO	90 (82)
6 PhCH=CHCH ₂ OH	40	PhCH=CHCHO	77 (67)
7 PhCH ₂ CH(Ph)OH	40	PhCH ₂ COPh	91 (88)
8 Cyclohexanol	60	cyclohexanone	82 (78) ^c
9 2-Methylcyclohexanol	60	2-methylcyclohexanone	85 (76) ^c
10 (–)-Menthol	60	menthone	85 (78)

^a Reaction using 1–1.2 equiv. of zeofen in CH₂Cl₂. The physical data of the isolated products are in agreement with those of authentic samples. ^b Yields are based on GLC analysis. Figures in parenthesis are yields of isolated products. ^c Yields based on isolation of the 2,4-DNPH derivative.

Table 2 Oxidation of alcohols using zeofen under microwave irradiation in a solventless system^a

Alcohol	t/s	Product	Yield ^b (%)
1 PhCH ₂ OH	10	PhCHO	99 (92)
2 4-MeC ₆ H ₄ CH ₂ OH	20	4-MeC ₆ H ₄ CHO	97 (88) ^c
3 2-NO ₂ -5-MeC ₆ H ₃ CH ₂ OH	20	2-NO ₂ -5-MeC ₆ H ₃ CHO	95 (82) ^c
4 (DL)-PhCH(Me)OH	30	PhCO(Me)	90 (82)
5 Ph ₂ CHOH	20	Ph ₂ CO	92 (80)
6 PhCH=CHCH ₂ OH	60	PhCH=CHCHO	81 (87)
7 PhCH ₂ CH(Ph)OH	60	PhCH ₂ COPh	91 (88)
8 Cyclohexanol	120	cyclohexanone	85 (81) ^c
9 2-Methylcyclohexanol	120	2-methylcyclohexanone	84 (78) ^c
10 (-)-Menthol	120	menthone	88 (80)

^a Reaction using 1–1.2 equiv. of zeofen under microwave irradiation in a solventless system. The physical data of the isolated products were in agreement with those of authentic samples. ^b Yields are based on GLC analysis. Figures in parentheses are yields of isolated products. ^c Yields based on isolation of the 2,4-DNPH derivative.

perature in solvent and under solvent-free conditions using a house-hold microwave oven. Solvent-free microwave irradiation has an advantage over conventional heating, offering a practical and environmentally benign protocol, decreasing reaction time and in some cases giving cleaner reactions and easier work up.

Research supported by the National Research Council of Iran (NRCI) as a National Research project under grant number 3708.

Notes and references

- 1 I. Delaude and P. Lazlo, *J. Org. Chem.*, 1996, **61**, 6360.
- 2 R. S. Varma, R. K. Saini and R. Dahuja, *Tetrahedron Lett.*, 1997, **38**, 7823.
- 3 G. Sh. Zhang, Cr. Z. Shi, M. F. Chen and K. Cai, *Synth. Commun.*, 1997, **27**, 3691.
- 4 R. S. Varma and R. K. Saini, *Tetrahedron Lett.*, 1998, **39**, 1481.
- 5 *Comprehensive Organic Synthesis (Oxidation)*, ed. B. M. Trost, Pergamon, New York, 1991, vol. 7.

- 6 A. J. Fatiadi, in *Organic Synthesis by Oxidation with Metal Compounds*, ed. W. J. Mijs and C. R. H. I. de Jonge, Plenum, New York, 1986, pp. 119–260.
- 7 J. Einhorn, C. Einhorn, F. Rajajczak and J. L. Piere, *J. Org. Chem.*, 1996, **61**, 7452.
- 8 J. Mazart, A. N. Ait Aijou and S. Ait-Mohand, *Tetrahedron Lett.*, 1994, **39**, 1989.
- 9 M. M. Heravi, D. Ajami and K. Tabar-Hydar, *Monatsh. Chem.*, 1998, **129**, 1305.
- 10 M. M. Heravi, D. Ajami, K. Tabar-Hydar and M. M. Mojtahedi, *J. Chem. Res. (S)*, 1998, 620.
- 11 B. J. Page and G. W. Loar, in *Kirk-Othmer Encyclopedia of Chemical Technology*, 4th edn., Wiley, New York, 1998, vol. 6, pp. 263–311.
- 12 A. H. Reidies, in *Kirk-Othmer Encyclopedia of Chemical Technology*, 3rd edn., Wiley, New York, 1981, vol. 14, pp. 844–895.
- 13 *Chromium in the Natural and Human Environments*, ed. J. O. Nriagn and E. Nieboer, Wiley, New York, 1988.
- 14 *Chemistry of Waste Minimization*, ed. J. H. Clark, Chapman and Hall, London, 1995.
- 15 R. S. Varma and R. Dehija, *Tetrahedron Lett.*, 1997, **38**, 2043.
- 16 A. McKillop and D. W. Young, *Synthesis*, 1979, 401 and 481.
- 17 M. Balogh and P. Lazlo, *Organic Chemistry using Clays*, Springer Verlag, Berlin, 1993.
- 18 M. M. Heravi, M. Tajbakhsh, S. Y. Beheshtiha and H. A. Oskooie, *Indian J. Heterocyclic Chem.*, 1996, **6**, 143; M. M. Heravi, H. A. Oskooie and M. Mafi, *Synth. Commun.*, 1997, **27**, 1725; M. M. Heravi, M. Tajbakhsh and B. Mohajerani, *Synth. Commun.*, 1999, **29**, 135.
- 19 M. M. Heravi, K. Aghapoor, M. A. Nooshabadi and M. M. Mojtahedi, *Monatsh. Chem.*, 1997, **128**, 1143; K. Aghapoor, M. M. Heravi and M. A. Nooshabadi, 1998, **37B**, 84.
- 20 M. M. Heravi, R. Kiakoojori and K. Tabar-Hydar, *J. Chem. Res.*, 1998, 656; M. M. Heravi, D. Ajami and K. Tabar-Hydar, *Synth. Commun.*, 1999, **29**, 163; M. M. Heravi, D. Ajami, M. M. Mojtahedi and M. Ghassemzadeh, *Tetrahedron Lett.*, 1999, **40**, 561.
- 21 HZSM-5 zeolite was prepared by calcination of NH₄ZSM5 zeolite at 500 °C for 8 h. Si:Al = 40:1 and pore diameters are 5.1 × 55 Å. We are grateful to Dr A. R. Garakani for a gift of zeolite.
- 22 A. Cornilis and P. Lazlo, *Synthesis*, 1985, 909.
- 23 R. A. Abramovitch, *Org. Prep. Proced. Int.*, 1991, **23**, 683; S. Cadadick, *Tetrahedron*, 1995, **51**, 10 403; C. R. Strauss and R. W. Trainer, *Aust. J. Chem.*, 1995, **48**, 1665.
- 24 R. Gedye, F. Smith, K. Weslaway, A. Humera, L. Baldisera and L. R. Laberge, *Tetrahedron Lett.*, 1986, **26**, 279.
- 25 R. S. Varma, R. Dahija and R. K. Saini, *Tetrahedron Lett.*, 1997, **38**, 7029 and references from this group cited therein.

Communication 9/000871