

β -Hydroxysulfoxides as chiral cyclic ketone equivalents: enantioselective synthesis of polysubstituted cyclohexanones, cyclohexenones and cyclohexenediones†

M. Carmen Carreño,* Manuel Pérez-González, María Ribagorda, Álvaro Somoza and Antonio Urbano

Departamento de Química Orgánica (C-I), Universidad Autónoma de Madrid, Cantoblanco, 28049-Madrid, Spain. E-mail: carmen.carrenno@uam.es

Received (in Cambridge, UK) 17th September 2002, Accepted 28th October 2002

First published as an Advance Article on the web 18th November 2002

The β -hydroxysulfoxide moiety, after oxidation to sulfone, acts as a masked carbonyl group in a cyclic system, opening an easy access to differently substituted enantiomerically pure cyclic ketones by means of aluminium-mediated conjugate additions, stereoselective reductions and elimination by retrocondensation in basic medium.


The utilization of chiral cyclohexanones and 2-cyclohexenones in asymmetric synthesis continues to attract considerable interest as an important class of starting materials and synthetic intermediates *en route* to a wide range of natural products like cyclitols,¹ carbasugars^{1b,2} or terpenes.³ The stereocontrolled transformation of cyclic ketones is a general way of access to these important compounds. Currently, this is achieved through the conversion of ketones into their chiral acetals⁴ or hydrazones.⁵ Recently, enantiopure α -acetoxy sulfones generated catalytically have been introduced into the synthetic arsenal as chiral aldehyde equivalents.⁶ Although some of these methodologies allow an efficient control of the asymmetric transformations, new chiral carbonyl equivalents are welcome.

In connection with a program devoted to asymmetric synthesis mediated by sulfoxides,⁷ we have recently reported a general synthetic approach to (*S,R*)-4-hydroxy-4-[(*p*-tolylsulfonyl)methyl]-2,5-cyclohexadienones **1** and studied the stereoselectivity of their conjugate additions.⁸ This study revealed that the β -hydroxy sulfoxide exercises a differentiation between the diastereotopic faces of the double bonds directing conjugate additions of organoaluminium reagents from the face containing the OH group in a highly diastereoselective way and with efficient desymmetrization of the cyclohexadienone moiety. Synthetic applications of these results required the elimination of the chiral auxiliary once the *p*-quinol system had been adequately functionalized. We now report the liberation of ketone functionalities through the oxidation of the corresponding β -hydroxy sulfoxides and elimination of the resulting sulfone. This sequence shows that the readily accessible β -hydroxy sulfoxide moiety can be regarded as a chiral ketone equivalent. In spite of the extended use of β -hydroxy sulfoxides in asymmetric synthesis,^{7a} this is the first example where such a fragment is used as a chiral ketone equivalent.†

Derivatives **2a–d**, bearing the enantiopure β -hydroxy sulfoxide moiety, were prepared through the highly stereoselective organoaluminium conjugate addition⁸ on the corresponding enantiopure *p*-quinols **1a–b**, as unique diastereomers in good yields (Scheme 1). Attempts to eliminate methyl *p*-tolyl sulfonide from **2** through a retrocondensation in basic medium were unsuccessful, probably due to the poor quality of the methyl sulfinyl anion as nucleofuge. We then transformed sulfonides **2** into sulfones **3**. Treatment of sulfone **3a** with different bases led to the expected elimination of methyl *p*-tolyl sulfone but with simultaneous aromatization to the corresponding 2-methylhydroquinone. Reduction of **3a** with DI-

BALH afforded carbinol (*4R*)-**4** resulting from the exclusive axial hydride attack on the most stable conformation **A** of **3a** fixed by the equatorial (*p*-tolylsulfonyl)methyl substituent.

Attempts to recover the masked carbonyl group from **4** in the presence of different bases were again unsuccessful due to difficulties in the isolation of the final 4-hydroxyenone. Fortunately, the TBDMS protected derivative **5** could be transformed into cyclohexenone **6** in the presence of several bases under different conditions. The best results were obtained with 2 equivalents of the weak base Cs₂CO₃ in CH₃CN at rt (Table 1, entry 1) which gave a clean mixture of cyclohexenone **6** and methyl *p*-tolyl sulfone, from which **6** { $[\alpha]_D^{20} = +67.0$ (c 0.4, acetone)} was isolated with 89% yield in enantiomerically pure form, as determined after desilylation into the corresponding carbinol.§



Scheme 1 Reagents and conditions: i, *m*-CPBA, CH₂Cl₂, rt, 1–2 h, 85–95%; ii, DIBALH, THF, –78 °C, 30 min, 95% from **3a**; iii, TBDMSOTf, 2,6-lutidine, CH₂Cl₂, rt, 2 h, 93%; iv, Cs₂CO₃, CH₃CN, rt; v, L-Selectride, THF, –78 °C, 3 h, 64%; vi, NaBH₄, CeCl₃, MeOH, –78 °C, 3–4 h, 64%.

† Electronic supplementary information (ESI) available: full experimental details and NMR spectra. See <http://www.rsc.org/suppdata/cc/b2/209121f/>

The stereoselective reductions of ketone **3b** on the now most stable conformation **B**, which avoids severe interactions between R¹ = Me and the equatorial (*p*-tolylsulfonyl)methyl substituent present in conformation **A** (Scheme 1), afforded carbinols **7** (L-Selectride, 95% equatorial attack) and **8** (NaBH₄, CeCl₃, 80% axial attack), whose treatment with Cs₂CO₃ (entries 2 and 3, respectively) allowed us to obtain in enantiomerically pure form§ both enantiomers of **9**, {[α]_D²⁰ = +48 (c 2, EtOH) and -44 (c 0.2, EtOH)}, known as Phorenol, a synthetic intermediate used for the enantioselective syntheses of several natural products.¹¹ Till now, these chiral synthons were mainly accessible through enzymatic resolutions.¹²

We then focused our attention on the base-induced elimination of methyl *p*-tolylsulfone on cyclohexenones **3b–d** *en route* to chiral cyclohexenediones (Scheme 1). Stirring sulfone **3b** in CH₃CN solution with an excess of Cs₂CO₃ gave, after flash chromatography, achiral 4-oxoisophorone **10b** (entry 4). The synthesis of enantiomerically pure¶ oxoisophorone analogues **10c**¹³ {[α]_D²⁰ = +0.6 (c 0.6, CHCl₃)} and **10d** {[α]_D²⁰ = -3.0 (c 0.8, CHCl₃)} could be achieved in a similar easy way from sulfones **3c** (entry 5) and **3d** (entry 6), respectively.

In order to know if the driving force for the easy elimination of methyl *p*-tolylsulfone was the formation of the conjugated cyclohexenone or cyclohexenedione fragment, we decided to explore the synthetic sequence *en route* to trisubstituted cyclohexanones **11** and **12** (Scheme 2). The precursors were obtained as follows. The stereoselective addition of Et₃Al on **2a** afforded cyclohexanone **13** as the unique diastereoisomer with the *R* absolute configuration at the new chiral center.⁸ *m*-CPBA oxidation of sulfoxide **13** gave sulfone **14** which was stereoselectively reduced to carbinol **15** (L-Selectride, 90% equatorial attack) or to the epimer **16** (NaBH₄, CeCl₃, 95% axial attack). The elimination of methyl *p*-tolyl sulfone was once again unsuccessful on free carbinols **15** and **16** and had to be carried out on the TBDMS protected derivatives **17** and **18**. Simply stirring a CH₃CN solution of **17** or **18** with Cs₂CO₃ (entries 7 and 8) afforded, respectively, the desired trisub-

stituted cyclohexanones **11** {[α]_D²⁰ = -1.7 (c 1.1, CHCl₃)} and **12** {[α]_D²⁰ = -4.5 (c 1.1, CHCl₃)} in enantiomerically pure form, as determined after desilylation into the corresponding carbinols.§

In summary, the enantiopure β-hydroxysulfoxide moiety can be considered as a chiral cyclic ketone equivalent which can be recovered after oxidation to a sulfone and mild basic treatment. An additional advantage of the β-hydroxysulfone as carbonyl protecting group in cyclic systems is that further transformations on such rigid systems occur in a stereocontrolled manner. Moreover, both the α-hydroxysulfoxides and the corresponding sulfones are very stable and can be easily handled and purified either by chromatography or crystallization. We are now applying this new efficient methodology to the total enantioselective synthesis of several natural products.

We thank DGICYT (Grant PB98-0062) and MCYT (Grant BQU2002-03371) for financial support, MEC for a fellowship to M. R. and CAM for a fellowship to A. S.

Notes and references


‡ β-Hydroxy sulfoximines have been used to resolve chiral racemic cyclic ketones.⁹

§ The optical purity was determined from the Mosher's esters.¹⁰

¶ The optical purity for **10c,d** was determined after reduction at C-4 using NaBH₄-CeCl₃ and formation of the Mosher's esters.¹⁰

Table 1 Treatment of cyclic β-hydroxysulfones with Cs₂CO₃ in CH₃CN at rt

Entry	β-Hydroxy sulfone	Cs ₂ CO ₃ equiv.	t/h	Cyclic ketone	Yield (%)	Ee (%)
1	5	2	17	(4 <i>R</i> ,6 <i>S</i>)- 6	89	>95
2	7	3	1	(4 <i>R</i>)- 9	90	>95
3	8	3	0.25	(4 <i>S</i>)- 9	93	>95
4	3b	2	0.3	10b	54	—
5	3c	2	0.5	(6 <i>S</i>)- 10c	70	>95
6	3d	2	0.25	(6 <i>R</i>)- 10d	68	>95
7	17	2	20	(2 <i>R</i> ,4 <i>S</i> ,6 <i>S</i>)- 11	89	>95
8	18	2	6	(2 <i>R</i> ,4 <i>R</i> ,6 <i>S</i>)- 12	82	>95



Scheme 2 Reagents and conditions: i, *m*-CPBA, CH₂Cl₂, rt, 1–2 h, 98%; ii, L-Selectride, THF, -78 °C, 2 h, 73%; iii, NaBH₄, CeCl₃, MeOH, -78 °C, 3–4 h, 75%; iv, TBDMSOTf, 2,6-lutidine, CH₂Cl₂, 0 °C, 2 h, 81–87%; v, Cs₂CO₃, CH₃CN, rt.

- Reviews: T. Hudlicky and M. Cebulak, *Cyclitols and Their Derivatives. A Handbook of Physical, Spectral, and Synthetic Data*, VCH, New York, 1993; L. Agrofoglio, E. Suhas, A. Farese, R. Condom, S. R. Challand, R. A. Earl and R. Guedj, *Tetrahedron*, 1994, **50**, 10611; T. Hudlicky, D. A. Entwistle, K. Pitzer and A. J. Thorpe, *Chem. Rev.*, 1996, **96**, 1195; Recent reference: S. E. De Sousa, P. O'Brien and C. D. Pilgram, *Tetrahedron Lett.*, 2001, **42**, 8081.
- Reviews: M. T. Crimmins, *Tetrahedron*, 1998, **54**, 9229; A. Berecibar, C. Grandjean and A. Siriwardena, *Chem. Rev.*, 1999, **99**, 779.
- Recent reference: G. P.-J. Hareau, M. Koiwa, S. Hikichi and F. Sato, *J. Am. Chem. Soc.*, 1999, **121**, 3640 and references therein.
- Reviews: A. Alexakis and P. Mangeney, *Tetrahedron: Asymmetry*, 1990, **1**, 477; B. C. Rossiter and N. M. Swingle, *Chem. Rev.*, 1992, **92**, 771; Recent references: F. Busqué, P. de March, M. Figueredo, J. Font and S. Rodríguez, *Tetrahedron: Asymmetry*, 2001, **12**, 3077; D. F. Taber, T. E. Christos, M. Rahimizadeh and B. Chen, *J. Org. Chem.*, 2001, **66**, 5911.
- Review: A. Job, C. F. Janeck, W. Bettray, R. Peters and D. Enders, *Tetrahedron*, 2002, **58**, 2253.
- B. M. Trost, M. L. Crawley and C. Bom Lee, *J. Am. Chem. Soc.*, 2000, **122**, 6120.
- Overview: M. C. Carreño, *Chem. Rev.*, 1995, **95**, 1717; Recent work: M. C. Carreño, A. Urbano and C. Di Vitta, *Chem. Eur. J.*, 2000, **6**, 906; M. C. Carreño, S. García-Cerrada, M. J. Sanz-Cuesta and A. Urbano, *Chem. Commun.*, 2001, 1452; M. C. Carreño, S. García-Cerrada and A. Urbano, *J. Am. Chem. Soc.*, 2001, **123**, 7929; M. C. Carreño, C. García Luzón and M. Ribagorda, *Chem. Eur. J.*, 2002, **8**, 208; M. C. Carreño, S. García-Cerrada and A. Urbano, *Chem. Commun.*, 2002, 1412; M. C. Carreño, M. Ribagorda and G. H. Posner, *Angew. Chem., Int. Ed.*, 2002, **41**, 2753; M. C. Carreño, M. Ribagorda, A. Somoza and A. Urbano, *Angew. Chem., Int. Ed.*, 2002, **41**, 2755.
- M. C. Carreño, M. Pérez González, M. Ribagorda and K. N. Houk, *J. Org. Chem.*, 1998, **63**, 3687.
- C. R. Johnson and J. R. Zeller, *J. Am. Chem. Soc.*, 1982, **104**, 4021.
- J. A. Dale and H. S. Mosher, *J. Am. Chem. Soc.*, 1973, **95**, 512.
- Carotenoids: H. Mayer and A. Ruettimann, *Helv. Chim. Acta*, 1980, **63**, 1451; R. Zell, E. Widmer, T. Lukac, H. G. W. Leuenberger, P. Schoenholzer and E. A. Broger, *Helv. Chim. Acta*, 1981, **64**, 2447; J. A. Haugan, E. Lobkovsky and S. Liaaen-Jensen, *Acta Chem. Scand.*, 1997, **51**, 1201; Abscisic acid: M. Soukup, T. Lukac, B. Lohri and E. Widmer, *Helv. Chim. Acta*, 1989, **72**, 361; α-ionone: H. Pfander and P. A. Semadeni, *Aust. J. Chem.*, 1995, **48**, 145.
- H. Kiyota, M. Nakabayashi and T. Oritani, *Tetrahedron: Asymmetry*, 1999, **10**, 3811 and references therein.
- This derivative has been recently described for the first time, albeit in racemic form: A. Aponick, R. S. Buzdygon, R. J. Tomko, A. N. Fazal, E. L. Shughart, D. M. McMaster, M. C. Myers, W. H. Pitcock and C. T. Wigal, *J. Org. Chem.*, 2002, **67**, 242.