

Catalytic one-pot synthesis of [5.5.5.6]fenestrane systems via a dicobalt octacarbonyl-catalyzed tandem cycloaddition of dienediynes†

Do Han Kim,^a Seung Uk Son,^a Young Keun Chung^{*a} and Sueg-Geun Lee^b

^a School of Chemistry and Center for Molecular Catalysis, Seoul National University, Seoul 151-747, Korea.

E-mail: ykchung@plaza.snu.ac.kr; Fax: (+82)2-889-0310; Tel: +82-2-880-6662

^b Korea Research Institute of Chemical Technology, PO Box 107, Yusong, Taejeon 305-600, Korea

Received (in Cambridge, UK) 5th September 2001, Accepted 20th November 2001

First published as an Advance Article on the web


Catalytic one-pot synthesis of fenestrane derivatives from dienediynes was developed: fenestranes were synthesized in high yields by a dicobalt octacarbonyl-catalyzed tandem cycloaddition of dienediynes.

One of the challenging uses of organometallic chemistry is to find reactions and strategies that allow for the facile conversion of simple compounds into complex materials, medicines, or molecules of theoretical interest.¹ Recently we have demonstrated² that the dicobalt octacarbonyl-catalyzed carbonylative cycloaddition of diene or triyne with/without diene or alkyne can produce up to four new 5-membered rings in one step. This success encouraged us to analyze the retrosynthesis of fenestranes,³ which has been studied in relation to the question of whether a tetrahedral carbon can be flattened.

Most of the previous syntheses of fenestranes are based on the Pauson-Khand reaction,⁴ photocycloaddition,⁵ Pd-catalyzed carbonylative cyclization,⁶ aldolization⁷ or cyclodehydration⁸ as the key step(s). They gave only small quantities of the desired tetracycles and suffered from a low selectivity and tedious separating problems. Now we report a catalytic synthesis of fenestrane derivatives using dicobalt octacarbonyl-catalyzed tandem cycloaddition of dienediynes under CO pressure.

Keese's group reported a one-pot synthesis of [5.5.5.5]fenestranes from enediynes *via* the Pauson-Khand reaction.^{4b} However, the reported yields were low.

Thus, at first we thought that enediynes (**1**) or triynes (**2**) might be appropriate precursors for the construction of a fenestrane




structure through two sequential [2 + 2 + 1] cycloaddition reactions. The inner triple bond of triynes might permit the construction of a quaternary center from two sequential carbon-carbon bond formation steps. Contrary to our expectation, a fenestrane structure was not formed. Instead, in the case of enediynes, polymeric materials were obtained and, in the case of triynes, an unnatural tetracyclic structure was obtained as the sole product.⁹

After many experimental trials, we chose dienediyne as a precursor to fenestrane.

For the cyclization, dienediynes derivatives **1a**, **1b**, and **1c** (Scheme 1) were prepared from the α -hydroxy diynes **2a**, **2b** and **2c**. Compound **2b** was prepared according to the procedures in the literature.¹⁰ The same procedure as used for the synthesis of **2b** has been applied to the synthesis of **2a** and **2c**. The reaction of α -hydroxy diynes with sodium hydride followed by hexa-2,4-dienyl bromide led to the isolation of **1a**, **1b**, and **1c** in high yields which were fully characterized.^{11‡}


Treatment of dienediynes **1a** (0.68 mmol) with dicobalt octacarbonyl (5 mol%) in dichloromethane at 130 °C under 30

atm of CO for 18 h yielded a fenestrane derivative **3a** in 84% yield [eqn. (1)].§




The structure of **3a** was established by ¹H and ¹³C NMR spectroscopic investigations (COSY, long range COSY, DEPT, HECTOR, long range HECTOR, phase sensitive NOESY, and 2D-INADEQUATE), and mass spectrometry.¶ The correlation peaks obtained from the COSY spectrum allow us to identify the consecutive connections of methyl-14 protons to methylene-9 proton, 9 to 10, 10 to 11, 11 to 12 and 12 to 13. The other connections were derived from the combinative information observed from the DEPT, 2D C–H correlation, and 2D long range C–H correlation spectra. No other products were observed. The reaction involves the formation of five carbon-carbon bonds.

Although this multibond formation reaction seems to be complex, it may be simple: the dicobalt octacarbonyl-catalyzed carbonylative [2 + 2 + 1] cycloaddition of diyne with carbon monoxide followed by a subsequent intramolecular [4 + 2] cycloaddition provides **3a**. Another feasible reaction pathway is that the diene unit reacts first with the closer triple bond and the thus formed cyclohexa-1,4-diene undergoes a Pauson-Khand reaction with the remote triple bond.¹² The central triple bond acts as a double functionality that could be used in both cyclizations. In the same way as in the synthesis of **3a**, the dicobalt octacarbonyl-catalyzed tandem cyclization of dien-



Scheme 1

† Electronic supplementary information (ESI) available: characterization of synthesized compounds. See <http://www.rsc.org/suppdata/cc/b1/b108034b/>



diynes **1b** and **1c** afforded **3b** and **3c** in 74% and 51% yields, respectively.

In conclusion, we have developed a new catalytic route to fenestrane derivatives *via* dicobalt octacarbonyl-catalyzed cycloaddition of dienediynes. The following significant points are noteworthy. All the reactions described here are catalytic with high conversion rates and experimentally a simple reaction, a one-pot reaction.

This work was supported by grant No. 2000-2-12200-001-1 from the Basic Research Program of the Korea Science and Engineering Foundation (KOSEF), KOSEF (1999-1-122-001-5), and KOSEF through the Center for Molecular Catalysis. SUS and DHK thank the Ministry of Education for the Brain Korea 21 Fellowship.

Notes and references

‡ Compound **2a** (0.20 g, 0.69 mmol) was dissolved in 15 ml of THF. The solution was cooled to 0 °C. To the solution was added NaH (40 mg, 50 wt% in oil). After the solution was stirred for 1 h, hexa-2,4-dienyl bromide (0.17 g, 1.05 mmol) was added to the solution. The resulting solution was stirred for 12 h and was quenched with diethyl ether and sat. NH₄Cl solution. The ether layer was separated, dried over anhydrous MgSO₄, concentrated and the residue separated on a silica gel eluting with hexane and diethyl ether (v/v, 10:1). Yield: 0.23 g (89%). **1a**: ¹H NMR (CDCl₃, 300 MHz): δ 7.39 (m, 4 H), 7.28 (m, 6 H), 6.27 (dd, 15.0, 10.0 Hz, 1 H), 6.00 (dd, 15.0, 10.0 Hz, 1 H), 5.70 (m, 2 H), 4.37 (dd, 13.0, 5.4 Hz, 1 H), 4.26 (s, 1 H), 4.08 (dd, 13.0, 7.2 Hz, 1 H), 2.65 (d, 17.0 Hz, 1 H), 2.47 (d, 17.0 Hz, 1 H), 1.71 (d, 6.2 Hz, 3 H), 1.20 (s, 3 H), 1.19 (s, 3 H) ppm; ¹³C NMR (CDCl₃, 75 MHz): δ 133.6, 131.7, 131.6, 130.9, 129.8, 128.2, 128.1, 127.5, 126.7, 124.1, 122.9, 87.9, 86.9, 82.6, 75.4, 69.7, 39.1, 29.4, 23.7, 22.8, 18.0 ppm; exact mass calc. 368.2140, obsd. 368.2137.

§ Compound **1a** (0.25 g, 0.68 mmol), 15 ml of CH₂Cl₂, and Co₂(CO)₈ (12 mg, 0.035 mmol) were put in a high pressure reactor (100 ml). After the solution was bubbled with nitrogen for 1 min, the reactor was pressurized with 30 atm of CO. The reactor was heated at 130 °C for 18 h. After the reactor was cooled to rt, excess gas was released and the reaction mixture was transferred into a one-neck flask (50 ml). Removal of the solvent followed by chromatography on a silica gel column eluting with hexane and diethyl ether (v/v, 5:1) gave **3a** in 84% yield (0.23 g, 0.58 mmol).

¶ ¹H NMR (CDCl₃, 300 MHz): δ 7.60–7.24(m, 10 H), 6.04 (m, 1 H), 5.73 (m, 1 H), 4.16 (dd, 8.7, 5.5 Hz, 1 H), 3.84 (dd, 8.7, 5.5 Hz, 1 H), 3.70 (s, 1 H), 2.90 (m, 1 H), 2.72 (m, 1 H), 2.68 (d, 15.0 Hz, 1 H), 2.43 (d, 15.0 Hz, 1 H), 1.22 (d, 7.3 Hz, 3 H), 1.04 (s, 3 H), 0.00 (s, 3 H) ppm; ¹³C NMR (CDCl₃, 75 MHz): δ 209.2, 180.7, 139.2, 137.9, 134.8, 131.3, 129.8, 128.9, 128.3, 127.7, 126.6, 89.6, 75.3, 70.3, 65.9, 48.0, 43.9, 42.7, 36.8, 25.9, 23.3,

18.9 ppm; IR ν_{CO} 1693 cm⁻¹; exact mass calc. 396.2089, obsd. 396.2088.

- (a) O. G. Kulinkovich and A. de Meijere, *Chem. Rev.*, 2000, **100**, 2789; (b) F. Sato, H. Urabe and S. Okamoto, *Chem. Rev.*, 2000, **100**, 2835; (c) J. Tsuji, *Transition Metal Reagents and Catalysts*, Wiley & Sons, Sussex, 2000; (d) *Transition Metals for Organic Synthesis*, ed. M. Beller and C. Bolm, Wiley-VCH, Weinheim, 1998; (e) L. S. Hegedus, *Transition Metals in the Synthesis of Complex Organic Molecules*, University Science Book, Mill Valley, CA, 1994.
- (a) S. H. Hong, D. S. Choi, Y. K. Chung and S. Lee, *Chem. Commun.*, 1999, 2099; (b) S. U. Son, S.-J. Paik, S. I. Lee and Y. K. Chung, *J. Chem. Soc., Perkin Trans. 1*, 2000, 141; (c) S. U. Son, D. S. Choi and Y. K. Chung, *Org. Lett.*, 2000, **2**, 2097; (d) S. U. Son, Y. K. Chung and S. Lee, *J. Org. Chem.*, 2000, **65**, 6142.
- Reviews on fenestrane chemistry: (a) D. Kuck, in *Advances in Theoretically Interesting Molecules*, ed. R. P. Thummel, JAI Press, Greenwich, CT, 1998, vol. 4, p. 81; (b) M. Thommen and R. Keese, *Synlett*, 1997, 231; (c) W. Luef and R. Keese, in *Advances in Strain in Organic Chemistry*, ed. B. Halton, JAI Press, Greenwich, CT, 1993, vol. 3, p. 229; (d) W. C. Agosta, in *The Chemistry of Alkanes and Cycloalkanes*, ed. S. Patai and Z. Rappoport, Wiley, New York, 1992, p. 927; (e) A. K. Gupta, X. Fu, J. P. Snyder and J. M. Cook, *Tetrahedron*, 1991, **47**, 3665; (f) K. Krohn, in *Organic Synthesis Highlights*, ed. J. Mulzer, H.-J. Altenbach, M. Braun, K. Krohn and H.-U. Reissig, VCH, Weinheim, 1991, p. 121; (g) B. R. Venepalli and W. C. Agosta, *Chem. Rev.*, 1987, **87**, 399; (h) R. Keese, in *Organic Synthesis; Modern Trends*, ed. O. Chizhov, Blackwell, Oxford, 1987, p. 43.
- (a) W. A. Smit, S. M. Buhanjuk, S. O. Simonyan, A. S. Shashkov, Y. T. Struchkov, A. I. Yanovsky, R. Caple, A. S. Gybin, A. G. Anderson and J. A. Whiteford, *Tetrahedron Lett.*, 1991, **32**, 2105; (b) M. Thommen, P. Gerber and R. Keese, *Chimia*, 1991, 21; (c) A. van der Waals and R. Keese, *J. Chem. Soc., Chem. Commun.*, 1992, 570; (d) M. Thommen, A. Veretenov, R. Guidetti-Grept and R. Keese, *Helv. Chim. Acta*, 1996, **79**, 461; (e) M. M. Bruendddl, S. G. Van Ornum, T.-M. Chan and J. M. Cook, *Tetrahedron Lett.*, 1999, **40**, 1113; (f) S. G. Van Ornummm, M. M. Bruendl, H. Cao, M. Reddy, D. S. Grubisha, D. W. Bennett and J. M. Cook, *J. Org. Chem.*, 2000, **65**, 1957; (g) H. Caa, S. G. Van Ornum and J. M. Cook, *Tetraahedron Lett.*, 2000, **41**, 5313.
- (a) W. G. Dauben and D. M. Walker, *Tetrahedron Lett.*, 1982, **23**, 711; (b) J. Mani and R. Keese, *Tetrahedron*, 1985, **41**, 5697; (c) M. T. Crimmins, S. W. Mascarella and L. D. Bredon, *Tetrahedron Lett.*, 1985, **26**, 997; (d) V. B. Rao, S. Wolff and W. C. Agosta, *Tetrahedron*, 1986, **42**, 1549; (e) P. Wender, T. M. Dore and M. A. deLong, *Tetrahedron Lett.*, 1996, **37**, 7687.
- R. Keese, R. Guidetti-Grept and B. Herzog, *Tetrahedron Lett.*, 1992, **33**, 1207.
- (a) X. Fu, G. Kubiak, W. Zhang, W. Han, A. K. Gupta and J. M. Cook, *Tetrahedron*, 1993, **49**, 1511; (b) O. Kubiak, X. Fu, A. K. Gupta and J. M. Cook, *Tetrahedron Lett.*, 1990, **31**, 4285.
- B. Bredenkötter, U. Hörke and D. Kuck, *Chem. Eur. J.*, 2001, **7**, 3387.
- S. U. Son, Y. A. Yoon, D. S. Choi, J. K. Park and Y. K. Chung, *Org. Lett.*, 2001, **3**, 1065.
- A. J. Pearson, R. J. Shively Jr. and R. A. Dubbert, *Organometallics*, 1992, **11**, 4096.
- See the Supporting Information.†
- A referee suggested this reaction pathway. The authors thank the referee for this suggestion.