Isomenthyl-derived 1,3-diol ligands catalyzing an asymmetric aldol reaction identified through parallel synthesis and screening

John M. Gardiner,*^{*a*} Philip D. Crewe,^{*a*} Gillian E. Smith^{*b*} and Kenneth Veale^{*b*}

^a Department of Chemistry, UMIST, Sackville Street, Manchester M60 1QD. Fax: 001 510 217 2371; Tel: +44 (0)161 200 4530; E-mail: john.gardiner@umist.ac.uk ^b GlaxoSmithKline, New Frontiers Science Park, Harlow, Essex CM19 5AW. Fax: +44 01279 622 348; Tel: +44 01279 622 061; E-mail: gillian e smith@gsk.com

Contents:

HPLC Data for Mukaiyama aldol products from parallel catalyst screen. Ligand generated in situ in each reaction tube is shown.

NMR (¹H, ¹³C, HMBC) for pure reference sample of Mukaiyama aldol product prepared using achiral conditions. This sample confirmed peaks by HPLC to represent enantiomers.

NMR data for independently prepared examples of 2 isomenthone aldol constituents included in the parallel synthesis sequence, and of 2 independently prepared examples of the diols included of the ligand array.











Entry 3





Entry 6







Entry 8





Entry 10



Run9





Run12











 Chronic Dates Transmission
 Chronic Dates Transmission

 EVEND
 1

 FTTADE
 2000 - 200 -



NMR Data for independently prepared isomenthone aldol products and diols



data for single diastereomer: configuration unassigned (reaction generates 80:20)



 Control (Data)
 Control (Data)

 Contro (Data)





Ph

 Nume
 Ourrent:
 Data Resources

 Corrent:
 Data Resources

 <t











S R S OH OH



