

Snapshots of an oxidatively induced α -hydrogen abstraction reaction to prepare a terminal and four-coordinate titanium imide†

Falguni Basuli, Brad C. Bailey, John C. Huffman and Daniel J. Mindiola*

Department of Chemistry and Molecular Structure Center, Indiana University, Bloomington, Indiana 47405, USA. E-mail: mindiola@indiana.edu; Fax: 001 812 855 8300; Tel: 001 812 855 2399


Received (in Purdue, IN, USA) 26th April 2003, Accepted 13th May 2003

First published as an Advance Article on the web 2nd June 2003

One electron oxidation of the bis-anilido titanium(III) complex $(\text{Nacnac})\text{Ti}(\text{NHAr})_2$ ($\text{Nacnac}^- = \text{ArNC}(\text{CH}_3)\text{CHC}(\text{CH}_3)\text{NAr}$, $\text{Ar} = 2,6\text{-}(\text{CHMe}_2)_2\text{C}_6\text{H}_3$) with AgOTf affords the cation $[(\text{Nacnac})\text{Ti}(\text{NHAr})_2][\text{OTf}]$ which is isolated and shown to gradually transform, by α -hydrogen abstraction, to the terminal and four-coordinate titanium imide $(\text{Nacnac})\text{Ti}=\text{NAr}(\text{OTf})$.

Low-coordinate transition metal imido complexes have experienced a remarkable growth in the last 30 years and constitute a landmark in inorganic chemistry.¹ Wolczanski² and Bergman³ have prepared perhaps the most reactive imido complexes of the early-transition metals. This inherent reactivity can lead to the activation of relatively inert bonds such as primary and secondary^{2,3} C–H bonds in alkanes and can be tailored to the unsaturated or electrophilic nature of the metal center. Parallel to the reactivity of early-transition metal imido systems, electron-rich late-transition metal imido complexes can also engage in C–H activation processes,⁴ hence research in this area continues to draw great attention. Our recent report of the synthesis of a four-coordinate titanium alkylidene complex⁵ and tertiary C–H bond activation reactions stemming from this complex⁶ stimulated the pursuit of an analogous titanium imide complex. Inspired by the work of Wolczanski we attempted to prepare low-coordinate titanium imido complexes by an oxidatively induced α -abstraction reaction. Herein we report the conversion of a Ti(III) bis-anilido complex $(\text{Nacnac})\text{Ti}(\text{NHAr})_2$, ($\text{Nacnac}^- = \text{ArNC}(\text{CH}_3)\text{CHC}(\text{CH}_3)\text{NAr}$, $\text{Ar} = 2,6\text{-}(\text{CHMe}_2)_2\text{C}_6\text{H}_3$),⁷ to the corresponding cation $[(\text{Nacnac})\text{Ti}(\text{NHAr})_2]^+$ and subsequent α -hydrogen abstraction to form a stable and terminal four-coordinate titanium imide complex.

Our approach to preparing a low-coordinate and terminal titanium imido complex involved a similar strategy for the synthesis of the analogous alkylidene complex $(\text{Nacnac})\text{Ti}=\text{CH}^t\text{Bu}(\text{OTf})$ (Scheme 1).⁵ Using Budzelaar's dichloride precursor $(\text{Nacnac})\text{TiCl}_2(\text{THF})$ ^{5,8} and 2 equiv of LiNHAr we prepared the bis-anilido titanium(III) complex $(\text{Nacnac})\text{Ti}(\text{NHAr})_2$ (**1**) in 92% yield as dark green blocks (Scheme 2).† In the absence of moisture and oxygen complex **1** is indefinitely stable in the solid state or in solution. X-band solution EPR spectra and magnetic measurements of **1** are consistent with a d^1 paramagnetic species.† The molecular structure of **1**† displays a four-coordinate Ti(III) complex in a tetrahedral environment. Salient features for the structure of **1** include titanium anilido distances of $\text{Ti}(1)\text{--N}(34)$, 1.961(2) Å and $\text{Ti}(1)\text{--N}(47)$, 1.966(2) Å.




Scheme 1 Synthesis of a four-coordinate titanium alkylidene via an oxidatively induced α -hydrogen abstraction reaction.

† Electronic supplementary information (ESI) available: complete experimental, spectroscopic, analytical, and crystallographic details for complexes **1–4**. See <http://www.rsc.org/suppdata/cc/b3/b304633h/>

A cyclic voltammogram of a solution of **1** (THF–TBAH) indicated a reversible oxidation wave at -0.89 V (referenced vs. $\text{FeCp}_2/\text{FeCp}_2^+$) for the Ti(III)/Ti(IV) couple.† Chemically, it was determined that treatment of a thawing THF solution of **1** with AgOTf caused a color change from green to orange–red concomitant with formation of an Ag^0 mirror. Immediate workup after 5 minutes led to isolation of the bis-anilido triflate salt $[(\text{Nacnac})\text{Ti}(\text{NHAr})_2][\text{OTf}]$ (**2**) as dark-red crystals in 71% yield (Scheme 2).† Complex **2** decomposes gradually in solution, and the yield of this complex is highly dependent on both time and temperature. Complex **2** was characterized by ^1H , ^{19}F , ^{13}C and IR spectroscopies.† Diagnostic features for **2** include two NH resonances centered at 11.1 and 10.4 ppm and one methyl environment for the β -carbon of the Nacnac backbone consistent with the molecule retaining C_s symmetry in solution. Complex **2** is insoluble and stable in most common organic solvents such as hexane, C_6H_6 and Et_2O , but soluble in THF and CH_2Cl_2 upon which it converts to a new complex and free aniline (*vide infra*). The insolubility of **2** in organic solvents suggests this complex to be a discrete salt where the OTf anion interacts weakly with the metal center.

Realizing the importance of the triflate anion in the stability of **2** we prepared a more stable salt of **2** using the weakly coordinating $\text{B}(\text{C}_6\text{F}_5)_4$ anion. Accordingly, oxidation of **1** with $[\text{FeCp}^*_2][\text{B}(\text{C}_6\text{F}_5)_4]$ ⁹ in Et_2O at -35 °C affords in 92% isolated yield the bis-anilido salt $[(\text{Nacnac})\text{Ti}(\text{NHAr})_2][\text{B}(\text{C}_6\text{F}_5)_4]$ (**3**) as brick-red microcrystals (Scheme 2).† Complex **3** shows nearly identical spectroscopic features to **2** in solution (δ for NH protons are 11.4 and 10.9 in the ^1H NMR spectrum of **3**).† The molecular structure of **3** was determined by single crystal X-ray diffraction studies and exhibits a four-coordinate titanium cation confined in a tetrahedral environment (Fig. 1).‡ Important crystallographic features for **3** include short Ti– $\text{N}_{\text{anilido}}$ bonds lengths of $\text{Ti}(1)\text{--N}(33)$, 1.863(2) Å and $\text{Ti}(1)\text{--N}(46)$, 1.875(2) Å, which is consistent with metal–ligand multiple bond character.¹ Most notably, the Ti– $\text{N}_{\text{anilido}}$ bond length is considerably shorter when compared to the neutral and d^1 complex **1** (*vide supra*). Other important crystallographic features for **3** include significantly different $\text{Ti}(1)\text{--N}(33)\text{--C}(34)$ and $\text{Ti}(1)\text{--N}(46)\text{--C}(47)$ angles of $132.8(9)^\circ$ and $157.5(9)^\circ$, respectively. The α -hydrogens on the anilido nitrogens were located in the E-map and are indicative of agostic interactions with the Ti(IV) center.†



Scheme 2 Synthesis of complexes **1–4** starting from Budzelaar's precursor $(\text{Nacnac})\text{TiCl}_2(\text{THF})$.




Fig. 1 Molecular structure of **3** (cation only) and **4** showing atom-labeling scheme with thermal ellipsoids at the 50% probability level. H-atoms and aryl groups with the exception of *ipso*-carbons on the nitrogen atoms have been omitted for clarity. §

Intrigued by the instability of **2** we reasoned that decomposition of this complex lead to α -hydrogen abstraction concomitant with formation of a strong Ti=N bond and free aniline. In fact, if the reaction of **1** with AgOTf is allowed to proceed for 1 hour or longer, subsequent work-up of the mixture affords the titanium imido complex (Nacnac)Ti=NAr(OTf) (**4**) as red prisms in 67% yield (Scheme 2).[†] Hence, complex **2** is an intermediate to **4** inasmuch as stirring solutions of isolated **2** at room temperature in CH₂Cl₂ afford **4** and free aniline in quantitative yield.[†] The role of the anion is important in the stability of the cation since treatment of **3** with Ti(OTf)¹⁰ in CH₂Cl₂ also promotes α -hydrogen abstraction to give **4** as evidenced by the ¹H NMR spectrum of the reaction mixture (Scheme 2).[†] The choice of solvent also plays an important role in the α -abstraction process since THF appears to accelerate the formation of **4**, relative to CH₂Cl₂. Single crystals of **4** were grown from pentane at -35 °C and the molecular structure is depicted in Fig. 1.¶ The structure of complex **4** shows a rare example of a four-coordinate titanium imido^{2,5} complex with a short Ti(1)–N(33) bond length of 1.705(5) Å and a nearly linear Ti–N_{imido}–C_{ipso} linkage (170.5(1)°). ¹H and ¹³C NMR spectra are in accordance with **4** retaining C_s symmetry in solution, which is also consistent with the molecular structure. Complex **4** is isostructural to the reported alkylidene derivative (Nacnac)Ti=CH^tBu(OTf).⁶

Our results suggest that both an electron deficient metal center in addition to coordination of a fifth ligand (–OTf) promotes α -hydrogen abstraction. Coordination of –OTf likely induces α -hydrogen abstraction by steric crowding of the anilido ligands. Alternatively, Lewis bases such as THF appear to enhance hydrogen abstraction, likely participating as proton carriers. Schrock and co-workers have observed closely related α -hydrogen abstraction reactions stemming from 5-coordinate d⁰ molybdenum species.¹¹ In contrast to alkyl groups on titanium,⁵ the anilido lone pair of electrons greatly increases the lifetime of the intermediate. This allows us to acquire a snapshot of long-lived intermediates associated with the oxidatively induced α -hydrogen abstraction process. The present work defines the steps by which one electron oxidation can lead to α -

hydrogen abstraction and formation of a low-coordinate titanium-imido complex.

For financial support of this research we thank Indiana University-Bloomington, the Camille and Henry Dreyfus Foundation, and the Ford Foundation. D.J.M. would like to thank Mr. X. Hu, Dr. R. Isaacson, and Professors K. Meyer and K. G. Caulton for insightful discussions.

Notes and references

‡ Crystal data for **3**·Et₂O, C₈₁H₈₇BF₂₀N₄OTi: Monoclinic, P2₁/c, *a* = 13.8639(6), *b* = 36.266(6), *c* = 13.9675(6) Å, β = 93.2700(10)°, *Z* = 4, μ (Mo–K α) = 0.210 mm^{–1}, *V* = 7591.3(6) Å³, *D_c* = 1.375 mg mm^{–3}, GoF on *F*² = 0.962, *R*(*F*) = 4.43% and *R*(*wF*) = 11.16%. Out of a total of 111677 reflections collected 17467 were independent and 9392 were observed (*R*_{int} = 10.95 %) with *I* > 2 σ (orange prism, 0.25 × 0.25 × 0.07 mm, 27.55° \geq θ \geq 2.07°).

§ CCDC 209461–209463. See <http://www.rsc.org/suppdata/cc/b3/b304633h/> for crystallographic data in .cif or other electronic format.

¶ Crystal data for **4**·pentane, C₄₇H₇₀F₃N₃O₃STi: Triclinic, P $\bar{1}$, *a* = 10.271(4), *b* = 12.965(7), *c* = 18.451(3) Å, α = 93.783(7)°, β = 106.097(9)°, γ = 96.880°, *Z* = 2, μ (Mo–K α) = 0.282 mm^{–1}, *V* = 2330.9(6) Å³, *D_c* = 1.228 mg mm^{–3}, GoF on *F*² = 0.933, *R*(*F*) = 4.00% and *R*(*wF*) = 9.78%. Out of a total of 51577 reflections collected 10763 were independent and 7221 were observed (*R*_{int} = 8.34 %) with *I* > 2 σ (orange prism, 0.30 × 0.25 × 0.25 mm, 27.60° \geq θ \geq 2.07°). The crystal data shows a pseudo-inversion center and was merohedrally twinned (domain ratio 57 : 43).

- D. E. Wigley, *Prog. Inorg. Chem.*, 1994, **113**, 2985; W. A. Nugent and J. M. Mayer, *Metal-Ligand Multiple Bonds*, John Wiley & Sons, New York, 1988.
- C. C. Cummins, S. M. Baxter and P. T. Wolczanski, *J. Am. Chem. Soc.*, 1988, **110**, 8731; C. C. Cummins, C. P. Schaller, G. D. Van Duyne, P. T. Wolczanski, A. W. Chan and R. Hoffmann, *J. Am. Chem. Soc.*, 1991, **113**, 2985; C. P. Schaller and P. T. Wolczanski, *Inorg. Chem.*, 1993, **32**, 131; J. L. Bennett and P. T. Wolczanski, *J. Am. Chem. Soc.*, 1994, **116**, 2179; C. P. Schaller, C. C. Cummins and P. T. Wolczanski, *J. Am. Chem. Soc.*, 1996, **118**, 591; J. L. Bennett and P. T. Wolczanski, *J. Am. Chem. Soc.*, 1997, **119**, 10696; D. F. Schafer and P. T. Wolczanski, *J. Am. Chem. Soc.*, 1998, **120**, 4881.
- P. J. Walsh, F. J. Hollander and R. G. Bergman, *J. Am. Chem. Soc.*, 1988, **110**, 8729.
- S. Thyagarajan, D. T. Shay, C. D. Incarvito, A. L. Rheingold and K. H. Theopold, *J. Am. Chem. Soc.*, 2003, **125**, 4440.
- F. Basuli, B. C. Bailey, J. Tomaszewski, J. C. Huffman and D. J. Mindiola, *J. Am. Chem. Soc.*, 2003, **125**, 6052.
- F. Basuli, B. C. Bailey, L. A. Watson, J. C. Huffman and D. J. Mindiola, manuscript in preparation.
- M. Stender, R. J. Wright, B. E. Eichler, J. Prust, M. M. Olmstead, H. W. Roesky and P. P. Power, *J. Chem. Soc., Dalton Trans.*, 2001, 3465.
- P. H. M. Budzelaar, A. B. von Oort and A. G. Orpen, *Eur. J. Inorg. Chem.*, 1998, 1485.
- D. J. Mindiola, K. Kitiachvili and G. L. Hillhouse, unpublished results. For synthesis see ESI.
- M. E. Woodhouse, F. D. Lewis and T. J. Marks, *J. Am. Chem. Soc.*, 1982, **104**, 5586.
- R. R. Schrock, *Chem. Rev.*, 2002, **102**, 145; R. Baumann, R. Stumpf, W. M. Davis, L.-C. Liang and R. R. Schrock, *J. Am. Chem. Soc.*, 1999, **121**, 7823; R. R. Schrock, J. S. Murdzek, G. C. Bazan, J. Robbins, M. DiMare and M. O'Reagan, *J. Am. Chem. Soc.*, 1990, **112**, 3875.