

New aerobic oxidation of benzylic compounds: efficient catalysis by *N*-hydroxy-3,4,5,6-tetraphenylphthalimide (NHTPPI)/CuCl under mild conditions and low catalyst loading†

Malek Nechab, Cathy Einhorn and Jacques Einhorn*

Laboratoire d'Etudes Dynamiques et Structurales de la Sélectivité/UMR 5616, Université Joseph Fourier, BP 53, 38041 Grenoble, Cedex 9, France. E-mail: Jacques.Einhorn@ujf-grenoble.fr; Fax: (+33) 4 76 51 48 36; Tel: (+33) 4 76 51 48 39

Received (in Cambridge, UK) 27th February 2004, Accepted 28th April 2004

First published as an Advance Article on the web 28th May 2004


Efficient aerobic oxidation of benzylic compounds using NHTPPI, a new NHPI analogue, as a key catalyst combined with CuCl, have been achieved under mild conditions and using as little as 1 mol% catalyst.

In recent years *N*-hydroxyphthalimide (NHPI, **1a**) has been recognized as a valuable catalyst for the aerobic oxidation of various organic compounds under mild conditions.¹ These oxidations proceed *via* intermediate phthalimide *N*-oxyl (PINO, **1b**) radical, able to abstract a hydrogen atom from the organic substrate.² The newly formed carbon centered radical then readily reacts with dioxygen, to give ultimately oxygenated compounds. PINO, however, is not indefinitely stable: its self-decomposition obeys second-order kinetics and has been studied by Masui³ and, more recently, by Espenson.^{2c,e} Self-decomposition of PINO makes it necessary to use quite large amounts of NHPI (generally 10 mol%) to achieved good conversions in the oxidation processes. Recently, various NHPI analogues have been prepared with the aim of tuning the catalyst performances,^{2e,4} or of developing asymmetric versions.⁵ These analogues, however, have mostly been used at catalyst loading ranging from 5 to 10 mol%.

Herein, we report a new development in NHPI type catalysis, allowing highly efficient benzylic oxidation at mild temperature, with low catalyst loading. In a first series of experiments, we have investigated the aerobic oxidation of indan, chosen as a model substrate, under various experimental conditions but limiting the amount of NHPI to 1 mol%. CuCl†⁶ has been found to be, by far, the most active co-catalyst among a large variety of metal salts tested (Co(acac)₂, Mn(acac)₂, Mn(acac)₃, V(acac)₂, V(acac)₃, Mn(OAc)₃, CoCl₂, MnCl₂, CuCl, CuCl₂): 62% of indan-1-one was obtained in acetonitrile at 35 °C after 6 h, using 5 mol% of CuCl. We have next replaced **1a** by several analogues (Table 1).§

Compared to **1a** (run 1), analogues bearing electron donating or electron withdrawing substituents (**2–5**)^{4b,7} were less effective (runs 2–5). Lower effectiveness was also observed for tetrachloro- and naphthalenic analogues **6** and **7**⁷ (runs 6 and 7). Our previously

described chiral *N*-hydroxyimides **8** and **9**,⁵ both gave poorer results than **1a** (runs 8 and 9). Finally, *N*-hydroxy-3,4,5,6-tetraphenylphthalimide (**10a**, NHTPPI) revealed a very good catalytic activity (run 10), yielding 80% of indan-1-one after 6 h.



To our knowledge NHTPPI has not been described previously. It can be prepared in a single step from commercially available tetraphenylphthalic anhydride (ESI†). It is more soluble than NHPI in most of the classical organic solvents. The main features of the NHTPPI/CuCl catalysed oxidation system are as follows: acetonitrile is the solvent of choice, benzonitrile, trifluoromethylbenzene, acetic acid or ethyl acetate being less appropriate (probably due to a lack of solubility of CuCl). The optimum amount of CuCl has been found to be around 5 mol%. Larger amounts resulted in no improvement of the reaction rate or of the final yield of indan-1-one. Smaller amounts reduce the reaction rate: if the quantity of CuCl was diminished to 0.5 mol%, 45% indan-1-one was formed after 6 h but the reaction then became exceedingly sluggish. The effect of temperature on the oxidation of indan using 1 mol% NHTPPI and 5 mol% CuCl is outlined in Fig. 1.

At 20 °C, 24 h are necessary to get 78% yield of indan-1-one, with a very slow further evolution. At 35 °C a 80% yield was obtained after only 6 h with no further evolution. At 50 °C the

Table 1 Aerobic oxidation of indan using various NHPI analogues (1 mol%) as catalysts in acetonitrile at 35 °C

Run	Catalyst	Yield ^a (%)
1	1a	62
2	2	45
3	3	46
4	4	30
5	5	11
6	6	34
7	7	48
8	8	38
9	9	15
10	10a	80

^a Determined by GC after 6h

† Electronic supplementary information (ESI) available: experimental details. See <http://www.rsc.org/suppdata/cc/b4/b403004d/>




Fig. 1 Oxidation of indan catalysed by NHTPPI (1 mol%)/CuCl (5 mol%) at various temperatures.

reaction stopped at 55% yield in less than 6 h. At 80 °C it stopped at 33% yield in less than 1 h. These results clearly indicate that catalysis by NHTPPI is more rapid at higher temperature, but that better turnover numbers of the catalyst are obtained at lower temperature.

A series of benzylic compounds has next been oxidized using 1 or 2 mol% NHTPPI and 5 mol% CuCl (Table 2). Indan-1-one or 2-methylindan-1-one have been obtained in 77 and 65% isolated yields, which represent more than 90% yields based on indan or 2-methylindan reacted (runs 1 and 2). Tetralin was similarly oxidized to tetralone with 74% isolated yield (run 3). In the same conditions, the oxidation of indan-1-ol was very slow, yielding only 15% of indan-1-one after 24 h. Therefore, a direct pathway leading from indan to indan-1-one seem likely (run 4).⁸ 4-Methoxy-1-ethylbenzene and fluorene gave 60% and 42% conversions after 6 h and 24 h oxidation. An additional 1 mol% NHTPPI raised the conversions to 79 and 84% and allowed isolation of 70% of 4-methoxyacetophenone and of 79% fluorenone (runs 5 and 6). Oxidation of a methyl substituent was less efficient: 4-methylanisol gave 25% of *p*-anisaldehyde, accompanied by 9% of *p*-anisic acid after 48 h reaction (run 7). Selective oxidation of one ethyl group of 1,4-diethylbenzene was observed, resulting in the isolation of 52% 4-ethylacetophenone using 1 mol% NHTPPI (run 8). This results, clearly indicates that a carbonyl substituent deactivates benzylic sites to oxidation. Finally, xanthone and isochromanone were obtained with excellent 96 and 86% isolated yields from xanthone and isochroman with only 1 mol% NHTPPI (run 9 and 10). In each case the main oxidation compound was accompanied by only small amounts (1–3%) of the corresponding alcohol.

To obtain some information on the cause of enhanced catalytic properties of NHTPPI compared to NHPI, we investigated generation and self decomposition of the corresponding radical **10b** in acetonitrile at 35 °C in the absence of substrate, measuring the absorbance changes by UV-vis spectrophotometry. Ammonium hexanitratocerate(IV) (CAN)⁹ has been found to be the most convenient oxidant to generate **10b** in CH₃CN: when CAN was added progressively to a solution of **10a** a new absorption band with a maximum at 425 nm appeared. It increased until 1 equiv. of CAN had been added.

This band has been attributed to radical **10b**, as its fast appearance was followed by slower decreasing accounting for self-

decomposition of the radical. The absorbance change was monitored at 425 nm (ESI†). Second-order kinetics fitting^{2c} gave second-order rate constant $k_d = 0.162 \text{ L mol}^{-1} \text{ s}^{-1}$ at 35 °C. For $[\mathbf{10b}]_0 = 0.5 \text{ mmol L}^{-1}$, the half life ($t_{1/2}$) of **10b** is 205 min.

For a comparison, self-decomposition of **1b** (PINO) in the same conditions of temperature and concentration, monitored at 380 nm gave $k_d = 0.777 \text{ L mol}^{-1} \text{ s}^{-1}$ and $t_{1/2} = 43 \text{ min}$ (ESI†). So, the half life of **10b** was found to be 4.7 times longer than that of PINO at 35 °C. This result has to be compared with recent results of Espenson: he showed that PINO radicals substituted either by electron donating or by electron withdrawing substituents have lower kinetic stabilities than unsubstituted PINO, accounting for lower efficiency of the corresponding NHPI catalysts.^{2e} In our case indeed, enhanced kinetic stability of the radical species is accompanied by enhanced catalytic activity of NHTPPI.

In conclusion we have developed a new catalytic system for aerobic oxidation using NHTPPI/CuCl. Enhanced kinetic stability of the corresponding radicals allows efficient catalysis with low catalyst loading. Further extension of this approach is underway.

We thank Professor J.-L. Pierre for fruitful discussions.

Notes and references

† To our knowledge the use of CuCl in association with *N*-hydroxyimides for catalytic oxidations has been restricted to our own work on NHPI chiral analogues.⁵ Its precise role has not yet been established. However, this role can tentatively be related to the formation of μ -oxocopper(II) species by reaction of CuCl with dioxygen in acetonitrile.⁶ These species may generate PINO or **10b** radicals from NHPI or NHTPPI. Further investigations concerning this point are underway.

§ Valuable NHPI analogues have been described by Ishii, bearing several *N*-hydroxyimide functionalities in the same molecule,^{4d,f} or a masked *N*-hydroxyimide functionality.^{4e} In the present study, we have restricted our choice to NHPI analogues with a single free *N*-hydroxyimide moiety.

Table 2 Oxidation of various substrate catalysed by NHTPPI/CuCl^a

Run	Substrate	Time/h	Conv. ^b (%)	Yield ^c (%)
1	Indan	6	83	77 (93)
2	2-Methylindan	27	73	65 (90)
3	Tetralin	18	84	74 (88)
4	Indan-1-ol	24	15	15 (100)
5	4-Methoxy-1-ethylbenzene	23	79	70 (88) ^d
6	Fluorene	48	84	79 (94) ^d
7	4-Methylanisol	48	34	25 (70) ^e
8	1,4-Diethylbenzene	8	69	52 (75)
9	Xanthone	18	98	96 (98)
10	Isochroman	7	93	86 (92)

^a Standard procedure: 1 mmol of substrate, 0.01 mmol of NHTPPI, 0.05 mmol of CuCl in 10 ml of acetonitrile with O₂ at atmospheric pressure and at 35 °C. ^b Determined by GC. ^c Yield of pure oxidation product after column chromatographic purification. Numbers in parentheses give the yield based on the substrate reacted. ^d 0.02 mmol of NHTPPI were used. ^e In addition to 25% of *p*-anisaldehyde, 9% of *p*-anisic acid was also obtained.

- Y. Ishii, S. Sakaguchi and T. Iwahama, *Adv. Synth. Catal.*, 2001, **343**, 379.
- (a) R. Arnaud, A. Milet, C. Adamo, C. Einhorn and J. Einhorn, *J. Chem. Soc., Perkin Trans.*, 2002, **2**, 1967; (b) R. Amorati, M. Lucarini, V. Mugnaini, G.-F. Pedulli, F. Minisci, F. Recupero, F. Fontana, P. Astolfi and L. Greci, *J. Org. Chem.*, 2003, **68**, 1747; (c) N. Koshino, B. Saha and J. H. Espenson, *J. Org. Chem.*, 2003, **68**, 9364; (d) N. Koshino, Y. Cai and J. H. Espenson, *J. Phys. Chem. A*, 2003, **107**, 4262; (e) B. Saha, N. Koshino and J. H. Espenson, *J. Phys. Chem. A*, 2004, **108**, 425.
- C. Ueda, M. Noyama, H. Ohmori and M. Masui, *Chem. Pharm. Bull.*, 1987, **35**, 1372.
- (a) K. Gorgy, J.-C. Lepretre, E. Saint-Aman, C. Einhorn, J. Einhorn, C. Marcadal and J.-L. Pierre, *Electrochem. Acta*, 1998, **44**, 385; (b) B. B. Wentzel, M. P. J. Donners, P. L. Alster, M. C. Feiters and R. J. M. Nolte, *Tetrahedron*, 2000, **56**, 7797; (c) N. Sawatari, T. Yokota, S. Sakaguchi and Y. Ishii, *J. Org. Chem.*, 2001, **66**, 7889; (d) A. Shibamoto, S. Sakaguchi and Y. Ishii, *Tetrahedron Lett.*, 2002, **43**, 8859; (e) N. Sawatari, S. Sakaguchi and Y. Ishii, *Tetrahedron Lett.*, 2003, **44**, 2053; (f) N. Hirai, N. Sawatari, N. Nakamura, S. Sakaguchi and Y. Ishii, *J. Org. Chem.*, 2003, **68**, 6587.
- C. Einhorn, J. Einhorn, C. Marcadal-Abadi and J.-L. Pierre, *J. Org. Chem.*, 1999, **64**, 4542.
- (a) M. Maumy and P. Capdevielle, *Bull. Soc. Chim. Fr.*, 1995, **132**, 734; (b) P. Capdevielle and M. Maumy, *Tetrahedron Lett.*, 1983, **24**, 5611.
- C. Einhorn, J. Einhorn and C. Marcadal-Abadi, *Synth. Commun.*, 2001, **31**, 741.
- For similar observations, see: C. Einhorn, J. Einhorn, C. Marcadal and J.-L. Pierre, *Chem. Commun.*, 1997, 447.
- S. Sakaguchi, T. Hirabayashi and Y. Ishii, *Chem. Commun.*, 2002, 516.