

Synthetic studies on (–)-lemonomycin: stereocontrolled construction of the 3,8-diazabicyclo[3.2.1] skeleton

Kentaro Rikimaru, Kazuki Mori, Toshiyuki Kan and Tohru Fukuyama*

Received (in Cambridge, UK) 30th September 2004, Accepted 1st November 2004

First published as an Advance Article on the web 8th December 2004

DOI: 10.1039/b415030a

Stereoselective synthesis of the pentacyclic key intermediate **22** for (–)-lemonomycin (**1**) has been accomplished using the Ugi 4-CC reaction with our novel isocyanide **8**, a cross-metathesis of **13** and allylsilane and a subsequent intramolecular Hosomi–Sakurai type reaction.


(–)-Lemonomycin (**1**) is a tetrahydroisoquinoline alkaloid¹ isolated from *Streptomyces candidus* (LL-AP191).² Although the isolation of **1** was achieved in 1964, the structure determination was not reported until 2000 by researchers at Wyeth–Ayerst.³ It was also discovered that the compound possessed interesting antibiotic activity against methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus faecium* (VREF), as well as cytotoxicity against a human colon tumor cell line (HCT-116).³ Because of its potent biological activity and challenging structure, lemonomycin has become an attractive target for synthesis, and indeed the first total synthesis of **1** was achieved by Stoltz and co-workers in 2003.⁴

During the course of our total synthesis of ecteinascidin 743,⁵ we established an efficient synthetic strategy for the tetrahydroisoquinoline alkaloids *via* the Ugi four-component condensation (4-CC)⁶ and the intramolecular Mizoroki–Heck reaction. Furthermore, an application of this protocol was demonstrated by the stereoselective construction of the highly strained 3,8-diazabicyclo[3.2.1]octane framework of (+)-naphthyridinomycin (**2**).⁷ In a subsequent synthetic investigation, we developed an efficient synthetic methodology for (–)-lemonomycin (**1**) and report herein the stereocontrolled synthesis of the tetracyclic key intermediate **22** for **1**.

The heart of our synthetic plan is illustrated in Scheme 1. Incorporation of a unique amino sugar moiety and transformation to a labile quinone and hemiaminal moiety would be carried out at later stages of the total synthesis. Thus, the tetracyclic compound **3** was designed as a key intermediate in our total synthesis of **1**. According to our synthetic study on **2**,⁷ the preparation of **3** from **4** would be readily achieved by hydration of the enamide of **4** and subsequent electrophilic cyclization between an electron-rich aromatic ring and an aldehyde. Since there is a significant difference between **1** and **2** in the stereochemistry at the C-15 position, the stereoselective construction of the bicyclo[3.2.1] system bearing an *exo*-oriented side chain in **4** would be a crucial step for the total synthesis of **1**.⁸


Our synthesis started from the two chiral amino acid derivatives **5**^{7,9} and **6**¹⁰ (see Scheme 2). In our synthetic investigation of the related alkaloids,^{5,7} condensation of the amino acid derivatives was

accomplished by the Ugi 4-CC reaction with *p*-methoxyphenyl isocyanide and acetaldehyde to access easily the corresponding diketopiperazine. However, we found that the Ugi 4-CC reaction of the isocyanide **8** and glyoxyaldehyde dimethylacetal **7** in trifluoroethanol¹¹ was more suitable for an efficient preparation of **13**. Thus, after conversion to the dipeptide **9**, the transformation to the cyclic enamide **10** was performed by treatment with CSA (10-camphorsulfonic acid) and quinoline. Although cleavage of the amide bond derived from the isocyanide of the Ugi adducts often requires harsh conditions, we discovered recently that the amide bond derived from the phenyl carbonate-type isocyanide was readily converted *via* the corresponding *N*-acyloxazolidinone derivatives.¹² Thus, upon treatment of the amidocarbonate **10** with *t*-BuOK, the oxazolidinone formation proceeded smoothly to provide **11** with release of phenol. The imide **11** was readily converted to the hydroxymethyl derivative **12** by simple reduction with NaBH₄. Changing the protecting group from TIPS to Ac proceeded smoothly in one step to provide **13** in high yield. For the crucial construction of the bicyclo[3.2.1] system, we found that an acyliminium ion-mediated cyclization was more suitable to obtain the desired stereochemistry.¹³ The elaboration of the cyclization precursor **15** was achieved by a cross-metathesis of **13** and allyltrimethylsilane in the presence of 2 mol% of the Grubbs 2nd generation catalyst (**14**).^{14,15} Upon treatment of **15** with BF₃·Et₂O, generation of the conjugate acyliminium cation **16** and subsequent cyclization of the allylsilane moiety proceeded immediately to



Scheme 1 Structure and synthetic strategy of lemonomycin (**1**).

*fukuyama@mol.f.u-tokyo.ac.jp



Scheme 2 Reagents and conditions: a) CF₃CH₂OH, rt to 50 °C; b) CSA, quinoline, PhMe, reflux, 73% (2 steps); c) *t*-BuOK, MS4A, THF, 0 °C; d) NaBH₄, THF–H₂O (3 : 1); e) TBAF, THF, 50 °C; Ac₂O, Py, 68% (3 steps); f) Grubbs 2nd generation cat. (**14**) (2 mol%), allyltrimethylsilane, CH₂Cl₂, reflux, 51%; g) BF₃·OEt₂, CH₂Cl₂, –78 °C, 95%; h) TMSOTf, CH₂Cl₂, 0 °C; i) CbzCl, DMAP, MeCN; j) K₂CO₃, MeOH; k) TIPSOTf, 2,6-lutidine, CH₂Cl₂, 0 °C, 55% (4 steps); l) DMDO (dimethyldioxirane), Na₂SO₄, MeOH, –78 °C to rt; CSA; m) NaBH₃CN, TFA–THF (9 : 1), 0 °C; n) KOSiMe₃, MeCN, 0 °C; BnBr, 50 °C, 53% (4 steps); o) DMP (Dess–Martin periodinane), CH₂Cl₂; p) TFA–CH₂Cl₂ (1 : 9), 76% (2 steps).

provide **17** with complete stereoselectivity¹⁶ in almost quantitative yield.

With the desired enamide **17** in hand, we next focused on the stereoselective construction of the tetracyclic system of **22**. According to our synthetic study on **2**,⁷ the cyclic enamide was oxidized by DMDO¹⁷ in methanol followed by addition of CSA to give the hemiaminal. However, acidic reduction of the aminal moiety, which was demonstrated in the synthetic studies on naphthyridinomycin, led to decomposition. Thus, after exchange of the protecting groups from Boc to Cbz and from Ac to TIPS ether, DMDO oxidation of **18** provided **19**. The acyliminium ion-mediated reduction of **19** occurred from the less hindered *exo*-face of the molecule to afford **20** as a single isomer with the correct stereochemistry. Since the reactivity of the aromatic ring was dependent on its electron-donating nature, the protecting group of the phenol **20** was changed from the mesyl to the corresponding benzyl group. After oxidation of the alcohol **21** with Dess–Martin periodinane, treatment of the resulting aldehyde with TFA allowed the desired cyclization to proceed smoothly providing **22**, which implies all the carbon atoms with the requisite stereochemistry needed for the lemonomycin (**1**) aglycon.

In summary, we have accomplished the efficient synthesis of the tetracyclic backbone **22** of (–)-lemonomycin (**1**), including the

highly strained bicyclo[3.2.1]octane framework. The present synthesis features the Ugi 4-CC reaction with our novel isocyanide **8**, which allows the easy preparation of the cyclic enamide **13**. The combination of cross-metathesis for a facile preparation of the allylsilane **15** and the intramolecular Hosomi–Sakurai type reaction provided the bicyclo[3.2.1]octane framework **17** with complete stereoselectivity. Further conversion of **22** to (–)-lemonomycin (**1**) is currently under investigation in our laboratory.

This work was financially supported by the 21st Century COE Program and a Grant-in-Aid from the Ministry of Education, Science, Sports, Culture and Technology, Japan.

Kentaro Rikimaru, Kazuki Mori, Toshiyuki Kan and Tohru Fukuyama*
 Graduate School of Pharmaceutical Sciences, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.
 E-mail: fukuyama@mol.f.u-tokyo.ac.jp; Fax: +81-3-5802-8694;
 Tel: +81-3-5841-4777

Notes and references

- For a recent review on tetrahydroisoquinoline alkaloids, see: J. D. Scott and R. M. Williams, *Chem. Rev.*, 2002, **102**, 1669.
- H. A. Whaley, E. L. Patterson, M. Dann, A. J. Shay and J. N. Porter, *Antimicrob. Agents Chemother.*, 1964, **8**, 83.

- 3 H. He, B. Shen and G. T. Carter, *Tetrahedron Lett.*, 2000, **41**, 2067.
- 4 E. R. Ashley, E. G. Cruz and B. M. Stoltz, *J. Am. Chem. Soc.*, 2003, **125**, 15000.
- 5 (a) A. Endo, A. Yanagisawa, M. Abe, S. Tohma, T. Kan and T. Fukuyama, *J. Am. Chem. Soc.*, 2002, **124**, 6552; (b) T. Kan, *J. Syn. Org. Chem., Jpn.*, 2003, **61**, 949.
- 6 For reviews on Ugi 4-CC reaction and other multicomponent reactions with isocyanides, see (a) C. Hulme and V. Gore, *Curr. Med. Chem.*, 2003, **10**, 51; (b) A. Dömling and I. Ugi, *Angew. Chem. Int. Ed.*, 2000, **39**, 3168; (c) G. Gokel, G. Lüdke and I. Ugi, in *Isonitrile Chemistry*, ed. I. Ugi, Academic, New York, 1971, p. 145.
- 7 K. Mori, K. Rikimaru, T. Kan and T. Fukuyama, *Org. Lett.*, 2004, **6**, 3095.
- 8 The selective production of both stereochemistries would be significant, since both stereochemistries were observed in this family of natural products. While the *exo*-oriented compounds belong to the quinocarcin family (lemonomycin, quinocarcin and tetrazomine), the corresponding *endo*-oriented compounds belong to the naphthyridinomycin family (naphthyridinomycin and dnacins). See ref. 2.
- 9 Preparation of the arylglycinol derivative **5** was performed according to the method recently developed by us. See: (a) S. Tohma, A. Endo, T. Kan and T. Fukuyama, *Synlett*, 2001, 1179; (b) S. Tohma, K. Rikimaru, A. Endo, K. Shimamoto, T. Kan and T. Fukuyama, *Synthesis*, 2004, 909.
- 10 Preparation of the crotyl glycine derivative **6** was performed according to the Pleixats method. See: A. López and R. Pleixats, *Tetrahedron: Asymmetry*, 1998, **9**, 1967.
- 11 Use of the Ugi 4-CC reaction in methanol or other solvents resulted in low yield due to the large amount of by-products, generated from solvolysis and/or hydrolysis of the intermediates.
- 12 K. Rikimaru, A. Yanagisawa, T. Kan and T. Fukuyama, *Synlett*, 2004, 41.
- 13 A similar acyliminium cation mediated intramolecular cyclization has been reported and the *exo*-oriented bicyclo[3.2.1]octane ring system was also obtained. See: (a) T. Fukuyama and J. J. Nunes, *J. Am. Chem. Soc.*, 1988, **110**, 5196; (b) J. J. N. Veerman, R. S. Bon, B. T. B. Hue, D. Girones, F. P. J. T. Rutjes, J. H. van Maarseveen and H. Hiemstra, *J. Org. Chem.*, 2003, **68**, 4486.
- 14 M. Scholl, S. Ding, C. W. Lee and R. H. Grubbs, *Org. Lett.*, 1999, **1**, 953.
- 15 A similar cross-metathesis with the allyl derivative of **13** (without the terminal methyl group) resulted in a low yield due to decomposition.
- 16 The stereochemistry was confirmed by a strong NOE between H-4 and H-15 in the Boc derivative of **19**.
- 17 R. W. Murray and M. Singh, *Org. Synth.*, 1998, **Coll. Vol. IX**, 288.