

# Carboxylation of anisole derivatives with CO and O<sub>2</sub> catalyzed by Pd(OAc)<sub>2</sub> and molybdovanadophosphates

Shinichiro Ohashi, Satoshi Sakaguchi and Yasutaka Ishii\*

Received (in Corvallis, OR, USA) 3rd August 2004, Accepted 11th October 2004

First published as an Advance Article on the web 2nd December 2004


DOI: 10.1039/b411934g

Anisole and its homologues were carboxylated under the influence of CO and O<sub>2</sub> catalyzed by Pd(OAc)<sub>2</sub> combined with molybdovanadophosphates (HPMoV) under mild conditions to give the corresponding carboxylic acids in fair to good yields; for instance, anisole underwent the carboxylation under a mixed gas of CO (0.5 atm) and O<sub>2</sub> (0.5 atm) in the presence of Pd(OAc)<sub>2</sub> (5 mol%) and H<sub>3</sub>PMo<sub>10</sub>V<sub>2</sub>O<sub>40</sub>·*n*H<sub>2</sub>O (2 mol%) to form an isomeric mixture of anisic acids in good yield.

Benzenes react stoichiometrically with Pd(II) salts to undergo chlorination,<sup>1</sup> acetoxylation,<sup>2</sup> carbonylation,<sup>3</sup> carboxylation,<sup>4</sup> etc. The direct carboxylation of benzene with CO was first reported by Fujiwara *et al.* by the use of a stoichiometric amount of Pd(OAc)<sub>2</sub>.<sup>5</sup> Thereafter, they reported the Pd-catalyzed carboxylation of benzene derivatives with CO using *t*-BuOOH<sup>6</sup> or K<sub>2</sub>S<sub>2</sub>O<sub>8</sub><sup>7</sup> as reoxidants in trifluoroacetic acid. Pd(II)-catalyzed carboxylation of hydrocarbons was reviewed by their group.<sup>8</sup> However, carboxylation using trifluoroacetic acid as a solvent is difficult to carry out on a large scale, since trifluoroacetic acid is an intractable compound. If molecular oxygen as an oxidant and acetic acid as a solvent can be used in place of K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> and trifluoroacetic acid, respectively, in the carboxylation of benzene derivatives, such a method would provide an important practical synthetic route to the corresponding benzoic acids. Recently, we have found that Pd(OAc)<sub>2</sub> combined with molybdovanadophosphoric acids (HPMoV) is an efficient catalytic system for the direct activation of the C–H bond of arenes using molecular oxygen as the reoxidant. Thus, the Heck–Mizoroki reaction of benzene with acrylate was first achieved by using the Pd(OAc)<sub>2</sub>/HPMoV/O<sub>2</sub> system.<sup>9</sup> We have now found that the Pd(OAc)<sub>2</sub>/HPMoV system catalyzes efficiently the carboxylation of anisole derivatives under the influence of a mixed gas of CO and O<sub>2</sub> to give the corresponding anisic acids in fair to good yields (Scheme 1).

In order to confirm the optimum reaction conditions, anisole (**1**) was allowed to react under a mixed gas of CO and O<sub>2</sub> in the presence of catalytic amounts of Pd(OAc)<sub>2</sub> and HPMoV in acetic acid under various conditions (Table 1).

The reaction of **1** under a 1 : 1 mixture of CO (0.5 atm) and O<sub>2</sub> (0.5 atm) in the presence of catalytic amounts of Pd(OAc)<sub>2</sub> (5 mol%) and H<sub>3</sub>PMo<sub>10</sub>V<sub>2</sub>O<sub>40</sub>·15.2H<sub>2</sub>O (HPMo<sub>10</sub>V<sub>2</sub>) (2 mol%) in acetic acid at 70 °C for 15 h afforded a 74 : 26 isomeric mixture of *p*- and *o*-anisic acids (*p*- and *o*-**2**) in 88% conversion and 97% selectivity (Run 2).<sup>†</sup> This is the first successful direct carboxylation of benzene derivatives like **1** using CO and O<sub>2</sub> catalyzed by Pd(OAc)<sub>2</sub> and HPMoV. It is important to note that the



Scheme 1

carboxylation of **1** by the use of CO and O<sub>2</sub> can be achieved by the Pd(OAc)<sub>2</sub>/HPMoV system, since the same reaction using PdCl<sub>2</sub>/CuCl<sub>2</sub><sup>5</sup> under these conditions provided no carboxylated products at all (Run 10). The carboxylation was carried out under varying pressures of CO and O<sub>2</sub> (Runs 1 to 3). It was found that the carboxylation of **1** to **2** was considerably influenced by the partial pressures of CO and O<sub>2</sub>. The best result was obtained when a 1 : 1 mixed gas of CO and O<sub>2</sub> was employed as shown in Run 2. The reaction under a 1 : 2 mixture of CO (0.33 atm) and O<sub>2</sub> (0.67 atm) resulted in the decrease of conversion of **1** and the selectivity of **2**, although the product ratio was almost the same (Run 1). Under a 2 : 1 mixture of CO (0.67 atm) and O<sub>2</sub> (0.33 atm), the conversion of **1** to **2** was considerably decreased (Run 3). When the amount of Pd(OAc)<sub>2</sub> was halved under these conditions, the conversion of **1** was decreased to 44%, but the selectivity to **2** was

Table 1 Carboxylation of anisole (**1**) to anisic acid (**2**) with CO and O<sub>2</sub> by Pd(II) combined with HPMo<sub>10</sub>V<sub>2</sub> under various conditions<sup>a</sup>

Run	Pd(II)	CO : O <sub>2</sub> (atm)	Solv.	Conv. (%)	Select. <sup>b</sup> (%)
1	Pd(OAc) <sub>2</sub>	0.33 : 0.67	AcOH	80	88 (23 : 77)
2	Pd(OAc) <sub>2</sub>	0.50 : 0.50	AcOH	88	97 (26 : 74)
3	Pd(OAc) <sub>2</sub>	0.67 : 0.33	AcOH	17	20 (26 : 74)
4 <sup>c</sup>	Pd(OAc) <sub>2</sub>	0.50 : 0.50	AcOH	44	96 (26 : 74)
5 <sup>d</sup>	Pd(OAc) <sub>2</sub>	0.50 : 0.50	AcOH	83	99 (23 : 77)
6 <sup>e</sup>	Pd(OAc) <sub>2</sub>	0.50 : 0.50	AcOH	74	91 (23 : 77)
7	Pd(OAc) <sub>2</sub>	0.50 : 0.50	EtCOOH	81	85 (23 : 77)
8	Pd(OAc) <sub>2</sub>	0.50 : 0.50	CH <sub>3</sub> CN	4	n.d.
9	Pd(OAc) <sub>2</sub>	0.50 : 0.50	DMF	11	n.d.
10 <sup>f</sup>	PdCl <sub>2</sub>	0.50 : 0.50	AcOH	no reaction	
11	Pd(SO <sub>4</sub> ) <sub>2</sub>	0.50 : 0.50	AcOH	no reaction	

<sup>a</sup> **1** (2 mmol) was allowed to react in the presence of Pd(OAc)<sub>2</sub> (0.1 mmol), HPMo<sub>10</sub>V<sub>2</sub> (0.04 mmol) in solvent (7 mL) at 70 °C for 15 h. <sup>b</sup> Numbers in parentheses show the ratio of *o*-**2** and *p*-**2**. <sup>c</sup> Pd(OAc)<sub>2</sub> (0.05 mmol) was used. <sup>d</sup> At 80 °C. <sup>e</sup> At 90 °C. <sup>f</sup> CuCl<sub>2</sub> (0.04 mmol) was used instead of HPMo<sub>10</sub>V<sub>2</sub>.

\*ishii@ipcku.kansai-u.ac.jp

not changed (Run 4). The conversion of **1** to **2** was slightly decreased in the reaction at 80 °C, but the reaction at 90 °C under these conditions resulted in the decrease of the conversion of **1** (Runs 5 and 6). The formation of *m*-anisic acid was not observed at all in the present carboxylation.

The reaction was examined in several solvents such as propionic acid, CH<sub>3</sub>CN and DMF. In a previous paper on the direct coupling of benzene with ethyl acrylate by Pd(OAc)<sub>2</sub> and HPMoV, propionic acid was a better solvent than acetic acid. However, the reaction in propionic acid resulted in a slight lowering of the conversion and selectivity (Run 7). It was found that CH<sub>3</sub>CN and DMF were not suited for the present carboxylation (Runs 8 and 9). PdCl<sub>2</sub> and PdSO<sub>4</sub> combined with HPMo<sub>10</sub>V<sub>2</sub> did not catalyze at all the present carboxylation (Runs 10 and 11).

We next tried the carboxylation of **1** by Pd(OAc)<sub>2</sub> combined with several heteropoly acids (Table 2).

In the absence of HPMoV as a co-catalyst under the same conditions, no reaction took place using Pd(OAc)<sub>2</sub> (Run 1). The carboxylation of **1** by Pd(OAc)<sub>2</sub> was found to be considerably affected by the vanadium content in HPMoV used (Runs 2 to 7). No reaction was promoted by Pd(OAc)<sub>2</sub> combined with H<sub>3</sub>PMo<sub>12</sub>O<sub>40</sub> (HPMo<sub>12</sub>) not containing vanadium (Run 2). The carboxylation of **1** was promoted by Pd(OAc)<sub>2</sub> combined with H<sub>4</sub>PMo<sub>11</sub>V<sub>1</sub>O<sub>40</sub>·*n*H<sub>2</sub>O (HPMo<sub>11</sub>V<sub>1</sub>) and H<sub>6</sub>PMo<sub>9</sub>V<sub>3</sub>O<sub>40</sub>·*n*H<sub>2</sub>O (HPMo<sub>9</sub>V<sub>3</sub>) containing vanadium<sup>10</sup> to give **2** in 19% and 43% yields, respectively (Runs 3 and 4). Although the reaction was examined using molybdotungstophosphoric acids (HPMo<sub>11</sub>W<sub>1</sub> and HPMo<sub>10</sub>W<sub>2</sub>) and vanadotungstophosphoric acid (HPW<sub>10</sub>V<sub>2</sub>) under these conditions, no carboxylation was observed (Runs 6 to 8). These results show that vanadium in heteropolyoxometalates is an essential component to catalyze the present carboxylation of **1**. Among heteropoly acids examined, HPMo<sub>10</sub>V<sub>2</sub> was the best combination with Pd(OAc)<sub>2</sub>. Although several vanadium compounds such as V<sub>2</sub>O<sub>5</sub> and VO(acac)<sub>2</sub> were employed instead of HPMoV, no carboxylation took place (Runs 9 and 10)

On the basis of these results, the carboxylation of several anisole derivatives was carried out by the use of Pd(OAc)<sub>2</sub> and HPMo<sub>10</sub>V<sub>2</sub> in acetic acid at 70 °C for 15 h (Table 3).

In order to obtain information on the reaction path, the carboxylation of **1** was compared with that of *o*-, *m*- and *p*-methylanisoles (*o*-, *m*-, and *p*-**3**). The reactivity in the carboxylation was found to decrease in the order of *o*-**3**, *m*-**3**, and *p*-**3**. The

**Table 2** Carboxylation of anisole (**1**) to anisic acid (**2**) with CO and O<sub>2</sub> catalyzed by Pd(OAc)<sub>2</sub> and heteropoly acids<sup>a</sup>

Run	HPA	Conv. (%)	Select. (%)
1	—	no reaction	—
2	HPMo <sub>12</sub>	2	—
3	HPMo <sub>11</sub> V <sub>1</sub>	30	64
4	HPMo <sub>9</sub> V <sub>3</sub>	53	81
5	HPMo <sub>8</sub> V <sub>4</sub>	27	57
6	HPMo <sub>11</sub> W <sub>1</sub>	1	—
7	HPMo <sub>10</sub> W <sub>2</sub>	2	—
8	HPW <sub>10</sub> V <sub>2</sub>	no reaction	—
9	V <sub>2</sub> O <sub>5</sub>	no reaction	—
10	VO(acac) <sub>2</sub>	no reaction	—


<sup>a</sup> **1** (2 mmol) was allowed to react under CO (0.5 atm) and O<sub>2</sub> (0.5 atm) by Pd(OAc)<sub>2</sub> (0.1 mmol) and HPA (0.04 mmol) in AcOH (7 mL) at 70 °C for 15 h.

**Table 3** Carboxylation of various anisoles and phenol with CO and O<sub>2</sub> by the Pd(OAc)<sub>2</sub>/HPMo<sub>10</sub>V<sub>2</sub> system<sup>a</sup>

Run	Substrate	Products	Conv. (%)	Select. (%)
1			90	73
2 <sup>b</sup>			82	72
3			49	64
4			90	75
5			89	92
6 <sup>c</sup>			90	59

<sup>a</sup> Substrate (2 mmol) was allowed to react under CO (0.5 atm) and O<sub>2</sub> (0.5 atm) by Pd(OAc)<sub>2</sub> (0.1 mmol) and HPMo<sub>10</sub>V<sub>2</sub> (0.04 mmol) in AcOH (7 mL) at 70 °C for 15 h. <sup>b</sup> *m*-**4** : *m*-**4'** = *ca.* 50 : 50. <sup>c</sup> In AcOH (4 mL) at 60 °C; *p*-**10** : *o*-**10** = *ca.* 90 : 10.

selectivity in the carboxylation could be satisfactorily explained by the electronic effect of the methoxy substituent. For instance, the carboxylation of *m*-**3** took place at the 4-position possessing the highest electron density. The low reactivity of *p*-**3** is due to the occupation of its *para*-position by the methyl group. Additionally, *p*-hydroxybenzoic acid (*p*-**10**) and *o*-hydroxybenzoic acid (*o*-**10**) were obtained in 59% selectivity at 90% conversion in the reaction of phenol (**9**) (Run 6). Unfortunately, however, benzene is inert for the present carboxylation, giving a small amount of benzoic acid (< 5%).



**Scheme 2**

The carboxylation is rationally explained by a similar reaction path proposed by Fujiwara<sup>7b</sup> (Scheme 2). The *ortho-para* orientation of the carboxylation of **1** by the present reaction system suggests that the reaction involves the electrophilic substitution of aromatic C–H bonds by Pd(II)<sup>‡</sup> followed by the CO insertion of the Ar–Pd(II) species leading to an arylpalladium species (**A**). The subsequent reductive elimination of Pd(0) and acetic anhydride from **A** leads to anisic acid. The Pd(0) is reoxidized by the action of HPMoV and O<sub>2</sub> to the parent Pd(II) species.

In conclusion, we have developed Pd-catalyzed direct carboxylation of anisole derivatives using CO and O<sub>2</sub> under mild conditions.

**Shinichiro Ohashi, Satoshi Sakaguchi and Yasutaka Ishii\***  
 Department of Applied Chemistry, Faculty of Engineering & High  
 Technology Research Center, Kansai University, Suita, Osaka,  
 564-8680, Japan. E-mail: ishii@ipcku.kansai-u.ac.jp

## Notes and references

<sup>†</sup> Representative procedure: To an AcOH solution (7 mL) of Pd(OAc)<sub>2</sub> (0.1 mmol) and H<sub>5</sub>PMo<sub>10</sub>V<sub>2</sub>O<sub>40</sub>·15.2H<sub>2</sub>O (0.04 mmol) was added **1** (2 mmol). Then, the reaction mixture was stirred at 70 °C for 15 h under a 1 : 1 mixed gas (1 atm) of CO and O<sub>2</sub>. After the reaction, the GC and GC-MS analyses were performed. The conversions and yields of products were estimated from the peak areas based on the internal standard technique using GC.

<sup>‡</sup> Although we could not determine the real active Pd(II) species at this stage, a PdL<sub>2</sub> or a cationic Pd<sup>+</sup>L (L: OAc) may be formed under these reaction conditions.

- 1 P. Henry, *J. Org. Chem.*, 1971, **36**, 1886.
- 2 L. Ebersson and L. Gomez-Gonzales, *J. Chem. Soc., Chem. Commun.*, 1971, 263.
- 3 P. Henry, *Tetrahedron Lett.*, 1968, 2285.
- 4 T. Sakakibara and Y. Odaira, *J. Org. Chem.*, 1976, **41**, 2049.
- 5 Y. Fujiwara, T. Kawauchi and H. Taniguchi, *J. Chem. Soc., Chem. Commun.*, 1980, 220.
- 6 T. Jintoku, T. Taniguchi and Y. Fujiwara, *Chem. Lett.*, 1987, 1159.
- 7 (a) T. Jintoku, Y. Fujiwara, I. Kawata, T. Kawauchi and T. Taniguchi, *J. Organomet. Chem.*, 1990, **385**, 297; (b) Y. Taniguchi, Y. Yamaoka, K. Nakata, K. Takai and Y. Fujiwara, *Chem. Lett.*, 1995, 345 and literature cited therein.
- 8 C. Jia, T. Kitamura and Y. Fujiwara, *Acc. Chem. Res.*, 2001, **34**, 633; Y. Fujiwara, K. Takaki and Y. Taniguchi, *Synlett*, 1996, 591.
- 9 T. Yokota, M. Tani, S. Sakaguchi and Y. Ishii, *J. Am. Chem. Soc.*, 2003, **125**, 1476.
- 10 D. M. Fenton and P. J. Steinwand, *J. Org. Chem.*, 1972, **37**, 2034; O. Hamed, A. El-Qisairi and P. M. Henry, *J. Org. Chem.*, 2001, **66**, 180.