

# Reversible 1,4-cycloaddition of singlet oxygen to *N*-substituted 2-pyridones: 1,4-endoperoxide as a versatile chemical source of singlet oxygen†

Masakatsu Matsumoto,\* Masayo Yamada and Nobuko Watanabe

Received (in Cambridge, UK) 27th September 2004, Accepted 22nd October 2004

First published as an Advance Article on the web 2nd December 2004

DOI: 10.1039/b414845b

*N*-substituted pyridones (**1**) easily undergo singlet oxygenation to give exclusively the corresponding endoperoxides (**2**), which decompose to give pyridones again while liberating  $^1\text{O}_2$  in high yield.

$\alpha$ -Pyranones are known to undergo 1,4-cycloaddition with singlet oxygen ( $^1\text{O}_2$ ),<sup>1–3</sup> to afford 1,4-endoperoxides, which decompose thermally into 1,2-diacetylenes while extruding  $\text{CO}_2$  (Scheme 1). We report here that a)  $^1\text{O}_2$  adds also to nitrogen-analogues of  $\alpha$ -pyranone, namely, *N*-substituted pyridones (**1**) to give 1,4-endoperoxides (**2**) exclusively, b) the thus-obtained peroxides (**2**) liberate  $^1\text{O}_2$  in high yield, differing from the case of 1,4-endoperoxides of  $\alpha$ -pyranones, and c) the peroxides (**2**) are a promising chemical  $^1\text{O}_2$  source with characteristics different from those of 1,4-endoperoxides of substituted polynuclear aromatics.

Commercially available 2-hydroxypyridine and its 4-methyl-analogue were easily alkylated with benzyl bromide, 4-bromomethylbiphenyl, and *t*-butyl bromoacetate to give the corresponding *N*-substituted 2-pyridones (**1a–1e**) in high yields. When a solution of *N*-benzyl-2-pyridone (**1a**) (250 mg) in dichloromethane (10 mL) was irradiated with a Na-lamp (940 W) together with a catalytic amount of tetraphenylporphine (TPP) under an oxygen atmosphere at  $-78\text{ }^\circ\text{C}$  for 2 h, 1,4-endoperoxide


(**2a**) was produced exclusively. Chromatographic purification (silica gel/hexane–dichloromethane) was attained easily to give peroxide (**2a**) as colorless leaflets (mp  $67.0\text{--}68.0\text{ }^\circ\text{C}$ , dec.), the structure of which was determined by  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ , IR, and mass spectral analysis.‡ The other pyridones (**1b–1e**) gave also exclusively the corresponding 1,4-endoperoxides (**2b–2e**) on similar singlet oxygenation.

All endoperoxides (**2a–2e**) synthesized here were stable enough for handling at room temperature and storing for  $> 1$  year in a freezer though they decomposed into the corresponding pyridones (**1a–1e**) exclusively in a solvent such as  $\text{CDCl}_3$  at  $40\text{ }^\circ\text{C}$  with half-life of 2–6 h. When the thermolysis of **2a–2e** was carried out in the presence of typical substrates, namely, dihydrofuran (**3**) for 1,2-addition of  $^1\text{O}_2$ ,<sup>4</sup> isobenzofuran (**4**) for 1,4-addition of  $^1\text{O}_2$ ,<sup>5</sup> and 2,3-dimethylbut-2-ene (**5**) for “ene” reaction of  $^1\text{O}_2$ ,<sup>6</sup> dioxetane (**6**),<sup>§</sup> diketone (**7**), and hydroperoxide (**8**) were produced, respectively. These results showed that the retro-1,4-cycloaddition proceeded to liberate  $^1\text{O}_2$ .

Thus, we attempted to estimate the efficiency of  $^1\text{O}_2$  generation by means of a trapping experiment of  $^1\text{O}_2$  with an olefin (**5**) for the thermolysis of endoperoxides (**2a–2e**). First of all, we examined whether the re-combination of  $^1\text{O}_2$  with pyridones, formed during the thermolysis of **2a**, could be prevented substantially by the use of a highly reactive olefin (**5**). Sensitized photooxygenation of a mixture of **1a** ( $1.5 \times 10^{-2}\text{ M}$ ) and an equimolar amount of **5** in  $\text{CDCl}_3$  at  $30\text{--}40\text{ }^\circ\text{C}$  was confirmed to give exclusively a hydroperoxide (**8**) without formation of **2a** even after *ca.* 40% conversion of **5**.

A solution of endoperoxide (**2a**) ( $1.5 \times 10^{-2}\text{ M}$ ) and an olefin (**5**) ( $7.5 \times 10^{-2}\text{ M}$ ) in  $\text{CDCl}_3$  was heated, and the time-course of the formation of **8** and **1a** and of the decrease of **2a** was followed by means of  $^1\text{H-NMR}$  at 30, 40, and  $50\text{ }^\circ\text{C}$ .¶ Similar kinetic experiments were carried out for thermolysis of the other endoperoxides (**2b–2e**). All endoperoxides (**2a–2e**) were found to decompose into the corresponding 2-pyridones (**1a–1e**) by first-order kinetics and to afford  $^1\text{O}_2$  in high yields even at high conversion ( $\sim 80\%$ ). These results are summarized in Table 1 together with activation parameters for the thermolysis, which were estimated from the Arrhenius plots.

Reversible 1,4-cycloaddition of  $^1\text{O}_2$  to a conjugate system has been recognized as one rather characteristic feature for polynuclear aromatic hydrocarbons.<sup>7</sup> In fact, endoperoxides of 9,10-diphenylanthracenes and 1,4-dialkyl-naphthalenes are well known to generate  $^1\text{O}_2$  in yields from 30 to 95% on heating.<sup>8,9</sup> Turro and his co-workers have found for the thermolysis of these endoperoxides that a)  $\Delta S^\ddagger$  values range from positive to slightly negative,



Scheme 1

† Electronic supplementary information (ESI) available: experimental section. See <http://www.rsc.org/suppdata/cc/b4/b414845b/>  
\*matsumo-chem@kanagawa-u.ac.jp


**Table 1** Activation parameters and efficiency of  $^1\text{O}_2$  generation for thermolysis of *N*-substituted 2-pyridone endoperoxides (**2a–2e**) in  $\text{CDCl}_3$

	$\Delta G^\ddagger /$ $\text{kcal mol}^{-1}$	$\Delta H^\ddagger /$ $\text{kcal mol}^{-1}$	$\Delta S^\ddagger /$ $\text{cal K}^{-1} \text{mol}^{-1}$	$^1\text{O}_2 / \%$	
				A <sup>a</sup>	B <sup>b</sup>
<b>2a</b>	24.3	22.8	-5.3	96 ± 1	81 ± 1
<b>2b</b>	24.1	23.0	-3.7	96 ± 1	73 ± 1
<b>2c</b>	24.7	23.2	-5.0	97 ± 1	84 ± 4
<b>2d</b>	24.6	24.1	-1.4	89 ± 2	74 ± 6
<b>2e</b>	25.2	25.2	0.1	84 ± 1	82 ± 1


<sup>a</sup> Mean of  $^1\text{O}_2$  yields for < 20% conversion of **2** at 40 °C. <sup>b</sup> Mean of  $^1\text{O}_2$  yields at 30, 40, and 50 °C.  $^1\text{O}_2$  yield at each temperature was estimated from the data at five points for 30–80% conversion of **2**.

and 2) positive  $\Delta S^\ddagger$  values correlate with the relatively low yields of  $^1\text{O}_2$ , and slightly negative or near zero  $\Delta S^\ddagger$  values correlate with the high yields of  $^1\text{O}_2$ .<sup>8,9</sup> They have also suggested that these tendencies can be interpreted in terms of a diradical mechanism which leads to a low yield of  $^1\text{O}_2$  and a concerted mechanism which leads to high yields of  $^1\text{O}_2$ , respectively.

$\Delta S^\ddagger$  values ranged from slightly positive to negative for endoperoxides of pyridones (**2a–2e**). Furthermore, column A in Table 1 shows that the liberation of  $^1\text{O}_2$  occurred nearly quantitatively for most cases of thermolysis of **2** at low conversion of **2** (< 20%). These relationships between the yield of  $^1\text{O}_2$  and the  $\Delta S^\ddagger$  value are illustrated in Fig. 1. One would realize from Fig. 1 that endoperoxides of pyridones (**2**) (squares) belong to the same class which includes endoperoxides of polynuclear aromatics for the mechanism of  $^1\text{O}_2$  liberation. On the other hand, the efficiency of  $^1\text{O}_2$  generation decreased considerably at high conversion of **2** except **2e** (~80%), though the absolute yields of  $^1\text{O}_2$  stayed at a high level. When the thermolysis of **2a** was carried out in the presence of an equimolar amount of 2-pyridone (**1a**),  $^1\text{O}_2$  yield decreased to *ca.* 80% at low conversion of **2a**. Thus, the decrease of  $^1\text{O}_2$  yields is likely to be attributable to the fact that 2-pyridone (**1**) produced during the thermolysis of **2** quenches  $^1\text{O}_2$  physically, though detailed experiments would be required to confirm such a conclusion.



**Fig. 1** Relationship between  $^1\text{O}_2$  yield and activation entropy for thermolysis of endoperoxides of polynuclear aromatics and pyridone endoperoxides (**2**).



**Scheme 2**

Chemical  $^1\text{O}_2$  sources which are less complicated mechanistically are quite useful for  $^1\text{O}_2$  reactions of biological substrates, mechanistic investigations of  $^1\text{O}_2$  reactions with organic substrates, and as an essential chemical species for COIL (chemical oxygen–iodine laser).<sup>10</sup> As described earlier, the present endoperoxides (**2**) would be useful as a new chemical source of  $^1\text{O}_2$ . Comparing **2** with 1,4-endoperoxides of alkylnaphthalene derivatives<sup>11,12</sup> as chemical  $^1\text{O}_2$  sources, one advantageous feature of **2** probably lies in the easy preparation and modification of their precursors, namely, *N*-substituted 2-pyridones (**1**), since the synthesis of most substituted naphthalenes is unexpectedly burdensome. A good example is the preparation of a polymer-supported  $^1\text{O}_2$  generator.

Functionalized polymer represented by Merrifield's resin is often used in a wide variety of reactions in the solid phase. 2-Pyridone was easily joined to Merrifield's resin to give **1f** by means of nucleophilic substitution similar to the synthesis of **1a–1e**. The thus-prepared pyridone supported on polymer (**1f**) was also easily converted into the corresponding endoperoxide (**2f**) by sensitized photooxygenation similar to the case of **1a–1e** except that a suspension of the substrate was used (Scheme 2).<sup>\*\*</sup> When endoperoxide supported on Merrifield's resin (**2f**) was heated with olefin (**5**) at 40 °C for 1 h,  $^1\text{O}_2$  was estimated to be liberated in 85% yield based on produced hydroperoxide (**8**) and the consumed endoperoxide.

In conclusion, we have shown here that *N*-substituted pyridones (**1**) easily undergo singlet oxygenation to give exclusively the corresponding endoperoxides (**2**), which decompose to give pyridones again while liberating  $^1\text{O}_2$  in high yields. Endoperoxides (**2**) have been shown to be a new type of chemical  $^1\text{O}_2$  source, preparation and structural modification of which are very easy and the efficiency of  $^1\text{O}_2$  liberation from which is high. Finally, it should be noted that a water-soluble and neutral  $^1\text{O}_2$  generator, such as an amide derivative (**2g**), was also obtained.

Masakatsu Matsumoto,\* Masayo Yamada and Nobuko Watanabe  
 Department of Materials Science, Kanagawa University, Tsuchiya,  
 Hiratsuka, Kanagawa, 259-1293, Japan.  
 E-mail: matsumo-chem@kanagawa-u.ac.jp

## Notes and references

‡ Selected data for **2a**:  $^1\text{H-NMR}$  (400 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{H}}$  4.42 (d,  $J = 15.3$  Hz, 1H), 4.87 (d,  $J = 15.3$  Hz, 1H), 5.08 (ddd,  $J = 5.8, 2.1, 0.6$  Hz, 1H), 5.53 (ddd,  $J = 5.2, 2.1, 0.6$  Hz, 1H), 6.71 (ddd,  $J = 7.8, 5.2, 2.1$  Hz, 1H), 6.76 (ddd,  $J = 7.8, 5.8, 2.1$  Hz, 1H), 7.19–7.24 (m, 2H), 7.28–7.39 (m, 3H) ppm;  $^{13}\text{C-NMR}$  (100 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{C}}$  46.9, 78.0, 83.2, 127.8, 127.9, 128.6, 128.8, 134.2, 135.1, 167.9 ppm; IR (KBr): 3089, 3031, 2932, 1691, 1426, 1350, 1158  $\text{cm}^{-1}$ ; Mass (EI, %): 217 ( $\text{M}^+$ , 25), 201 (70), 185 (76), 184 (34), 106 (34), 91 (100), 65 (20).

§ When an endoperoxide (**2**) ( $1.5 \times 10^{-2}$  M) and a dihydrofuran (**3**) ( $3.0 \times 10^{-3}$  M) were heated in  $\text{CDCl}_3$  for 2 h, a dioxetane (**6**) was produced in 12–56% yield.

¶ The reaction was carried out in an NMR sample tube without stirring.  
|| As Merrifield's resin, 1% cross-linked polystyrene (200–400 mesh, 1.2 mmol  $\text{Cl}^-/\text{g}$ ) was used. The 2-pyridone unit was estimated to be supported nearly quantitatively on the polymer from the difference between the pyridone used initially and that recovered.

\*\* Completion of singlet oxygenation was checked by monitoring the increase of the C=O peak ( $1710 \text{ cm}^{-1}$ ) due to endoperoxide of pyridone and disappearance of the C=O peak ( $1661 \text{ cm}^{-1}$ ) due to *N*-substituted 2-pyridone for the IR spectrum.

1 W. Adam and I. Erden, *Angew. Chem.*, 1978, **90**, 223.

2 (a) J. P. Smith and G. B. Schuster, *J. Am. Chem. Soc.*, 1978, **100**, 2564; (b) J. P. Smith, A. K. Schrock and G. B. Schuster, *J. Am. Chem. Soc.*, 1982, **104**, 1041.

3 W. Adam and I. Erden, *J. Am. Chem. Soc.*, 1979, **101**, 5692–5696.

4 (a) M. Matsumoto, H. Murakami and N. Watanabe, *Chem. Commun.*, 1998, 2319–2320; (b) M. Matsumoto, *J. Photochem. Photobiol. C*, 2004, **5**, 27–53.

5 A. J. Bloodworth and H. J. Eggelte, 'Endoperoxides', in *Singlet Oxygen*, ed. A. A. Frimer, Vol. **II**, CRC Press, Boca Raton, 1985, pp. 93–203.

6 (a) K. Gollnick and H. J. Kuhn, 'Ene-reactions with Singlet Oxygen', in *Singlet Oxygen*, ed. H. H. Wasserman and R. W. Murray, Academic Press, New York, 1979, pp. 287–427; (b) A. A. Frimer and L. M. Stephenson, 'The Singlet Oxygen "Ene" Reaction', in *Singlet Oxygen*, ed. A. A. Frimer, Vol. **II**, CRC Press, Boca Raton, 1985, pp. 67–92.

7 E. L. Clennan and C. S. J. Foote, 'Endoperoxides', in *Organic Peroxides*, ed. W. Ando, Wiley, New York, 1992, pp. 255–318.

8 N. J. Turro, M.-F. Chow and J. Rigaudy, *J. Am. Chem. Soc.*, 1977, **101**, 1300–1301.

9 N. J. Turro, M.-F. Chow and J. Rigaudy, *J. Am. Chem. Soc.*, 1979, **103**, 7218–7224.

10 For example, F. Wani, M. Endo and T. Fujioka, *Appl. Phys. Lett.*, 1999, **75**, 20, 3081–3083.

11 (a) I. Saito, T. Matsuura and K. Inoue, *J. Am. Chem. Soc.*, 1981, **103**, 188; (b) I. Saito, T. Matsuura and K. Inoue, *J. Am. Chem. Soc.*, 1983, **105**, 3200; (c) K. Inoue, T. Matsuura and I. Saito, *J. Photochem.*, 1984, **25**, 511.

12 I. Saito, R. Nagata and T. Matsuura, *J. Am. Chem. Soc.*, 1985, **107**, 6329–6334.