IN THIS ISSUE

ISSN 1359-7345 CODEN CHCOFS (7) 813-956 (2005)

In this issue...

Tracking cancer cells

The uptake and distribution of zinc bis(thiosemicarbazone) complexes in living cancer cells can be studied by fluorescence microscopy. See Dilworth *et al.*, pp. 845–847.





Cover

See Richard G. Compton, Craig E. Banks, Trevor J. Davies and Gregory G. Wildgoose, pp. 829–841. This feature article addresses the reactivity of carbon surfaces, with particular emphasis on the importance of edge-plane sites for both electron transfer and surface modification.

CHEMICAL TECHNOLOGY

T5

Chemical Technology highlights the latest applications and technological aspects of research across the chemical sciences.

Chemical Technology

February 2005/Volume 2/Issue 2

www.rsc.org/chemicaltechnology


FEATURE ARTICLE

829

Electrocatalysis at graphite and carbon nanotube modified electrodes: edge-plane sites and tube ends are the reactive sites

Craig E. Banks, Trevor J. Davies, Gregory G. Wildgoose and Richard G. Compton*

This feature article addresses the reactivity of carbon surfaces, with particular emphasis on the importance of edge-plane sites for both electron transfer and surface modification. Novel carbon nanotube and graphitic materials are contrasted and a critical overview of the recent literature is presented.



EDITORIAL STAFF

Managing editor

Sarah Thomas

Deputy editor

Sula Armstrong

Assistant editors

Meriel Dyche, Lorna Jack, Alison Stoddart, Katherine Vickers

Publishing assistants

Jayne Drake, Jayne Gough, Lois Kershaw,

Crystallographic data editor

Kirsty Anderson

Team leader, serials production

Technical editors

Celia Clarke, Sandra Jones, Caroline Moore, Elinor Richards, Michael Smith, Ken Wilkinson

Editorial secretaries

Sonya Spring, Julie Thompson, Rebecca Gotobed

Publisher, journals and reviews

Adrian Kybett

Chemical Communications (print: ISSN 1359-7345; electronic: ISSN1364-548X) is published 48 times a year by the Royal Society of Chemistry, Thomas Graham House, Science Park, Milton Road, Cambridge, UK CB4 0WF. All orders accompanied by payment should be sent directly to Portland Customer Services, Commerce Way, Colchester, Essex, UK CO2 8HP. Tel +44 (0)1206 226050; E-mail sales@rscdistribution.org

2005Annual (print + electronic) subscription price: £1595; US\$2635, 2005 Annual (electronic) subscription price: £1435; US\$2370. Customers in Canada will be subject to a surcharge to cover GST. Customers in the EU subscribing to the electronic version only will be charged VAT. If you take an institutional subscription to any RSC journal you are entitled to free, site-wide web access to that journal. You can arrange access via Internet Protocol (IP) address at www.rsc.org/ip. Customers should make payments by cheque in sterling payable on a UK clearing bank or in US dollars payable on a US clearing bank. Periodicals postage paid at Rahway, NJ, USA and at additional mailing offices. Airfreight and mailing in the USA by Mercury Airfreight International Ltd., 365 Blair Road, Avenel, NJ 07001, USA. US Postmaster: send address changes to Chemical Communications, c/o Mercury Airfreight International Ltd., 365 Blair Road, Avenel, NJ 07001. All despatches outside the UK by Consolidated Airfreight. PRINTED IN THE UK.

© The Royal Society of Chemistry, 2005. Apart from fair dealing for the purposes of research or private study for non-commercial purposes, or criticism or review, as permitted under the Copyright, Designs and Patents Act 1988 and the Copyright and Related Rights Regulations 2003, this publication may only be reproduced, stored or transmitted, in any form or by any means, with the prior permission in writing of the Publisher or in the case of reprographic reproduction in accordance with the terms of licences issued by the Copyright Licensing Agency in the UK. US copyright law is applicable to users in the USA. The Royal Society of Chemistry takes reasonable care in the preparation of this publication but does not accept liability for the consequences of any errors or omissions. Inclusion of an item in this publication does not imply endorsement by The Royal Society of Chemistry of the content of the original documents to which that item refers.

ChemComm

Chemical Communications

www.rsc.org/chemcomm

EDITORIAL BOARD

Chairman

Roeland J. M. Nolte, Nijmegen, The Netherlands nolte@sci.kun.nl

Jerry L. Atwood, Columbia, USA rsc.chemcomm@missouri.edu

Shankar Balasubramanian, Cambridge, UK sb10031@cam.ac.uk

Hans-Ulrich Blaser, Solvias AG, Switzerland hans-ulrich.blaser@SOLVIAS.com

P. Andrew Evans, Bloomington, USA chemcomm@indiana.edu

Makoto Fujita, Tokyo, Japan mfujita@appchem.t.u-tokyo.ac.jp

Alois Fürstner, Mülhelm, Germany fuerstner@mpi-muelheim.mpg.de

David Haddleton, Warwick, UK D.M.Haddleton@warwick.ac.uk

chemcomm@mit.edu Dermot O'Hare, Oxford, UK chemcomm@chem.ox.ac.uk Colin Raston, Perth, Australia clraston@chem.uwa.edu.au Ferdi Schüth, Mülheim, Germany schueth@mpi-muelheim.mpa.de T. Don Tilley Berkeley, USA chemcomm@berkeley.edu

Donald Hilvert, Zurich, Switzerland

Mir Wais Hosseini, Strasbourg, France

hosseini@chimie.u-strasbg.fr

Barbara Imperiali, Cambridge, USA

hilvert@org.chem.ethz.ch

Scientific editors

The Scientific Editors welcome enquiries from potential authors regarding the submission and scientific content of papers. For more information please see http://www.rsc.org/authors

Associate editors

Manuscripts can be submitted online at www.rsc.org/resource Manuscripts from the Americas should be submitted to the appropriate Associate Editor.

Supramolecular

Jerry L. Atwood

Organic

P. Andrew Evans

Editorial advisory board

Varinder Aggarwal, Bristol, UK Takuzo Aida, Tokyo, Japan Frank Allen, CCDC, Cambridge, UK Dario Braga, Bologna, Italy Jillian M. Buriak, Alberta, Canada Derrick Clive, Alberta, Canada Marcetta Darensbourg, College Station, USA Gregory C. Fu, Cambridge, USA Tohru Fukuyama, Tokyo, Japan Lutz Gade, Heidelberg, Germany Philip Gale, Southampton, UK George W. Gokel, St Louis, USA Craig Hawker, Santa Barbara, USA Andrew B. Holmes, Melbourne, Australia Amir Hoveyda, Boston, USA Kazuyuki Kuroda, Tokyo, Japan

Dermot O'Hare Donald Hilvert Mir Wais Hosseini Alois Fürstner

Chemical biology

Barbara Imperiali

Inorganic, Organometallic and Materials

T. Don Tilley

Manuscripts from other regions should be submitted to the Managing Editor Dr Sarah Thomas (chemcomm@rsc.org) For information on how to submit your manuscript see http://www.rsc.org/authors

Jérôme Lacour, Geneva, Switzerland David MacMillan, Pasadena, USA E. W. 'Bert' Meijer, Eindhoven, The Netherlands Jason Micklefield, Manchester, UK Achim Müller, Bielefeld, Germany Catherine Murphy, South Carolina, USA Atsuhiro Osuka, Kyoto, Japan Ian Paterson, Cambridge, UK Maurizio Prato, Trieste, Italy Christopher A. Reed, Riverside, USA Robin Rogers, Alabama, USA Michael Sailor, San Diego, USA Jonathan Sessler, Austin, USA Jonathan W. Steed, Durham, UK Carsten Tschierske, Halle, Germany Herbert Waldmann, Dortmund, Germany Henry N. C. Wong, Hong Kong, PR China

Advertisement sales: Tel +44 (0)1223 432243

Fax +44 (0)1223 426017; E-mail advertising@rsc.org

Royal Society of Chemistry: Registered Charity No. 207890.


Authors may reproduce/republish portions of their published contribution without seeking permission from the RSC, provided that any such republication is accompanied by an acknowledgement in the form: (Original Citation)-Reproduced by permission of The Royal Society of Chemistry.



Demonstration of the importance of oxygenated species at the ends of carbon nanotubes for their favourable electrochemical properties

Alison Chou, Till Böcking, Nagindar K. Singh and J. Justin Gooding*

Definitive evidence is presented for the favourable electrochemical properties of carbon nanotube modified electrodes arising from the ends of SWNTs due to oxygenated carbon species in general, and carboxylic acid moieties in particular, produced during acid purification.



Potential / mV versus Ag/AgCI


845



Fluorescence studies of the intra-cellular distribution of zinc bis(thiosemicarbazone) complexes in human cancer cells

Andrew R. Cowley, Jason Davis, Jonathan R. Dilworth,* Paul S. Donnelly, Rachel Dobson, Adrian Nightingale, Josephine M. Peach, Ben Shore, David Kerr and Len Seymour

The intrinsic fluorescence of zinc complexes of bis(thiosemicarbazone) ligands is used to track for the first time their distribution in a range of living human cancer cells; this is shown to be dependent on both cell type and ligand substituents.




848

Porous functionalised silica particles: a potential platform for biomolecular screening

Angus P. R. Johnston, Bronwyn J. Battersby, Gwendolyn A. Lawrie and Matt Trau*

Using a novel emulsion synthesis technique, porous, functionalized silica particles possessing controlled morphology have been developed. Fluorescent dyes can be covalently attached to the particles, which act like an optical barcode.




851



Hydrolysis of acetylcholinesterase inhibitors organophosphorus acid anhydrolase enzyme immobilization on photoluminescent porous silicon platforms

Sonia E. Létant, Staci R. Kane, Bradley R. Hart, Masood Z. Hadi, Tu-Chen Cheng, Vipin K. Rastogi and John G. Reynolds*

We report on the immobilization and consequent enzymatic behavior of an OPAA enzyme on luminescent porous silicon devices, and on the utilization of this new platform to hydrolyze *p*-nitrophenyl-soman.



COMMUNICATIONS

854


20 18 16 Concentration (mM) 14 12 10 8 6 4 2 0 0 200 800 1000 400 600 Time (s)

Opening the 'black box': oscillations in organocuprate conjugate addition reactions

Michael D. Murphy, Craig A. Ogle* and Steven H. Bertz*

Complex oscillations are observed in the reactions of butyl Gilman reagents with 2-cyclohexenone, a typical substrate.

857




Resorcinarene assemblies as synthetic receptors

Liam C. Palmer, Alexander Shivanyuk, Masamichi Yamanaka and Julius Rebek, Jr.

The host–guest complexes of resorcinarenes are re-examined in solution through modern spectroscopic methods. The assemblies are determined to be hexameric capsules and the guest exchange rates are measured by EXSY NMR spectroscopy. The neutral and cationic guest exchange rates correspond to barriers of approximately 20 kcal mol⁻¹.

859




High facial selectivity observed in amine coordination to chiral oxazaborolidinones

Toshiro Harada,* Yuuki Yamamoto and Takahiro Kusukawa

High top-face selectivity can be rationalized by an electrostatic stabilizing interaction between the negatively charged sulfonyl oxygen atom and the positively charged pyridyl moiety.


862



The antituberculosis, antitumor, multibranched dodecafuranoarabinan of *Mycobacterium* species has been assembled from a single n-pentenylfuranoside source


Jun Lu and Bert Fraser-Reid*

An n-pentenyl furanosyl-1,2-orthoester can function as a donor or be rearranged leading to an n-pentenyl furanoside acceptor which is glycosylated by its progenitor, regioselectively or doubly, thereby enabling rapid fabrication of a multibranched dodecasaccharide, known to possess a wide variety of biological interactions.



Shuhei Furukawa, Masaaki Ohba and Susumu Kitagawa*

A novel 2-D coordination polymer with honeycomb (6,3) net topology is constructed from a paramagnetic paddlewheel diruthenium tetrabenzoate, $[Ru^{II}_{2}(O_{2}CPh)_{4}]$, as a linear linker motif and triazine as a three-connected node.




868

1-Butyl-3-methylimidazolium 3,5-dinitro-1,2,4-triazolate: a novel ionic liquid containing a rigid, planar energetic anion

Alan R. Katritzky,* Shailendra Singh, Kostyantyn Kirichenko, John D. Holbrey,* Marcin Smiglak, W. Matthew Reichert and Robin D. Rogers*

The novel crystalline salt, 1-butyl-3-methylimidazolium 3,5-dinitro-1,2,4-triazolate, has been synthesized and characterized. The salt melts at 35 °C and demonstrates how ionic liquids can be readily prepared with functionalized heterocyclic energetic anions.




871

The first example of a liquid crystalline side-chain polymer with bent-core mesogenic units: ferroelectric switching and spontaneous achiral symmetry breaking in an achiral polymer

Christina Keith, R. Amaranatha Reddy and Carsten Tschierske*


The first bent-core side-chain polymeric liquid crystal, a dimethylsiloxane diluted polysiloxane, was synthesised and shows an optically isotropic ferroelectric switching polar smectic C phase consisting of a conglomerate of homogeneously chiral domains with opposite handedness.



Single-walled carbon nanotubes displaying multivalent ligands for capturing pathogens

Lingrong Gu, Tara Elkin, Xiuping Jiang, Huaping Li, Yi Lin, Liangwei Qu, Tzuen-Rong J. Tzeng, Ronalda Joseph and Ya-Ping Sun*

A single-walled carbon nanotube was exploited for its semi-flexible pseudo-one-dimensional nanostructure as a unique scaffold to display multivalent carbohydrate ligands, with a specific demonstration showing that galactosylated carbon nanotubes were effective in the capturing of pathogenic Escherichia coli in solution.



Native AMDase


Ar C'''/CO_2H CO_2H CO_2H

Inversion of enantioselectivity of asymmetric biocatalytic decarboxylation by site-directed mutagenesis based on the reaction mechanism

Yoichiro Ijima, Kaori Matoishi, Yosuke Terao, Nobuhide Doi, Hiroshi Yanagawa and Hiromichi Ohta*

Arylmalonate decarboxylase is an unique enzyme which catalyses the enantioselective decarboxylation of α -aryl- α -methylmalonate to give optically pure arylpropionate. The enantioselectivity of this enzyme was dramatically changed by the introduction of two mutations (G74C/C188S) based on the reaction mechanism.

880




Selective polymorph transformation *via* solvent-drop grinding

Andrew V. Trask, Ning Shan,* W. D. Samuel Motherwell, William Jones,* Shaohua Feng, Reginald B. H. Tan and Keith J. Carpenter

A method of inducing specific polymorph transformations is exemplified with two single-component systems, whereby a given crystal form undergoes conversion when subjected to solid state grinding in the presence of a minor quantity of a certain solvent.

883




Photoactivated luminescent CdSe quantum dots as sensitive cyanide probes in aqueous solutions

Wei Jun Jin, María T. Fernández-Argüelles, José M. Costa-Fernández, Rosario Pereiro and Alfredo Sanz-Medel*

The first application of quantum dots as highly sensitive and selective luminescent probes for anions determination in aqueous solutions is described. The synthesized nanoparticles proved to be one of the most sensitive detection systems available for monitoring the highly toxic cyanide anion.

886



Reversible Michael addition of thiols as a new tool for dynamic combinatorial chemistry


Baolu Shi and Michael F. Greaney*

Glutathione reacts with six ethacrynic acid derivatives to form a twelve membered dynamic combinatorial library of Michael adducts and enone starting materials. The library reaches equilibrium in three hours, is freely reversible and can be switched on or off by a simple pH change.

Dye-sensitized solar cells based on composite solid polymer electrolytes

Moon-Sung Kang, Jong Hak Kim, Young Jin Kim, Jongok Won, Nam-Gyu Park and Yong Soo Kang*

Dye-sensitized solar cells employing composite polymer electrolytes show a remarkable energy conversion efficiency of $\sim 3.84\%$ at $100~\text{mW}~\text{cm}^{-2}$. The increased efficiency is primarily due to the addition of small-sized oligomeric poly(propylene glycol) into high molecular weight PEO with electrolytes of KI and I_2 for both enhancing the ionic conductivity and more importantly enlarging the interfacial contact area between dyes and polymer electrolytes.




892

Solvent-free, direct synthesis of supramolecular nano-capsules

Jochen Antesberger, Gareth W. V. Cave, Matthew C. Ferrarelli, Michael W. Heaven, Colin L. Raston and Jerry L. Atwood*

The single step direct synthesis of pyrogallol[4]arene *via* a solvent free protocol yields the pure product as a self-assembled nano-capsule, comprised of six macrocyclic building blocks.




895

Molecular recognition in a uradinyl-functionalized stable radical

Patrick Taylor, Paul M. Lahti,* Joseph B. Carroll and Vincent M. Rotello

2-(6-Uradinyl)-4,4,5,5-tetramethyl-4,5-dihydro-1H-imidazole-l-oxyl (1) binds to hydrogen-bonding complement 2,6-di(propylamido)pyridine (DAP) in chloroform with association constant $K_a = 220 \text{ M}^{-1}$ at 33 °C; ESI-MS shows not only 1:DAP complementary dyad formation, but also 1:(DAP)₂ formation at higher relative concentrations of DAP.




898

Ni(II)1,4,8,11-tetraazacyclotetradecane electrocatalytic films prepared on top of surface anchored PAMAM dendrimer layers. A new type of electrocatalytic material for the electrochemical oxidation of methanol

Miguel A. González-Fuentes, J. Manríquez, S. Gutiérrez-Granados, A. Alatorre-Ordaz and Luis A. Godínez*

The presence of interfacially anchored PAMAM dendrimers on a Au modified electrode, enhances the electrocatalytic activity of Ni–cyclam towards the electrochemical electroxidation of methanol.



$$R^{1} R^{2} + R^{3} R^{2} R^{4} R^{3} R^{2}$$

$$R^{4} R^{3} R^{2} R^{3} R^{3} R^{2}$$

$$R^{2} R^{3} R^{2}$$

Low-valent Nb(III)-mediated synthesis of 1,1,2-trisubstituted-1*H*-indenes from aliphatic ketones and aryl-substituted alkynes

Yasushi Obora, Masahiro Kimura, Makoto Tokunaga and Yasushi Tsuji*

A variety of 1,1,2-trisubstituted-1*H*-indenes are synthesized by the reaction of aliphatic ketones, aryl-substituted alkynes and NbCl₃(DME) in 1,2-dichloroethane under reflux conditions.

903

Chloroindate(III) ionic liquids: recyclable media for Friedel-Crafts acylation reactions

Martyn J. Earle, Ullastiina Hakala, Christopher Hardacre, Johanna Karkkainen, Barry J. McAuley, David W. Rooney, Kenneth R. Seddon, Jillian M. Thompson and Kristina Wähälä

Chloroindate(III) ionic liquids are versatile reaction media for Friedel-Crafts acylation reactions; the system is catalytic and totally recyclable.

906



O
$$AcO_2$$
 $BocN$ MeO_2C $BocN$ MeO_2C N

A general method for making bicyclic compounds with nitrogen at a bridgehead—application to the halichlorine spiro subunit

Derrick L. J. Clive,* Maolin Yu and Zhiyong Li

N-Protected β-amino aldehydes having the nitrogen in a ring are easily converted into Morita–Baylis–Hillman adducts; O-acetylation and N-deprotection result in spontaneous cyclization to bicyclic structures having nitrogen at a bridgehead.

909



NaI-catalyzed highly regioselective ring-opening [1 + 2] cycloaddition reaction of cyclopropenes with imines: highly stereoselective synthesis of *cis*-vinylic aziridines

Shengming Ma,* Junliang Zhang, Lianghua Lu, Xin Jin, Yangjun Cai and Hairong Hou

cis-Vinylic aziridines were prepared highly stereoselectively via a NaI-catalyzed regioselective ring-opening [1 + 2] cycloaddition reaction of cyclopropenes with imines.



Regioselective silylphosphination of methyl vinyl ketone with complexes containing cyclic and linear iron-silicon-phosphorus reacting sites

Masaaki Okazaki,* Kyeong A Jung and Hiromi Tobita*


Methyl vinyl ketone was incorporated into iron-siliconphosphorus reacting sites in three different fashions. Structures of the products were fully characterized by X-ray diffraction

915

Determination of the ¹³C NMR chemical shifts in an a_{2n} type iron(III) porphyrin cation radical

Akito Hoshino and Mikio Nakamura*

The chemical shifts of all the carbons in an a_{2u} type iron(III) porphyrin radical cation have been determined by the titration method.




918



One-step solution-based catalytic route to fabricate novel α-MnO₂ hierarchical structures on a large scale

Zhengquan Li, Yue Ding, Yujie Xiong, Qing Yang and Yi Xie*

A novel α-MnO₂ hierarchical structure, a core-shell structure with spherically aligned nanorods, was conveniently synthesized by a room-temperature solution-based catalytic route.




921



The use of tin-decorated mesoporous carbon as an anode material for rechargeable lithium batteries

I. Grigoriants, L. Sominski, Hongliang Li, Ilan Ifargan, D. Aurbach and A. Gedanken*

We report on the preparation of mesoporous carbon particles, decorated with metallic tin nanoparticles. These composites were proved to be a promising active mass for anodes in rechargeable Li batteries.



The role of carbon dioxide in chemoselective hydrogenation of halonitroaromatics over supported noble metal catalysts in supercritical carbon dioxide

Shinichiro Ichikawa, Mizuki Tada, Yasuhiro Iwasawa and Takao Ikariya*

Chemoselective hydrogenation of halogenated nitrobenzenes over Pt/C catalysts proceeds effectively in supercritical carbon dioxide (scCO₂) to produce halogenated anilines with an excellent selectivity in which the rate of the hydrogenation is markedly enhanced and the dehalogenation reaction is significantly suppressed.

927

Lanthanum-exchanged zeolites as active and selective catalysts for the generation of singlet oxygen from hydrogen peroxide

Joos Wahlen, Dirk De Vos, Sigrid De Hertogh, Véronique Nardello, Jean-Marie Aubry, Paul Alsters and Pierre Jacobs*

Lanthanum(III)-exchanged zeolites Beta and USY are active and selective catalysts for the generation of singlet oxygen from H₂O₂ showing superior activity and oxidant efficiency compared to unsupported La-catalysts, *e.g.* La(OH)₃.

930



A remarkable side-product from the synthesis of an octaalkylphthalocyanine: formation of a tetrabenzotriazaporphyrin

Andrew N. Cammidge,* Michael J. Cook,* David L. Hughes, Fabien Nekelson and Muhibur Rahman

Subtle changes to the experimental protocol of the lithium alkyloxide induced conversion of 3,6-dihexylphthalonitrile into 1,4,8,11,15,18,22,25-octahexylphthalocyanine can suppress or enhance the unexpected formation of the correspondingly substituted tetrabenzotriazaporphyrin side-product.

933



2-Arylallyl as a new protecting group for amines, amides and alcohols

José Barluenga,* Francisco J. Fañanás, Roberto Sanz, César Marcos and José M. Ignacio

A new protecting allyl-type group, 2-arylallyl, for amines, amides and alcohols is reported. Selective deprotection with *t*-BuLi probably involves an intermolecular carbolithiation.

Fine-tuning furan toxicity: fast and quantitative DNA interchain cross-link formation upon selective oxidation of a furan containing oligonucleotide

Sami Halila, Trinidad Velasco, Pierre De Clercq and Annemieke Madder*


In situ oxidation of a duplex formed between a furan modified oligonucleotide and its complement transforms the furan moiety into a reactive enal that triggers fast and clean interchain cross-link formation.

939

N-Alkyl-N-methylpyrrolidinium salts as templates for hexagonally meso-ordered silicate thin films

Andrew M. Dattelbaum, Sheila N. Baker and Gary A. Baker*

A series of *N*-alkyl-*N*-methylpyrrolidinium bromide salts (C14, C16, or C18) have been explored as templates for the synthesis of ordered, mesoporous silica films. "Soft" annealing based on deep-UV calcination is also reported.



942

Low-dimensional chainlike assemblies of TiO₂ nanorod-stabilized Au nanoparticles

P. D. Cozzoli,* E. Fanizza, M. L. Curri, D. Laub and A. Agostiano

Chainlike Au nanoparticle assemblies are formed upon UV-irradiating $HAuCl_4$ in the presence of TiO_2 nanorods as stabilizers in apolar media.

TiO₂ nanorods


HAuCl₄

945

A ditopic fluorescent sensor for potassium fluoride

Suvi J. M. Koskela, Thomas M. Fyles and Tony D. James*

Ditopic fluorescent sensors for potassium fluoride have been prepared using a boronic acid receptor (fluoride) and crown ether receptors (potassium)




Alkali metal cation intercalation properties of titanate nanotubes

Renzhi Ma,* Takayoshi Sasaki and Yoshio Bando

Hydrothermally synthesized titanate nanotubes show high alkali metal cation intercalation reactivity evidenced by high resolution microscopic examination, elemental mapping and profiling techniques as well as wet chemical analysis.

951



New reactions of $\beta\text{-diketiminatol}$ anthanoid complexes: sterically induced self-deprotonation of $\beta\text{-diketiminatol}$ ligands

Peter B. Hitchcock, Michael F. Lappert* and Andrey V. Protchenko

Attempted synthesis of sterically demanding bis- or tris- β -diketiminato complexes of lanthanoids resulted in ligand deprotonation and the formation of complexes containing both a "normal" and a deprotonated ligand; one of these on protonation gave the first cationic β -diketiminato—Ln complex.

AUTHOR INDEX

Agostiano, A., 942 Alatorre-Ordaz, A., 898 Alsters, Paul, 927 Antesberger, Jochen, 892 Atwood, Jerry L., 892 Aubry, Jean-Marie, 927 Aurbach, D., 921 Baker, Gary A., 939 Baker, Sheila N., 939 Bando, Yoshio, 948 Banks, Craig E., 829 Barluenga, José, 933 Battersby, Bronwyn J., 848 Bertz, Steven H., 854 Böcking, Till, 842 Cai, Yangjun, 909 Cammidge, Andrew N., 930 Carpenter, Keith J., 880 Carroll, Joseph B., 895 Cave, Gareth W. V., 892 Cheng, Tu-Chen, 851 Chou, Alison, 842 Clive, Derrick L. J., 906 Compton, Richard G., 829 Cook, Michael J., 930 Costa-Fernández, José M., 883 Cowley, Andrew R., 845 Cozzoli, P. D., 942 Curri, M. L., 942 Dattelbaum, Andrew M., 939 Davies, Trevor J., 829 Davis, Jason, 845 De Clercq, Pierre, 936 De Hertogh, Sigrid, 927 De Vos, Dirk, 927 Dilworth, Jonathan R., 845 Ding, Yue, 918 Dobson, Rachel, 845 Doi, Nobuhide, 877 Donnelly, Paul S., 845 Earle, Martyn J., 903 Elkin, Tara, 874 Fañanás, Francisco J., 933 Fanizza, E., 942 Feng, Shaohua, 880 Fernández-Argüelles, María T., Ferrarelli, Matthew C., 892 Fraser-Reid, Bert, 862 Furukawa, Shuhei, 865 Fyles, Thomas M., 945 Gedanken, A., 921 Godínez, Luis A., 898 González-Fuentes, Miguel A., Gooding, J. Justin, 842 Greaney, Michael F., 886 Grigoriants, I., 921 Gu, Lingrong, 874 Gutiérrez-Granados, S., 898 Hadi, Masood Z., 851 Hakala, Ullastiina, 903 Halila, Sami, 936 Harada, Toshiro, 859 Hardacre, Christopher, 903 Hart, Bradley R., 851 Heaven, Michael W., 892 Hitchcock, Peter B., 951 Holbrey, John D., 868 Hoshino, Akito, 915 Hou, Hairong, 909 Hughes, David L., 930 Ichikawa, Shinichiro, 924 Ifargan, Ilan, 921 Ignacio, José M., 933 Ijima, Yoichiro, 877 Ikariya, Takao, 924 Iwasawa, Yasuhiro, 924 Jacobs, Pierre, 927 James, Tony D., 945 Jiang, Xiuping, 874 Jin, Wei Jun, 883 Jin, Xin, 909 Johnston, Angus P. R., 848 Jones, William, 880 Joseph, Ronalda, 874 Jung, Kyeong A, 912 Kane, Staci R., 851 Kang, Moon-Sung, 889 Kang, Yong Soo, 889 Karkkainen, Johanna, 903 Katritzky, Alan R., 868 Keith, Christina, 871 Kerr, David, 845

Kim, Jong Hak, 889 Kim, Young Jin, 889 Kimura, Masahiro, 901 Kirichenko, Kostyantyn, 868 Kitagawa, Susumu, 865 Koskela, Suvi J. M., 945 Kusukawa, Takahiro, 859 Lahti, Paul M., 895 Lappert, Michael F., 951 Laub, D., 942 Lawrie, Gwendolyn A., 848 Létant, Sonia E., 851 Li, Hongliang, 921 Li, Huaping, 874 Li, Zhengquan, 918 Li, Zhiyong, 906 Lin, Yi, 874 Lu, Jun, 862 Lu, Lianghua, 909 Ma, Renzhi, 948 Ma, Shengming, 909 Madder, Annemieke, 936 Manríquez, J., 898 Marcos, César, 933 Matoishi, Kaori, 877 McAuley, Barry J., 903 Motherwell, W. D. Samuel, 880 Murphy, Michael D., 854 Nakamura, Mikio, 915 Nardello, Véronique, 927 Nekelson, Fabien, 930 Nightingale, Adrian, 845 Obora, Yasushi, 901 Ogle, Craig A., 854 Ohba, Masaaki, 865 Ohta, Hiromichi, 877 Okazaki, Masaaki, 912 Palmer, Liam C., 857 Park, Nam-Gyu, 889 Peach, Josephine M., 845 Pereiro, Rosario, 883 Protchenko, Andrey V., 951 Qu, Liangwei, 874 Rahman, Muhibur, 930 Rastogi, Vipin K., 851 Raston, Colin L., 892 Rebek, Jr., Julius, 857

Reddy, R. Amaranatha, 871 Reichert, W. Matthew, 868 Reynolds, John G., 851 Rogers, Robin D., 868 Rooney, David W., 903 Rotello, Vincent M., 895 Sanz, Roberto, 933 Sanz-Medel, Alfredo, 883 Sasaki, Takayoshi, 948 Seddon, Kenneth R., 903 Seymour, Len, 845 Shan, Ning, 880 Shi, Baolu, 886 Shivanyuk, Alexander, 857 Shore, Ben, 845 Singh, Nagindar K., 842 Singh, Shailendra, 868 Smiglak, Marcin, 868 Sominski, L., 921 Sun, Ya-Ping, 874 Tada, Mizuki, 924 Tan, Reginald B. H., 880 Taylor, Patrick, 895 Terao, Yosuke, 877 Thompson, Jillian M., 903 Tobita, Hiromi, 912 Tokunaga, Makoto, 901 Trask, Andrew V., 880 Trau, Matt, 848 Tschierske, Carsten, 871 Tsuji, Yasushi, 901 Tzeng, Tzuen-Rong J., 874 Velasco, Trinidad, 936 Wähälä, Kristina, 903 Wahlen, Joos, 927 Wildgoose, Gregory G., 829 Won, Jongok, 889 Xie, Yi, 918 Xiong, Yujie, 918 Yamamoto, Yuuki, 859 Yamanaka, Masamichi, 857 Yanagawa, Hiroshi, 877 Yang, Qing, 918 Yu, Maolin, 906 Zhang, Junliang, 909

FREE E-MAIL ALERTS

Contents lists in advance of publication are available on the web via www.rsc.org/chemcomm - or take advantage of our free e-mail alerting service (www.rsc.org/ej_alert) to receive notification each time a new list becomes available.

* Indicates the author for correspondence: see article for details.

Electronic supplementary information (ESI) is available via the online article (see http://www.rsc.org/esi for general information about ESI).

ADVANCE ARTICLES AND ELECTRONIC JOURNAL

Free site-wide access to Advance Articles and electronic form of this journal is provided with a full-rate institutional subscription. See www.rsc.org/ejs for more information.

An international, high quality journal covering the full breadth of synthetic, physical and biomolecular organic chemistry.

See for yourself - examples of forthcoming papers are listed below

For further details and free access to Issue 1, visit

www.rsc.org/obc

Emerging Area:

Asymmetric Organocatalysis

Benjamin List, Germany

Communications:

Synthesis and properties of 1,6,7,12,13,18,19,24-octamethyl-acenaphthyleno [b,l]tetraphenylene

Jay S. Siegel, Switzerland

Fluorescence enhancement of bis-acridine orange peptide, BAO, upon binding to double stranded DNA

Shigeori Takenaka, Japan

The Selective Aerobic Oxidation of Methylaromatics to Benzaldehydes using a unique combination of two Heterogeneous Catalysts

James H Clark, UK

Articles:

Synthesis and Photochemical Properties of Photoactivated
Antitumor Prodrugs Releasing 5-Fluorouracil

Zhouen Zhang, Japan

Neutral cumulene oxide CCCCO is accessible by one-electron oxidation of [CCCCO]-. in the gas phase

John H. Bowie, Australia

Preparation and unique circular dichroism phenomena of urea-functionalized self-folding resorcinarenes bearing chiral termini through asymmetric hydrogen bonding belts

Osamu Hayashida, Japan

Involvement of Proton Transfer in the Reductive
Repair of DNA Guanyl Radicals by Aniline Derivatives

Jamie R. Milligan, USA

Thermotropic liquid-crystalline peptide derivatives: oligo(glutamic acid)s forming hydrogen-bonded columns

Takashi Kato, Japan

Structure-activity relationships of galabioside derivatives as inhibitors of E. coli and S. suis adhesins: Nanomolar inhibitors of S. suis adhesins

Ulf J. Nilsson, Sweden