

Room temperature palladium catalysed coupling of acyl chlorides with terminal alkynes†

Russell J. Cox,^{*a} Dougal J. Ritson,^a Thomas A. Dane,^a John Berge,^b Jonathan P. H. Charmant^c and Anob Kantacha^c

Received (in Cambridge, UK) 27th September 2004, Accepted 29th October 2004


First published as an Advance Article on the web 10th January 2005

DOI: 10.1039/b414826f

Conditions are reported for the facile, high-yielding coupling of acyl chlorides with terminal alkynes in a reaction involving palladium and copper iodide; the reaction is tolerant of a wide variety of acyl chlorides and terminal alkynes and provides a convenient one-pot route to acetylenic ketones.

Acetylenic ketones are useful precursors for organic synthesis, especially as precursors during the synthesis of saturated polycyclic¹ and heterocyclic² compounds. However their synthesis is not straightforward because the addition of acetylenic nucleophiles to acyl chlorides or acid anhydrides can be accompanied by unwanted addition to the products, forming bis-acetylenic tertiary alcohols.³ An alternative route involving addition of acetylenic Grignard reagents, for example, to aldehydes can be low-yielding and requires a subsequent oxidation reaction.^{2,4}

We reasoned that because acyl chlorides are known to oxidatively add to Pd(0) species, forming an acyl Pd(II) intermediate,⁵ it should be possible to intercept this species with a suitable nucleophile (Scheme 1). For example, others have reported the use of boronic acids as nucleophiles in similar processes.⁶ In the classical Sonogashira reaction, terminal alkynes are activated as nucleophiles under basic conditions using a CuI co-catalyst. The precise identity of the Pd catalyst active in these reactions is not known, but could involve discrete Pd(0)L₂ species or Pd nanoclusters as has recently been demonstrated for Heck reactions.⁷ Sonogashira also reported that terminal acetylenes would react with acid chlorides when catalysed by Pd(0) and CuI, but the reactions were limited by unwanted reaction between the acid chlorides and the Et₃N solvent, reducing yields.⁸ More recently Najera and co-workers have reported a similar reaction, but at high temperature and in the absence of Cu, giving moderate isolated yields of products.⁹ We now report a simple procedure which operates at RT giving high isolated yields of products.



Scheme 1 Catalytic cycle for the Pd(0) catalysed coupling of acid chlorides.

† Electronic Supplementary Information (ESI) available: experimental procedures, characterisation for compounds 5–28 and crystal structure data for 14. See <http://www.rsc.org/suppdata/cc/b4/b414826f/>

*r.j.cox@bris.ac.uk

In order to reduce the chance of reaction between the acid chloride component and the base, we reduced the base concentration and used 1.25 eq. of NEt₃ in THF rather than neat base as the solvent. Using these conditions with PdCl₂(PPh₃)₂ (0.9 mol%) and CuI (3.0 mol%) in dry THF at RT, the reaction between phenylacetylene (1.0 eq.) and benzoyl chloride (1.5 eq.) was complete by TLC analysis within 10 minutes (Table 1) (See ESI for experimental procedures).† The product 5 was isolated in 96% yield after purification.

The reaction is tolerant of a wide range of functional groups on the acetylene (Table 1), in particular alkyl, ester, silyl, silyloxy and protected amino groups are compatible, all giving isolated purified products in high yields. We also tested a range of aromatic acyl chlorides (Table 2, entries 1–4). Once again, good to high yields of purified products were obtained rapidly. *p*-Nitrobenzoyl chloride yielded the highly crystalline 14 (Fig. 1).‡

In the case of the *p*-bromobenzoyl chloride, the expected product 15 was obtained, and no reaction of the bromide was observed. However, in the case of the *p*-iodobenzoyl chloride, the acyl chloride and iodo groups appeared to be equally reactive, and in the presence of two equivalents of acetylene, the bis-acetylene 16 was formed in high yield.





Table 1 Coupling reactions of benzoyl chloride



Entry	R	Product	Yield (%) ^a
1	C ₆ H ₅	5	96
2	(CH ₂) ₂ CO ₂ Me	6	89
3	(CH ₂) ₂ CO ₂ ^t Bu	7	84
4	(CH ₂) ₃ CH ₃	8	88
5	Si(CH ₃) ₂ ^t Bu	9	96
6	CH ₂ OSi(CH ₃) ₂ ^t Bu	10	94
7	CH ₂ NH(BOC) ^b	11	91
8	CH ₂ NHAc	12	70

^a Isolated purified yield. ^b 1 h reaction.

Table 2 Coupling reactions of phenylacetylene


Entry	R	Product	Yield (%) ^a
1	<i>p</i> MeOC ₆ H ₄	13	98
2	<i>p</i> O ₂ NC ₆ H ₄	14	50 ^b
3	<i>p</i> BrC ₆ H ₄	15	89
4	<i>p</i> IC ₆ H ₄	16	66 ^c
5	^t Bu ^c	17	86
6	C ₆ H ₁₁	18	96
7	(CH ₃) ₂ CH	19	97
8	CH ₃ (CH ₂) ₂	21/22	43 ^d



^a Isolated purified yield. ^b Recrystallised yield. ^c overnight reaction. ^d using DIPEA as the base. ^e using two equivalents of acetylene.

**Fig. 1** X-ray crystal structure of *p*-nitrophenyl ketone **14**.[‡]**Scheme 2** Reaction of linear aliphatic acid chlorides.

Pivaloyl chloride, cyclohexylcarboxylic acid chloride and 2-methylpropionyl chloride also reacted smoothly to give high yields of the acetylenic ketones **17**, **18** and **19**. However, in the case of butyryl chloride, oxetane **20** was obtained in 46% isolated yield (Scheme 2). This presumably results from [2 + 2] cycloaddition reactions of ketenes. Such compounds are known to arise from the direct reaction of NEt₃ with aliphatic acyl chlorides,¹⁰ and in control reactions, using our standard reaction conditions but lacking the Pd catalyst, we observed the formation of **20** by TLC and ¹H NMR.

However, in the presence of the more hindered base diisopropylethylamine (DIPEA, Hunig's base) the reaction of butyryl chloride was more productive, giving a mixture of the desired product **21**, together with both isomers of the corresponding enol **22** (Scheme 3) in 43% combined yield.

In an effort to broaden the range of potential acyl partners, we used isobutyryl chloroformate **23** in an attempt to synthesise acetylenic esters (Scheme 4). In the presence of Et₃N, the only product was the corresponding

**Scheme 3** Reaction of linear aliphatic acyl chlorides.**Scheme 4** Low-yielding coupling reactions. *Reagents and conditions:* (i) HCCPh, PdCl₂(PPh₃)₂, CuI, Et₃N or DIPEA, THF, RT, 8%; (ii) HCC(CH₂)₂CO₂^tBu, PdCl₂(PPh₃)₂, CuI, DIPEA, THF, RT, 4%.

O-isobutyryl-*N,N*-diethylcarbamate **24a** formed by attack of the base on the carbonyl. In the presence of DIPEA the desired product **25** was formed but in 8% yield: the balance of the reaction was the corresponding carbamate rotamers **24b** and **c**. Similar results were obtained when using methyl oxalylchloride **26**. Using DIPEA as the base, this reacted to give a mixture of the desired product **27** (4%), a decarbonylated product **28** (4%) and the balance of the reaction as amide rotamers **29a** and **b**. These reactions, however, did indicate the intermediacy of an acyl-palladium species. Oxalyl-palladium species are known to easily decarbonylate,¹¹ and decarbonylation and interception by the amine¹² clearly compete with transfer of the acetylene in these cases.

Thus the reaction of terminal acetylenes with acid chlorides under room temperature Pd(0) catalysed conditions is a rapid and high-yielding method for the construction of synthetically useful acetylenic ketones. The product yields are significantly improved over previous reports⁸ and a wide variety of substituents are tolerated on the acetylene. Aromatic and α -branched aliphatic acid chlorides bearing a variety of substituents can be used. Linear aliphatic acid chlorides also react in the presence of DIPEA. The reaction conditions are mild enough that aromatic bromo-substituents, which could be used for subsequent organo-metallic coupling reactions, do not react but aromatic iodides are reactive under these conditions.

We thank the School of Chemistry and University of Bristol for funding. TAD thanks EPSRC for an earmarked quota studentship (00316697). DR thanks GlaxoSmithKline and BBSRC for a CASE award (01/A2/B/07049).

Russell J. Cox,^{*a} Dougal J. Ritson,^a Thomas A. Dane,^a John Berge,^b Jonathan P. H. Charmant^c and Anob Kantacha^c

^aSchool of Chemistry, University of Bristol, Cantock's Close, Bristol, UK BS8 1TS. E-mail: r.j.cox@bris.ac.uk; Fax: +44 (0)117 9298611

^bGlaxoSmithKline, New Frontiers Science Park (North), 3rd Avenue, Harlow, Essex, UK CM19 5AW

^cStructural Chemistry Laboratory, School of Chemistry, University of Bristol, Cantock's Close, Bristol, UK BS8 1TS

Notes and references

‡ CCDC 252032. See <http://www.rsc.org/suppdata/cc/b4/b414826f/> for crystallographic data in .cif or other electronic format. Crystal data for compound **14**. C₁₅H₉NO₃, M = 251.23, monoclinic, $a = 7.6488(6)$, $b = 7.1371(6)$, $c = 10.9816(9)$ Å, $\beta = 97.617(2)^\circ$, $U = 594.20(8)$ Å³, $T = 173(2)$ K, space group $P2_1$ (no. 4), $Z = 2$, $d_f(\text{Mo-K}\alpha) = 0.099$ mm⁻¹, 3914 reflections measured, 2516 unique ($R_{\text{int}} = 0.0185$) which were used in all calculations. The final R_1 and wR_2 were 0.0310 and 0.0796 ($I > 2\sigma I$).

1 M. E. Krafft, L. V. R. Bonaga, A. S. Felts, C. Hirosawa and S. Kerrigan, *J. Org. Chem.*, 2003, **68**, 6039–6042; J.-C. Wang, S.-S. Ng and M. J. Krische, *J. Am. Chem. Soc.*, 2003, **125**, 3682–3683.

- 2 P. Wipf, Y. Aoyama and T. E. Benedum, *Org. Lett.*, 2004, **6**, 3593–3595.
- 3 M. Kunishima, D. Nakata, S. Tanaka, K. Hioki and S. Tani, *Tetrahedron*, 2000, **56**, 9927–9935; M. J. Piggott and D. Wege, *Aust. J. Chem.*, 2000, **53**, 749–754.
- 4 D. Rodriguez, M. F. Martinez-Esperon, L. Castedo, D. Dominguez and C. Saa, *Synlett*, 2003, **10**, 1524–1526.
- 5 P. Fitton, M. P. Johnson and J. E. McKeon, *J. Chem. Soc., Chem. Commun.*, 1968, 6.
- 6 G. W. Kabalka, R. R. Malladi, D. Tejedor and S. Kelly, *Tetrahedron Lett.*, 2000, **41**, 999–1001; H. Chen and M.-Z. Deng, *Org. Lett.*, 2000, **2**, 1649–1651.
- 7 M. R. Eberhard, *Org. Lett.*, 2004, **6**, 2125–2128.
- 8 Y. Tohda, K. Sonogashira and N. Hagihara, *Synthesis*, 1977, **11**, 777–778.
- 9 D. A. Alonso, C. Najera and M. C. Pacheco, *J. Org. Chem.*, 2004, **69**, 1615–1619.
- 10 L. Hintermann and A. Togni, *Helv. Chim. Acta*, 2000, **83**, 2425–2435; R. W. Holder, H. S. Freiman and M. F. Stefanchick, *J. Org. Chem.*, 1976, **41**, 20, 3303–3307; C. D. Hurd and C. A. Blanchard, *J. Am. Chem. Soc.*, 1950, **72**, 1461–1462.
- 11 E. D. Dobrzynski and R. E. Angelici, *Inorg. Chem.*, 1975, **14**, 59–63.
- 12 The presence of PdCl₂(PPh₃)₂ in these reactions appeared to increase the rate of formation, and the yield, of amide and carbamate products.