

Indium–bipyridine-catalyzed, enantioselective thiolysis of *meso*-epoxides†

Mecheril Valsan Nandakumar,^a Andreas Tschöp,^a Harald Krautscheid^b and Christoph Schneider^{*a}

Received (in Cambridge, UK) 12th March 2007, Accepted 3rd April 2007

First published as an Advance Article on the web 24th April 2007

DOI: 10.1039/b703625f

The indium–bipyridine-catalyzed, enantioselective ring-opening of *meso*-epoxides with aliphatic and aromatic thiols furnished 1,2-mercapto alcohols in good yields and excellent enantioselectivities; the crystal structure of the chiral catalyst reveals a pentagonal-bipyramidal coordination geometry around the indium center.

The catalytic asymmetric ring-opening of *meso*-epoxides has proven to be a valuable tool for the straightforward synthesis of enantiomerically highly enriched 1,2-difunctionalized fine chemicals.¹ In particular, chiral 1,2-amino alcohols² and 1,2-azido alcohols,³ 1,2-diol derivatives,⁴ 1,2-cyano alcohols⁵ and 1,2-halohydrins⁶ have become available in excellent optical purities using this strategy. Among the processes reported for the addition of thiols to *meso*-epoxides,⁷ a heterobimetallic gallium–lithium–BINOL complex has been established by Shibasaki *et al.* as the most enantioselective catalyst currently known for this reaction.^{7a} Unfortunately, the use of *tert*-butyl thiol is mandatory here to avoid undesired ligand exchange on the metal center which significantly limits the applicability of this process.

We have reported recently a highly enantioselective catalyst for the alcoholysis and aminolysis of *meso*-epoxides composed of scandium triflate and bipyridine **1a** (Fig. 1).^{8a,b} With just 5 mol% of the metal–ligand complex formed *in situ*, *meso*-epoxides were ring-opened with anilines and aliphatic alcohols to furnish 1,2-amino alcohols^{8c} and 1,2-diol monoethers,^{8d} respectively, in good yields and up to 97% ee. We have now extended this protocol to the enantioselective thiolysis of *meso*-epoxides and show here that an indium–bipyridine complex formed *in situ* from InBr₃ and bipyridine **1a** is an effective chiral catalyst for this endeavour.^{9,10}

We started our investigations with the reaction of *cis*-stilbene oxide (**2a**) and benzylthiol (**3a**) and screened a broad range of




Fig. 1 **1a** R = H, **1b** R = CH₃.

^aInstitut für Organische Chemie, Universität Leipzig, Johannisallee 29, 04103 Leipzig, Germany. E-mail: schneider@chemie.uni-leipzig.de; Fax: +49 341 9736599; Tel: +49 341 9736529

^bInstitut für Anorganische Chemie, Universität Leipzig, Johannisallee 29, 04103 Leipzig, Germany. E-mail: krautscheid@rz.uni-leipzig.de; Fax: +49 341 36199; Tel: +49 341 9736172

† Electronic supplementary information (ESI) available: Experimental procedures, HPLC data, and spectral data for all new compounds. See DOI: 10.1039/b703625f

Lewis acids and Lewis acid–chiral ligand combinations as catalysts for this reaction (Table 1). Eventually, we identified indium(III) halide–bipyridine **1a** complexes as the most suitable chiral catalysts which not only possessed sufficient Lewis acidity to activate the epoxide towards nucleophilic ring-opening but also readily tolerated the Lewis basic thiol. Thus, 10 mol% of InCl₃ and 11 mol% of bipyridine **1a** afforded 1,2-mercapto alcohol **4a** in 60% yield and 95% ee after 6 days at rt (entry 4). The reaction was markedly accelerated by employing 10 mol% of InBr₃ and 11 mol% bipyridine **1a**, which furnished **4a** in 80% yield and 92% ee after just 16 h at rt (entry 5). Interestingly, no product formation was observed in the absence of the chiral ligand or when the unsubstituted 2,2'-bipyridine, which lacks the additional hydroxyl groups, was employed as the ligand (entries 1–3). Also, the bis-*O*-methylated bipyridine **1b** failed to give a reactive indium(III) catalyst (entry 6). These results clearly indicate the combined effect of the metal center and the hydroxyl protons for the epoxide opening reaction, similar to what we had observed in the scandium–bipyridine-catalyzed alcoholysis and aminolysis of *meso*-epoxides.^{8a}

Employing 10 mol% of InBr₃–bipyridine **1a** as catalyst, *cis*-stilbene oxide (**2a**) was ring-opened with various aromatic as well as aliphatic thiols, furnishing 1,2-mercapto alcohols **4a–f** in typically good to very good yields and excellent enantioselectivities of up to 96% ee (Table 2, entries 1–6). There appears to be no structural or electronic limitation in the thiol component and straight-chain aliphatic thiols even give rise to some of the most enantioselective reactions included in this study (entries 4–6). Other aromatic *meso*-epoxides were subsequently ring-opened with thiophenol and benzylthiol, and furnished the corresponding 1,2-mercapto alcohols **4g–m** in generally good yields and high

Table 1 Optimization studies

Entry	Lewis acid	Ligand	Time	Yield (%) ^a	ee (%) ^b
1	InCl ₃	—	48 h	—	—
2	InBr ₃	—	48 h	—	—
3	InBr ₃	2,2'-Bipyridyl	48 h	—	—
4	InCl ₃	1a	6 d	60	95
5	InBr ₃	1a	16 h	80	92
6	InBr ₃	1b	6 d	—	—

^a Isolated yields of chromatographed material. ^b Determined from chiral HPLC analysis (see supplementary information).

Table 2 Indium–bipyridine-catalyzed ring opening of *meso*-epoxides with thiols^a

Entry	Ar	R	Yield (%) ^b	ee (%) ^{c,d}
1	C ₆ H ₅	C ₆ H ₅ CH ₂	80 (4a)	92
2	C ₆ H ₅	C ₆ H ₅	81 (4b)	96
3	C ₆ H ₅	4-Me-C ₆ H ₄	79 (4c)	96
4	C ₆ H ₅	CH ₃ CH ₂	91 (4d)	>95
5	C ₆ H ₅	n-Bu	90 (4e)	95
6	C ₆ H ₅	(CH ₃) ₂ CH(CH ₂) ₂	89 (4f)	96
7	2-Naphthyl	C ₆ H ₅	79 (4g)	96
8	2-Naphthyl	C ₆ H ₅ CH ₂	82 (4h)	96
9	3-Me-C ₆ H ₄	C ₆ H ₅	67 (4i)	91
10	4-Me-C ₆ H ₄	C ₆ H ₅	76 (4j)	85
11	4-Me-C ₆ H ₄	C ₆ H ₅ CH ₂	68 (4k)	92
12	4-Cl-C ₆ H ₄	C ₆ H ₅	84 (4l) ^e	92
13	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	67 (4m) ^e	93


^a Reaction conditions: 10 mol% InBr₃, 11 mol% bipyridine **1a**, 1.5 equiv. thiol, CH₂Cl₂, rt, 16 h. ^b Isolated yields of chromatographed material. ^c Determined from chiral HPLC analysis (see supplementary information). ^d The absolute configuration of **4b** was proven to be 1*R*,2*R* through conversion into the corresponding known sulfone;¹¹ the configurations of all other products were assigned by analogy. ^e Reaction time: 6 d.

enantioselectivities (entries 7–13). Unfortunately, aliphatic *meso*-epoxides currently give rise to only low enantioselectivity in the indium–bipyridine-catalyzed thiolysis.

The product 1,2-mercapto alcohols **4** may be employed as substrates for desulfurization reactions, furnishing chiral, highly enantiomerically enriched alcohols. Thus, nickel boride reduction¹² of **4a** afforded alcohol **5** in good yield and almost identical enantiomeric excess as **4a**, proving that the reaction had not affected the stereochemistry of the adjacent carbinol center (Scheme 1).

Accordingly, the sequential thiolysis–desulfurization of *meso*-epoxides constitutes an elegant alternative to the still elusive enantioselective hydride addition to epoxides.¹³

A single crystal of the indium(III)–bipyridine complex suitable for X-ray crystallography was obtained from tetrahydrofuran–H₂O solution.¹⁴ The cationic [InBr₂·**1a**·H₂O]⁺ complex (the bromide counter anion is omitted for clarity) depicted in Fig. 2 is characterized with a pentagonal–bipyramidal coordination geometry around the indium(III) center, which is tetracoordinated by the bipyridine ligand **1a** through both nitrogen and oxygen atoms. Significantly, both hydroxyl protons are still attached to the catalyst. The crystal structure very closely resembles the structure of the scandium(III)–bipyridine catalyst,¹⁵ and both catalysts also exhibit the same sense of asymmetric induction in epoxide-opening reactions.⁸ Additional high resolution ESI-MS measurements gave



Scheme 1 Desulfurization of 1,2-mercapto alcohol **4a**.




Fig. 2 ORTEP (50% ellipsoid) of the [InBr₂·**1a**·H₂O]⁺ moiety in the X-ray crystal structure of [InBr₂·**1a**·H₂O]Br·THF·4H₂O.

a molecular peak at 602.9533 for [M]⁺ which further corroborated the molecular formula InBr₂C₂₀H₂₈N₂O₂.

Considering the failure of any indium(III)–bipyridine catalyst lacking the additional hydroxyl groups, it is most likely that the chiral catalyst exhibits a combined Lewis acid–Brønsted acid activity, with the Lewis acidic indium center activating the epoxide and the hydroxyl protons possibly forming a hydrogen bond to the incoming thiol, thereby guiding it to the epoxide.

In conclusion, we have devised a novel chiral catalyst for the highly enantioselective thiolysis of aromatic *meso*-epoxides, with aromatic as well as aliphatic thiols furnishing 1,2-mercapto alcohols in good yields and excellent enantioselectivities. Most importantly, there is no restriction to a specific thiol. A crystal structure analysis of the chiral catalyst reveals a pentagonal–bipyramidal coordination geometry around the indium center with both nitrogen and oxygen atoms coordinating to the metal. Additional hydrogen-bonding between the catalyst and the incoming nucleophile appears to activate and direct the incoming nucleophile to the indium-bound epoxide.

Generous financial support was provided by the Deutsche Forschungsgemeinschaft (Schn 441/3-2). We thank Wacker AG for providing chemicals and Dr Claudia Birkemeyer (University of Leipzig) for recording the ESI-MS spectra of the indium–bipyridine complex.

Notes and references

- Reviews: (a) C. Schneider, *Synthesis*, 2006, 3919; (b) I. M. Pastor and M. Yus, *Curr. Org. Chem.*, 2005, **9**, 1; (c) E. N. Jacobsen and M. H. Wu, in *Comprehensive Asymmetric Catalysis*, ed. E. N. Jacobsen, A. Pfaltz and H. Yamamoto, Springer, Berlin, 1999, vol. 3, p. 1309.
- (a) X. L. Hou, J. Wu, L. X. Dai, L. J. Xia and M. H. Tang, *Tetrahedron: Asymmetry*, 1998, **9**, 1747; (b) S. Sagawa, H. Abe, Y. Hase and T. Inaba, *J. Org. Chem.*, 1999, **64**, 4962; (c) A. Sekine, T. Ohshima and M. Shibasaki, *Tetrahedron*, 2002, **58**, 75; (d) G. Bartoli, M. Basco, A. Carlone, M. Locatelli, M. Massaccesi, P. Melchiorre and L. Sambri, *Org. Lett.*, 2004, **6**, 2173; (e) F. Carree, R. Gil and J. Collin, *Tetrahedron Lett.*, 2004, **45**, 7749; (f) F. Carree, R. Gil and J. Collin, *Org. Lett.*, 2005, **7**, 1023.
- (a) W. M. Nugent, *J. Am. Chem. Soc.*, 1992, **114**, 2768; (b) L. E. Martinez, J. L. Leighton, D. H. Carsten and E. N. Jacobsen, *J. Am. Chem. Soc.*, 1995, **117**, 5897.
- (a) E. N. Jacobsen, F. Kakiuchi, R. G. Konsler, J. F. Larrow and M. Tokunaga, *Tetrahedron Lett.*, 1997, **38**, 773; (b) S. Matsunaga, J. Das, J. Roels, E. M. Vogl, N. Yamamoto, T. Iida, K. Yamaguchi and M. Shibasaki, *J. Am. Chem. Soc.*, 2000, **122**, 2252.
- (a) B. M. Cole, K. D. Shimizu, C. A. Krueger, J. P. A. Harrity, M. L. Snapper and A. H. Hoveyda, *Angew. Chem.*, 1996, **108**, 1776, (*Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1668); (b) K. D. Shimizu,

- B. M. Cole, C. A. Krueger, K. W. Kuntz, M. L. Snapper and A. H. Hoveyda, *Angew. Chem.*, 1997, **109**, 1782, (*Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1703); (c) S. E. Schaus and E. N. Jacobsen, *Org. Lett.*, 2000, **2**, 1001.
- 6 (a) S. E. Denmark, P. A. Barsanti, K. T. Wong and R. A. Stavenger, *J. Org. Chem.*, 1998, **63**, 2428; (b) W. A. Nugent, *J. Am. Chem. Soc.*, 1998, **120**, 7139; (c) B. Tao, M. M. C. Lo and G. C. Fu, *J. Am. Chem. Soc.*, 2001, **123**, 353; (d) M. Nakajima, M. Saito, M. Uemura and S. Hashimoto, *Tetrahedron Lett.*, 2002, **43**, 8827; (e) E. Tokuoka, S. Kotani, H. Matsunaga, T. Ishizuka, S. Hashimoto and M. Nakajima, *Tetrahedron: Asymmetry*, 2005, **16**, 2391.
- 7 (a) T. Iida, N. Yamamoto, H. Sasai and M. Shibasaki, *J. Am. Chem. Soc.*, 1997, **119**, 4783; (b) M. H. Wu and E. N. Jacobsen, *J. Org. Chem.*, 1998, **63**, 5252; (c) J. Wu, X.-L. Hou, L.-X. Dai, L.-J. Xia and M.-H. Tang, *Tetrahedron: Asymmetry*, 1998, **9**, 3431.
- 8 (a) C. Schneider, A. R. Sreekanth and E. Mai, *Angew. Chem.*, 2004, **116**, 5809, (*Angew. Chem., Int. Ed.*, 2004, **43**, 5691); (b) for the first synthesis and application of bipyridine **1a** in asymmetric catalysis, see: C. Bolm, M. Zehnder and D. Bur, *Angew. Chem.*, 1990, **102**, 206, (*Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 191); (c) E. Mai and C. Schneider, *Chem.-Eur. J.*, 2007, **13**, 2729; (d) A. Tschöp, A. Marx, A. R. Sreekanth and C. Schneider, *Eur. J. Org. Chem.*, 2007, 2318; (e) for related work, see: S. Azoulay, K. Manabe and S. Kobayashi, *Org. Lett.*, 2005, **7**, 4593; (f) M. Boudou, C. Ogawa and S. Kobayashi, *Adv. Synth. Catal.*, 2006, **348**, 2585.
- 9 For non-enantioselective indium(III)-catalyzed thiolyses of epoxides, see: (a) J. S. Yadav, B. V. S. Reddy and G. Baishya, *Chem. Lett.*, 2002, **31**, 906; (b) F. Fringuelli, F. Pizzo, S. Tortoioli and L. Vaccaro, *Adv. Synth. Catal.*, 2002, **344**, 379; (c) F. Fringuelli, F. Pizzo, S. Tortoioli and L. Vaccaro, *Org. Lett.*, 2005, **7**, 4411.
- 10 After the work described herein had been carried out and while the manuscript was being prepared, a related report about a scandium-bipyridine-catalyzed process using mainly thiophenol appeared, see: C. Ogawa, N. Wang and S. Kobayashi, *Chem. Lett.*, 2007, **36**, 34.
- 11 S. Nakamura, R. Nakagawa, Y. Watanabe and T. Toru, *J. Am. Chem. Soc.*, 2000, **122**, 11340.
- 12 T. G. Back, D. L. Barton and K. Yang, *J. Org. Chem.*, 1993, **58**, 2407.
- 13 For a mechanistically different approach towards formal enantioselective hydride addition to *meso*-epoxides, see: A. Gansäuer, T. Lauterbach, H. Bluhm and M. Noltemeyer, *Angew. Chem.*, 1999, **111**, 3112, (*Angew. Chem., Int. Ed.*, 1999, **38**, 2909); A. Gansäuer, H. Bluhm, B. Rinker, S. Narayan, M. Schick, T. Lauterbach and M. Pierobon, *Chem.-Eur. J.*, 2003, **9**, 531.
- 14 Crystallographic data for the complex: single crystal from THF-H₂O, C₂₄H₄₆Br₃InN₂O₈, *M* = 845.18, orthorhombic, *a* = 8.3069(3), *b* = 17.0302(6), *c* = 23.6466(9) Å, *U* = 3345.2(2) Å³, *T* = 180(2) K, space group *P*2₁2₁2₁ (No. 19), *Z* = 4, μ (Mo-K α) = 4.333 mm⁻¹, *D*_c = 1.678 g cm⁻³, 26 247 reflections measured, 7293 unique (*R*_{int} = 0.0413), which were used in all calculations. The final *R*1 = 0.0407 and *wR*2 = 0.0892 (all data). G. M. Sheldrick, *SHELXS-97*, *SHELXL-97. Programs for crystal structure determination*, University of Göttingen, Germany, 1997. CCDC 640641. For crystallographic data in CIF or other electronic format, see DOI: 10.1039/b703625f.
- 15 S. Ishikawa, T. Hamada, K. Manabe and S. Kobayashi, *J. Am. Chem. Soc.*, 2004, **126**, 12236.