

Gold-catalysed intramolecular *trans*-allylsilylation of alkynes forming 3-allyl-1-silaindenes†

Takanori Matsuda, Sho Kadowaki, Yoshiyuki Yamaguchi and Masahiro Murakami*


Received (in Cambridge, UK) 25th March 2008, Accepted 23rd April 2008

First published as an Advance Article on the web 20th May 2008

DOI: 10.1039/b804721a

3-Allyl-1-silaindenes are synthesised from alkynes having an allylsilane moiety by a gold-catalysed intramolecular *trans*-allylsilylation reaction.

Enyne cycloisomerisation reactions catalysed by electrophilic transition metal complexes have emerged as a powerful and efficient method for the construction of complex cyclic structures.¹ Yamamoto and his co-workers reported an interesting example using an alkyne tethered to an allylsilane moiety.² The intramolecular *trans*-allylsilylation reaction is catalysed by a hafnium complex to afford carbocyclic compounds (eqn (1)). Our interest in the synthesis of silole (silacyclopentadiene) derivatives, which possess unique photo-physical and electronic properties,³ led us to examine the reaction of enynes linked by a silicon-containing tether (eqn (2)). Herein, we report a gold(I)-catalysed intramolecular *trans*-allylsilylation reaction of silicon-tethered enynes furnishing silole cores.^{4–6}




Gold complexes possess a highly electrophilic character which often works specifically on alkynes.⁷ Thus, we examined the use of a gold(I) complex to catalyse a reaction of allyldimethyl[2-(pent-1-ynyl)phenyl]silane (**1a**) (Scheme 1).⁸ Whereas a neutral chlorogold(I) complex was ineffective, cationic gold complexes having a bistriflimide counter anion⁹ catalysed an intramolecular *trans*-allylsilylation reaction at room temperature to produce 3-allyl-1-silaindene **2a**.¹⁰ The yield of **2a** was significantly affected by the ligand. 2-Di-*tert*-butylphosphinobiphenyl was found to be the ligand of choice,

giving **2a** in 97% yield.¹¹† A gold complex bearing IPr¹² ligand exhibited moderate catalytic activity for the reaction.

The reaction of various sila-enynes **1** having a range of the alkyne substituents (R) was examined using **3** as the catalyst, and the results are listed in Table 1. Methyl, *sec*-alkyl, methoxycarbonyl, alkynyl, and aryl groups were all suitable substituents to give the corresponding 2-substituted 3-allyl-1-silaindenes **1b–j** in moderate to good yield (entries 1–9). Higher catalyst loadings (6–12 mol%) were required to complete the reaction in the case of the cyclopentyl and aryl groups. Notably, a variety of functional groups such as a boryl group and a thienyl group were tolerated in the gold-catalysed reaction. Chloro- and bromoalkyne moieties could be contained in the substituents, producing 2-halo-1-silaindenes **2k** and **2l**, respectively (entries 10 and 11).¹³ On the other hand, the substrates possessing a hydrogen (*i.e.*, a terminal alkyne), a bulky *tert*-butyl group, and 2-pyridyl group failed to afford the corresponding 1-silaindenes.

Next, structural variation in the allylsilane moiety was investigated (Table 2). Whereas methylphenylsilylene-bridged sila-enyne **1m** provided 1-silaindene **2m** in good yield (entry 1), diethylsilylene-bridged substrate **2n** was less reactive (entry 2). Methallyl-, crotyl-, and cyclohexenylsilanes **1o–q** also gave the corresponding 1-silaindenes **2o–q** (entries 3–5). The result with **1p** indicated that the allylation took place regioselectively at the γ position. Reaction of allylgermane **1r** occurred in a similar manner to afford 3-allyl-1-germaindene **2r** in 85% yield (entry 6).


The synthesis of 1-silaindene derivatives possessing extended π -conjugation is exemplified in eqn (3) and (4). *m*-Phenylenebis(1-silaindene) **5a** was obtained by two-fold *trans*-allylsilylation of **4a**. An analogous two-fold allylsilylation successfully proceeded with substrate **4b** to afford 1,5-disila-*s*-indacene **5b**.



Scheme 1 Gold-catalysed *trans*-allylsilylation of sila-enyne **1a**.



Department of Synthetic Chemistry and Biological Chemistry, Kyoto University, Katsura, Kyoto 615-8510, Japan. E-mail: murakami@sbchem.kyoto-u.ac.jp; Fax: +81 (75) 383-2748

† Electronic supplementary information (ESI) available: Experimental procedures and characterisation data for new compounds. See DOI: 10.1039/b804721a

Table 1 Gold-catalysed *trans*-allylsilylation of sila-enynes **1b–l**^a

Entry	1 (R)	mol% Au	2	Yield ^b
1	1b (Me)	3	2b	58%
2	1c (cyclopentyl)	6	2c	71% ^c
3	1d (CO ₂ Me)	3	2d	89%
4	1e (C≡CTIPS ^d)	3	2e	84%
5	1f (4-MeC ₆ H ₄)	6	2f	75% ^e
6	1g (4-(pin)BC ₆ H ₄ ^f)	6	2g	81%
7	1h (3-MeOC ₆ H ₄)	6	2h	71%
8	1i (2-FC ₆ H ₄)	12	2i	77%
9	1j (5-methyl-2-thienyl)	6	2j	42%
10 ^g	1k (Cl)	3	2k	79%
11 ^g	1l (Br)	3	2l	77%

^a Sila-enynes **1b–l** (0.20 mmol) were reacted in the presence of the catalyst **3** (3–12 mol%) in CH₂Cl₂ (2.5 mL) at room temperature for 9 h. ^b Isolated yield by preparative TLC. ^c 33% yield with 3 mol% of **3**. ^d TIPS = triisopropylsilyl. ^e 13% conversion after 1 h. ^f Pin = pinacolato (–OCMe₂CMe₂O–). ^g Reaction time: 2 h.




A mechanism conceivable for the production of **2** from **1** is shown in Scheme 2.¹⁴ Initially, the alkyne moiety of **1** activated by electrophilic Au⁺ (in **A**) induces intramolecular addition of the pendant allyl group to produce seven-membered ring intermediate **B**. Subsequent ring opening accomplishes allylation to give alkenylgold species **C**, to which the cationic silyl moiety undergoes electrophilic ring closure. Finally, demetalation from alkygold species **D** furnishes sila-cyclo **2** and regenerates Au⁺.¹⁵

The produced 3-allyl-1-silaindenes **2** exhibited fluorescence upon ultraviolet excitation with 250 nm in hexane solution. Silaindene **2a** emitted fluorescence at 361 nm with quantum efficiency of 0.07, and **5b** at 408 nm with 0.33.

Table 2 Reaction of **1m–r** in the presence of **3**^a

Entry	1	2	Yield ^b
1	1m	2m	83%
2 ^c	1n	2n	53%
3	1o	2o	83%
4	1p ^d	2p	69%
5	1q	2q	46%
6	1r	2r	85%

^a Unless otherwise noted, all reactions were carried out in the presence of 3 mol% of **3** in CH₂Cl₂ at room temperature for 9 h. ^b Isolated yield by preparative TLC. ^c 6 mol% of **3** was used. ^d *cis* : *trans* = 93 : 7.

**Scheme 2** Mechanism of gold-catalysed intramolecular *trans*-allylsilylation.

In summary, we have developed a new method for the synthesis of 3-allyl-1-silaindenes by a gold-catalysed intramolecular *trans*-allylsilylation reaction of alkynes.


This work was supported by a Grant-in-Aid for Scientific Research for Young Scientist (B) (No. 19750074) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Notes and references


‡ General procedure: to a Schlenk tube containing [2-PhC₆H₄P(*t*-Bu)]₂AuNTf₂ (**3**, 7.0 mg, 9.0 μmol, 3 mol%) under an argon atmosphere was added a solution of allyldimethyl[2-(pent-1-ynyl)-phenyl]silane (**1a**, 72.7 mg, 0.30 mmol) in dichloromethane (0.37 mL). After stirring for 9 h at room temperature, the volatile materials were removed under reduced pressure. The residue was purified by preparative thin-layer chromatography (hexane) to give 3-allyl-1,1-dimethyl-2-propyl-1-silaindene (**2a**, 70.2 mg, 97%).

- 1 Reviews: (a) M. Méndez, V. Mamane and A. Fürstner, *Chemtracts*, 2003, **16**, 397; (b) A. M. Echavarren and C. Nevado, *Chem. Soc. Rev.*, 2004, **33**, 431; (c) C. Bruneau, *Angew. Chem., Int. Ed.*, 2005, **44**, 2328; (d) S. Ma, S. Yu and Z. Gu, *Angew. Chem., Int. Ed.*, 2006, **45**, 200; (e) C. Nieto-Oberhuber, S. López, E. Jiménez-Núñez and A. M. Echavarren, *Chem.–Eur. J.*, 2006, **12**, 5916; (f) L. Zhang, J. Sun and S. A. Kozmin, *Adv. Synth. Catal.*, 2006, **348**, 2271.
- 2 (a) K.-i. Imamura, E. Yoshikawa, V. Gevorgyan and Y. Yamamoto, *J. Am. Chem. Soc.*, 1998, **120**, 5339. For the intermolecular variants, see; (b) S. H. Yeon, J. S. Han, E. Hong, Y. Do and I. N. Jung, *J. Organomet. Chem.*, 1995, **499**, 159; (c) N. Asao, E. Yoshikawa and Y. Yamamoto, *J. Org. Chem.*, 1996, **61**, 4874; (d) E. Yoshikawa, V. Gevorgyan, N. Asao and Y. Yamamoto, *J. Am. Chem. Soc.*, 1997, **119**, 6781; (e) N. Asao, H. Tomeba and Y. Yamamoto, *Tetrahedron Lett.*, 2005, **46**, 27.
- 3 (a) T. Matsuda, S. Kadowaki, T. Goya and M. Murakami, *Org. Lett.*, 2007, **9**, 133; (b) T. Matsuda, S. Kadowaki and M. Murakami, *Chem. Commun.*, 2007, 2627; (c) T. Matsuda, Y. Yamaguchi and M. Murakami, *Synlett*, 2008, 561.
- 4 For recent examples of a gold-catalysed intramolecular *cis*-allylsilylation reaction of 4-sila-1,5-enynes, see: (a) S. Park and D. Lee, *J. Am. Chem. Soc.*, 2006, **128**, 10664; (b) Y. Horino, M. R. Luzung and F. D. Toste, *J. Am. Chem. Soc.*, 2006, **128**, 11364.
- 5 For related intramolecular allyl migration reactions, see: (a) A. Fürstner, F. Stelzer and H. Szillat, *J. Am. Chem. Soc.*, 2001, **123**,

- 11863; (b) A. Fürstner and P. W. Davies, *J. Am. Chem. Soc.*, 2005, **127**, 15024; (c) S. Cacchi, *J. Organomet. Chem.*, 1999, **576**, 42.
- 6 For cycloisomerisations of 1,7-enynes, see: (a) A. Ochida, H. Ito and M. Sawamura, *J. Am. Chem. Soc.*, 2006, **128**, 16486; (b) S. Porcel and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2007, **46**, 2672; (c) N. Cabello, C. Rodríguez and A. M. Echavarren, *Synlett*, 2007, 1753.
- 7 Reviews: (a) A. Hoffmann-Röder and N. Krause, *Org. Biomol. Chem.*, 2005, **3**, 387; (b) E. Jiménez-Núñez and A. M. Echavarren, *Chem. Commun.*, 2007, 333; (c) D. J. Gorin and F. D. Toste, *Nature*, 2007, **446**, 395; (d) A. Fürstner and P. W. Davies, *Angew. Chem., Int. Ed.*, 2007, **46**, 3410; (e) Y. Yamamoto, *J. Org. Chem.*, 2007, **72**, 7817; (f) A. S. K. Hashmi, *Chem. Rev.*, 2007, **107**, 3180.
- 8 A reaction in the presence of HfCl₄ (CH₂Cl₂, 0 °C) resulted in a complex mixture. No reaction occurred in the presence of PtCl₂ (toluene, 80 °C).
- 9 N. Mézailles, L. Ricard and F. Gagosz, *Org. Lett.*, 2005, **7**, 4133
- 10 Siloxane **6** was formed in 23% yield in addition to **2a** (61%) in the case of (Ph₃P)AuNTf₂



- 11 Unlike the previous Al- and Hf-catalysed *trans*-allylsilylation reaction of alkynes, the complex **3** failed to promote an intermolecular reaction of allyldimethyl(phenyl)silane with pent-1-ynylbenzene.
- 12 IPr = Bis(2,6-diisopropylphenyl)imidazol-2-ylidene.
- 13 Dehalogenated 1-silaindene **7** (ca. 14%) was also isolated as the side product. Performing the reaction for a longer period of time than 2 h increased the formation of **7**



- 14 It is premature for us to argue about the mechanism in more detail. A different mechanism can be also assumed to operate depending on the substrates employed.
- 15 For a related silyldeauration process, see: I. Nakamura, T. Sato, M. Terada and Y. Yamamoto, *Org. Lett.*, 2007, **9**, 4081.